QK
167
B3
1904

B 4 314 894

EXCHANGE



BIOLOGY
LIBRARY
G

EXCHANGE
NOV 8 1914

MICHIGAN FLORA

A LIST OF THE

FERN AND SEED PLANTS GROWING WITHOUT CULTIVATION

PREPARED BY

W. J. BEAL, Sc. M., Ph. D.,

AGRICULTURAL COLLEGE, MICHIGAN.

[Reprinted by permission, from the Fifth Report of the Michigan Academy of Science, 1904.]

PUBLISHED BY
THE STATE BOARD OF AGRICULTURE
AGRICULTURAL COLLEGE, MICH.

LANSING, MICH.
ROBERT SMITH PRINTING CO., STATE PRINTERS AND BINDERS
1904

MICHIGAN FLORA

A LIST OF THE

FERN AND SEED PLANTS GROWING WITHOUT CULTIVATION

PREPARED BY

W. J. BEAL, Sc. M., Ph. D.,

AGRICULTURAL COLLEGE, MICHIGAN.

[Reprinted by permission, from the Fifth Report of the Michigan Academy of Science, 1904.]

PUBLISHED BY
THE STATE BOARD OF AGRICULTURE
AGRICULTURAL COLLEGE, MICH.

LANSING, MICH.
ROBERT SMITH PRINTING CO., STATE PRINTERS AND BINDERS
1904

QR161
B3
1904

BIOLOGY
LIBRARY
G

UNIV. OF
CALIFORNIA

INTRODUCTION.

The first Michigan Flora entitled a "Catalogue of the Phænogamous and Vascular Cryptogamous Plants of Michigan, Indigenous, Naturalized and Adentive," was prepared by Charles F. Wheeler and Erwin F. Smith, and was printed in the report of the Michigan State Horticultural Society for 1880.

The second Michigan Flora, based on the first, was prepared by W. J. Beal and C. F. Wheeler and was printed in the report of the Michigan State Board of Agriculture for 1892. Of the second Michigan Flora one thousand separates were printed for distribution and the supply was exhausted in less than five years.

The copies of all the former edition of this Flora were distributed chiefly among botanists of all grades from Professors in Universities and Colleges, Normal Schools, teachers in High Schools, Academies and among amateurs. It is believed that this edition notwithstanding all its defects, will encourage many to study the wild plants of Michigan not forgetting the arrivals from other countries. Besides assisting the student in becoming familiar with names of species and their distribution, it should be useful in other respects.

Within the past few years a delightful department of botany has attracted much attention. It is emphatically outdoor work and is known as Ecology or the relations of plants to their environment.

This Flora may help the student in his investigations of plant groups or plant associations, noting those peculiar to certain kinds of soil, and others growing almost everywhere. It will aid in listing natives and exotics. Students can make many lists, such as those forming rosettes, those that climb, those that thrive in the woods in early spring, those that grow in strata or layers above each other or in zones within and without each other. It should aid in the study of plant dispersal by all sorts of methods and in the multitude of ways by which plants protect themselves.

The following is a list of the chapters found in the last edition of the Flora that are omitted in this edition:

Planting the Roadside and about the Home.
Planting a Grove.
Planting a Wild Garden.
Autumn Foliage.
Native Trees and Shrubs selected for the Color of their Leaves in Autumn.
Native small Trees and Shrubs distinguished for their Flowers.
Native Shrubs or Trees distinguished for their beautiful Fruit.
A list of native Trees and Shrubs distinguished for their showy or brilliant colored Bark.

293907

Native Climbing Plants.
Native Plants which are very light Colored.
A list of Small Evergreens.
Bronze Evergreens.
Native Bog and Marsh plants which are Promising for Cultivation.
Plants suitable for winter Bouquets.
Native Aquatic Plants most worthy of Cultivation.
A list of native Ferns promising for Cultivation.
The Procession of Flowers.
The best Kinds of Timber for Firewood.
The most durable Kinds of Timber for Post or Sills.
Favorite Kinds of Timber for the Cabinet Maker.
Timber for farm Implements.
Valuable Timber for large Boats.
Kinds of timber most employed for Boxes, Baskets and Barrels.
Timber for Paper Pulp.
Trees for Sugar.
Trees and Shrubs best suited for Screens or Wind-breaks.
Some of our wild fruits and nuts, with suggestions concerning their improvement.
Native plants for the Protection of Hillsides, Embankments and drifting Sands.
Plants for Carp Ponds.
Native tree-like or large Shrubs.
List of smaller Shrubs.
List of Rare or Local Plants.
List of Plants introduced from Europe and becoming Naturalized.
Parasitic Fungi.
Flowering Parasites and Saprophytes destitute of green leaves.
List of Michigan Plants which are admitted to the United States Pharmacopœia.
List of medicinal Plants not officinal.

As mentioned in both editions of the Flora above referred to, so in this, the third edition, much pains has been taken to examine authentic specimens which are preserved in some herbarium. For thirty years, till he resigned his position at the Agricultural College in 1902, Professor C. F. Wheeler was a most diligent collector of plants in the State. He kept full notes, not only of his findings but sought specimens and notes from many others.

In 1888, Professor L. H. Bailey, C. F. Wheeler, and the author accompanied by two students spent two weeks in collecting on a journey across the State from Harrisville in Alcona county to Frankfort in Benzie county.

In 1892, Professor Wheeler spent several weeks in the southeastern and in the southwestern counties and in the Upper Peninsula making collections for the exposition held in Chicago. In 1895, some weeks were spent collecting in Alpena and vicinity, and later he and B. O. Longyear collected in Ingham, Washtenaw, and Jackson counties. In the summer of 1900 Professor Wheeler spent about six weeks collecting at Chatham and vicinity in the Upper Peninsula.

Portions of every year since 1890 were occupied by Professor Wheeler in making great numbers of short excursions in behalf of the herbarium of the Agricultural College. During these years and previously, the author collected considerable in the counties of Iosco, Clare, Crawford, Grand Traverse, Lake, Muskegon, Calhoun, Eaton, Lenawee, Clinton, Ingham. More particularly since 1890, numerous collections made by others have been sent to the Agricultural College for identification and for preservation in the herbarium.

Notably among these collectors must be mentioned Miss Emma J. Cole

and H. C. Skeels of Grand Rapids; C. D. McLouth of Muskegon; G. H. Hicks of Grayling, Owosso and the Agricultural College; C. K. Dodge of Port Huron; George M. Bradford of Bay City; W. K. Brotherton of Oakland; Geo. W. Davis of Tekonsha; J. W. Stacy of Clarksville. Collections have been received, by gift or purchase, from Prof. C. A. Davis formerly of Alma, now of the University; of O. A. Farwell formerly of Keweenaw county, of Ypsilanti, and later of Detroit; and Rev. Francis Daniels formerly of Alto, Kent county, Manistee and Sturgis; W. S. Cooper of Alma.

After all has been said and done, the study of the flora of the state at best can only be considered as fairly begun. By far the greater areas have not yet been seen by any systematic botanist and very few regions have been visited by one who is an expert in some one or more of the more difficult families.

What species flourished in large areas will never be fully known, since man has cut off, burned over and plowed under tens of thousands of acres of the virgin wilderness! Swamps, marshes and lakes have been drained and the land occupied by farm crops. Many native plants are rapidly shifting from one place to another.

Chiefly through the agency of man, great numbers of weeds and other plants have been introduced from other states and from foreign countries and each has begun a vigorous warfare for all the room it can get.

The sequence of natural families in former Michigan Floras followed *Gray's Manual* which is essentially that of Auguste Pyrame De Candolle. Most of the reasons given for that arrangement have long since been considered untenable. In the Flora, I have followed *Britton's Manual of the Flora of the Northern States and Canada,* published in April, 1901. In this work the sequence of families is very nearly the same as that of Engler and Prantl, which is considered the most philosophical yet presented.

Some of the guiding principles for the system of Engler and Prantl, as they are stated by Britton and Brown's Flora, are as follows:

The more simple forms are, in general, distinguished from the more complex, (1) by fewer organs or parts; (2) by the less perfect adaptation of the organs to the purposes they subserve; (3) by the relative degree of development of the more important organs; (4) by the lesser degree of differentiation of the plant-body or of its organs; (5) by considerations of antiquity, as indicated by the geological record; (6) by a consideration of the phenomena of embryogeny. Thus, the Pteridophyta, which do not produce seeds and which appear on the earth in Silurian time, are simpler than the Spermatophyta; the Gymnospermæ in which the ovules are borne on the face of a scale, and which are known from the Devonian period onward, are simpler than the Angiospermæ, whose ovules are borne in a closed cavity, and which are unknown before the Jurassic.

In the Angiospermæ the similar types are those whose floral structure is nearest the structure of the branch or stem from which the flower has been metamorphosed, that is to say, in which the parts of the flower (modified leaves) are more nearly separate or distinct from each other, the leaves of any stem or branch being normally separated, while those are the most complex whose floral parts are most united.

The names of genera and species are the same as those used in *Britton's Manual* and where these differ from those in the sixth edition of

Gray's Manual, the latter are also inserted in the text. The species of each genus are arranged in alphabetical order. To economize space a considerable number of sections of the Flora last prepared have been omitted or much abbreviated.

In the preparation of this Flora thanks are due to Professor C. F. Wheeler, O. A. Farwell, Prof. C. A. Davis, C. K. Dodge, C. D. McLouth, G. M. Bradford, J. B. Dandeno.

W. J. BEAL.

Agricultural College, Mich., Dec. 1, 1904.

CONTENTS.

	Page.
Bibliography	9
Herbaria consulted	10
Local list consulted	11
Topography	11
Climate and distribution	13
Flora of the Jack Pine Plains	16
a. Those most common	16
b. Those less frequent	16
Plants indicating a fertile soil	18
Plants peculiar to the prairies	19
Overlapping of northern and southern species in the Grand River Valley	19
Northern species	19
Southern species	19
Comparison of the flora of the eastern and the western sides of the State in latitude 44° 40′	20
a. Northern plants found on the east side of the State and not on the west	20
b. Southern plants found on the west side of the State and not on the east	20
Plants supposed to have immigrated from the northeast	20
Plants supposed to have immigrated from the north and west	21
Trees of Michigan compared with those of Europe	21
Why has Michigan so many trees and Great Britain so few?	22
Native forage plants	23
Native bee plants	24
Weeds, native and introduced	25
List of weeds introduced from Europe and Asia	25
List of indigenous weeds	27
Native poisonous plants	72
Native plants fast disappearing	28
List of trees indigenous to Michigan	28
List of shrubs indigenous to Michigan	30
Flora catalogue	34
Index	137

BIBLIOGRAPHY.

The following publications have been consulted in the preparation of this work:

1839. Wright, John, M. D., Catalogue of the Phænogams and Filicoid Plants collected on the Geological Survey of Michigan. Legislative Report No. 23, pp. 17-44. Detroit.

1849. Burt, W. A., Catalogue of the Plants collected in the primitive region south of Lake Superior in 1846. D. Cooley, Jackson's Lake Superior, pp. 875-882. Washington, D. C.

1850. Agassiz, Louis, Lake Superior, its Physical Character, Vegetation and Animals, etc.

1851. Whitney, W. D., List of Plants of the Upper Peninsula in Report on the Geology of the Lake Superior Land District. J. W. Foster and J. D. Whitney, part 2.

1853. Cooley, D., M. D., A Manuscript List of the Plants growing spontaneously within ten miles of Cooley's corners, Washington, Macomb, Co.

1861. Winchell, N. H., Catalogue of Phænogamous and Acrogenous Plants found growing wild in the Lower Peninsula of Michigan and the Islands at the head of Lake Huron. Geological Report for 1860, pp. 245-330.

1873. Coleman, N., Catalogue of Flowering Plants of the Southern Peninsula of Michigan, with a few of the Cryptogamia. Grand Rapids, Miscellaneous Publications, No. 2, Kent Scientific Institute.

1876. Tuthill, F. H., Some notes on the Flora near Kalamazoo, Mich. Bot. Gaz. Vol. 1, pp. 13-14.

1876. Almendinger, E. C., Flora of Ann Arbor and Vicinity. Proceedings of the Ann Arbor Scientific Association, pp. 85-116.

1877. Palmer, Elmore, M. D., Catalogue of Phænogamous and Acrogenous Plants found growing wild in the State of Michigan.

1877. Spalding, V. M., List of Native Medicinal Plants of Michigan. Proceedings of the Michigan Pharmaceutical Association.

1878. Lyons, A. B., M. D., Medicinal Plants Indigenous in Michigan. Nov. 27, 1877, Detroit Lancet, February and March, 1878.

1880. Bailey, L. H., Jr., Michigan Lake Shore Plants at South Haven. Bot. Gaz. pp. 76-77 and pp. 90-91.

1882. Bailey, L. H., Jr., Limits of Michigan Plants. Bot. Gaz. pp. 106-108.

1882. Foerste, A. F., Plants of Belle Isle, Michigan. Bot. Gaz. pp. 202-203.

1884. Hill, E. J., The Menominee Iron Region and its Flora. I., pp. 208-211; II., pp. 225-229; Bot. Gaz.

1886. Campbell, D. H., Plants of the Detroit River. Bull. Torr. Bot. Club, Vol. XIII, pp. 93-94.

1888. Beal, W. J., List of Trees and Shrubs belonging to Michigan. First Report of State Forestry Commission, pp. 36-51.

1888. Beal, W. J., Flora of the Sandy Pine Plains of Michigan. Report Mich. Hort. Soc. pp. 49-55.

1890. Gray, Asa, Manual of the Botany of the Northern United States. 6th Ed. pp. 760. American Book Co., N. Y.

1890. Hill, E. J., Notes on the Flora of the Lake Superior Region. I., pp. 140-149; II., pp. 159-166; Bot. Gaz.

1890. Bailey, L. H., Jr., The Carices of the Upper Half of the Keweenaw Peninsula. Bull. Torr. Bot. Club, pp. 61-64, Vol. 17.
1890. Macoun, John, M. A., Catalogue of Canadian Plants. S. C., 1883-1890.
1891. Wheeler, C. F., Central Michigan Cyperaceae. Bull. Torr. Bot. Club, p. 148. Vol. 18.
1891. Beal, W. J., and Wheeler, C. F., Michigan Flora. Report of the Secretary of the State Board of Agriculture, pp. 471-689.
1893. Blodgett, H. T., Plants of Mason County, Mich. Asa Gray Bull. No. 3.
1893. Hicks, Gilbert H., New and Rare Michigan Plants. Asa Gray Bull. No. 3.
1894. Farwell, O. A., Contributions to the Botany of Michigan. Asa Gray Bull. Nos. 6, 7, et seq.
1894. Pieters, A. J., Plants of Lake St. Clair. Bull. Mich. Fish Commission, No. 2, 1893.
1894. Reighard, J. E., Biological Examinations of Lake St. Clair. Bull. Mich. Fish Commission No. 4.
1894. Davis, Charles A., The Flora of Michigan Lakes. First Report of the Mich. Acad. Sci. pp. 24-31.
1896. Cole, Miss Emma J., List of Plants collected in and about Grand Rapids.
1896-8. Britton, N. L., and Brown, Addison, An Illustrated Flora of the Northern United States, Canada and the British Possessions. In three Volumes, Vol. I. pp. 612, Vol. II. pp. 643, Vol III. pp. 588. Charles Scribner's Sons, N. Y.
1897. Dodge, C. K., Flora of St. Clair County, Michigan and the Western part of Lambton County, Ontario. Report of the Secretary of the Mich. State Hort. Soc. pp. 230-314.
1898. Davis, Charles A., Contributions to the Knowledge of the Flora of Tuscola County. First Report of the Mich. Acad. Sci. p. 116, 1898. Bot. Gaz., June 1898, pp. 453-58.
1900. Davis, Charles A., Botanical Notes on Huron County. Michigan Geological Survey Reports, VII. pp. 235-245.
1900. Farwell, O. A., A catalogue of the Flora of Detroit with additions, from the Eleventh Annual Report of the Commissioner of Parks and Boulevards, Detroit.
1901. Britton, N. L., Manual of the Flora of the Northern States and Canada. pp. 1080. Henry Holt & Co., N. Y.
1901. Livingston, Burton E., The Distribution of the Plant Societies of Kent County. Michigan Geological Survey, Third Annual Report, pp. 81-103.
1902. Clark, H. L., Notes on the Flora of Eaton County. Third Report of the Mich. Acad. Sci. pp. 51-52.
1903. Livingston, Burton E., The Relation of Soils to Natural Vegetation in Roscommon and Crawford Counties. Mich. Geol. Survey, Annual Report.
1903. Daniels, Francis Potter, Flora of the Vicinity of Manistee, Mich. Fourth Report of the Mich. Acad. Sci. pp. 125-144.
1903. Daniels, Francis Potter, Ecology of the Flora of Sturgis and vicinity, Mich. Fourth Report of the Mich. Acad. Sci. pp. 145-159.

HERBARIA CONSULTED.

The following Herbaria have been examined:

The Herbarium of the State Agricultural College is fortunate in possessing the collection of Dr. D. Cooley, an excellent botanist who lived many years in Washington, Macomb county. He was a valued correspondent of Dr. Gray, Dr. Torrey, W. S. Sullivant and other botanists of the early part of this century.

The Herbarium of Dr. D. Clark, of Flint, Mich., has lately become the property of the State Agricultural College. This collection contains sets of Bebbs' Willows, Olney's Carices and many specimens from the earlier American collectors, besides valuable collections of Michigan plants.

The large collection of Prof. C. F. Wheeler, which was destroyed by the burning of the Botanical Laboratory on the 23d of March, 1890.

Prof. V. M. Spalding kindly permitted us to examine the University Herbarium at Ann Arbor, in which are deposited the collections of Dr. Douglass Houghton, 1838; Miss Mary H. Clark, Miss E. C. Almendinger, Prof. M. W. Harrington, Prof. N. H. Winchell, Geo. L. Ames, M. D., F. E. Wood and others.

The collection of plants belonging to the Kent Scientific Institute at Grand Rapids under the charge of Mr. George D. Sones.

The collection of O. J. Stilwell, which belongs now to Prof. C. A. Davis of University of Michigan; also Prof. Davis' collection.

The collection of G. H. Hicks, of the Agricultural College, made in Northern and Central Michigan.

The collection of G. F. Comstock, made in Lenawee County, 1845-'50, now the property of Dr. W. J. Beal.

The collection of Dr. W. J. Beal, 1860-1870, now the property of the Michigan Agricultural College.

LOCAL LISTS CONSULTED.

To the following persons we are indebted for lists of the plants growing in their several localities:

Farwell, O. A., for full list of plants of the Keweenaw peninsula, Ypsilanti and Detroit, with copious notes and many specimens.

Beardslee, Prof. H. C., of the University School, Cleveland, O., and Kofoid, Prof. Chas. A., Leland Stanford Jr. University, for a very complete list of the plants of Cheboygan County, Mich., observed by them during the smmer of 1890, with full notes and many specimens.

Dodge, C. K., for a collection of the plants growing in the vicinity of Port Huron, with many notes on variation and distribution.

Dewey, L. H., for a list of the plants in the vicinity of Tecumseh, Mich.

Hull, Prof. W., for notes and specimens from Albion and vicinity.

Orth, S. P., for list of plants in the vicinity of Imlay City.

Mosely, E. H., for a list of plants observed near Union City.

Foerste, A. F., for a list of plants observed in the eastern part of St. Clair county.

Mitchell, Prof. I. N., for a list of plants collected in various parts of the State.

St. John, Prof. C. E., for a list of plants collected in Mason county and in the southeastern portion of the State.

Stacey, I. W., for a list of plants collected at Clarksville, Ionia county.

Daniels, F. P., for a list of plants collected at Manistee and Sturgis.

Cooper, W. S.. St. Clair county.

Davis, Charles A., lists of plants of Gratiot county in the vicinity of Alma, and about Ann Arbor.

Bradford, G. M.. Flora of Bay county.

Pepoon, H. S., catalogue of the plants of Michigan adjacent to Lakes Magician, Dewey, Cable and Crooked, Van Buren and Cass counties.

TOPOGRAPHY.

Michigan is peculiarly situated within the waters of the great lakes. N. latitude 41°45′ to 48°20′; W. longitude 82°25′ to 90°34′. It is divided into two parts, called the Upper and Lower Peninsulas. The greatest length of the northern portion from east to west is 318 miles, width 30 to

164 miles, forming about two-fifths of the State. The greatest length of the southern portion from north to south is 277 miles and its extreme width is 259 miles. The total area is 58,915 square miles, with a coast of over 1,600 miles.

The general elevation of the Upper Peninsula is 400 to 1,100 feet above Lake Superior, and that of the Lower Peninsula is 400 to 600 feet above the level of Lakes Michigan and Huron.

The two parts of the State present a striking contrast in many respects. The Upper Peninsula may be divided into two sections east and west of a line drawn through Marquette which present very marked surface and geological characteristics. The eastern portion slopes northward from its southern border to a watershed and thence falls rapidly to the shores of Lake Superior. This plateau contains many lakes and marshes, also fine forests of pine intermixed with groves of hardwood.

The western part is rugged and hilly, some of the hills rising 1,000 to 1,200 feet. In the extreme northwest are ranges which form the copper region; the central range extends from Keweenaw Point across to the Wisconsin line; on either side are the Porcupine mountains and the copper range proper.

South and east of the copper range lies the iron range of Marquette and Iron counties. The eastern portion of this peninsula is underlaid with stratified rocks belonging to the Silurian period, while the western part is occupied by the copper bearing rocks and those of the Huronian period. Glacial drift covers deeply a large portion of both the eastern and western sections.

The Lower Peninsula is generally level or rolling, sloping up in its northern portion to a central ridge or watershed which extends nearly northeast and southwest, the highest part of which, in Otsego county, is 1,100 feet above the lake level.

The shores along the west side of this peninsula are generally bold bluffs which are constantly wearing away, while on the Huron shore they are low and extending by additions of earth cast up by the waves.

The rivers are small but their number is great, and these, with the 5,000 lakes scattered along the watersheds of the State abundantly water all parts of it.

Dr. C. Rominger, a former State Geologist, writes of the geology of the Lower Peninsula as follows: "It forms the center-point of an oceanic bay which seems to have existed without any important alteration in its limits, from the beginning of the Silurian period to the end of the Carboniferous time. We find within the space supposed to have been the bay an uninterrupted series of marine deposits, following each other in the greatest regularity of superposition, which represent all the known formations deposited on this continent from the Silurian period on to the coal formation." The entire surface of the peninsula is covered deeply with glacial drift, consisting of sand, gravels and clays variously intermixed.

The topographical outlines of the Lower Peninsula are due to the joint action of moving ice and flowing water during and following the glacial period.

Beginning in Presque Isle county the lateral moraine of the Huron glacier passes southwest near the line between Montmorency and Alpena counties, thence south by west through Oscoda, Roscommon and Clare, meeting in Mecosta county, the east lateral moraine of Lake Michigan. These join and pass in a southwest direction through Kent, Barry, Kala-

mazoo and St. Joseph counties. The Saginaw bay branch of the Huron glacier unites with the Huron glacier proper, and forms a lateral moraine beginning in Huron county, passing through Sanilac, Lapeer, Oakland, Livingston and portions of Jackson, Washtenaw and Hillsdale counties. This peninsula is divided by these moraines into certain more or less clearly marked floral regions.

CLIMATE AND DISTRIBUTION.

"The sinuosities of the several isothermal lines will demonstrate at a glance the peculiar character of the climate of Michigan, and the fact that both in summer and winter, it is better adapted to the interests of agriculture and horticulture, and probably also to the comfort and health of its citizens, than the climate of any other northwestern state. The marked peculiarity of the climate of Michigan in these respects is attributable to the influence of the great lakes by which the state is nearly surrounded. It has long been known that considerable bodies of water exert a local influence in modifying climate and especially in averting frosts, but it has never been expected that Lake Michigan, for instance, impresses upon the climatic character of a broad region an influence truly comparable with that exerted by the great ocean."—ALEXANDER WINCHELL.

The following general notes on Climate and Distribution are from the preface to the first edition by E. F. Smith:

"The influence of climate on vegetation may be summed up in a few words. The climate of the Lower Peninsula is not as severe as that of the Upper, nor so even, but is subject to frequent, sudden, and extreme changes of temperature—as great a variation during the winter season as 53° Fahr. in less than 24 hours having been recorded. Such rapid changes more or less affect vegetation, especially the tender branches of cultivated trees, which are sometimes seriously injured. In one or two instances a like effect on our forest trees has been noticed. The annual range of temperature is about 116°, and the annual mean 46°. Of rainfall, including what falls in form of snow, we have, yearly, about thirty inches. Our snowfall is much less, for the same latitude, than that of New York and New England. In the center of the peninsula, we seldom have more than a few inches at a time.

"The proximity of the Great Lakes exerts a marked influence in equalizing the temperature and the effects are marked upon our flora.

"Trees, like *Liriodendron Tulipifera, Asimina triloba, Cercis Canadensis, Gleditsia triacanthos, Cornus florida, Nyssa multiflora,* and *Morus rubra,* which belong to Ohio and Central Illinois, have crept northward, favored by the mild influence of the lake winds, through the central and western part of the Lower Peninsula, often beyond the middle, and the same is true of smaller and less noticeable plants.

"As might be expected from the uniform surface of the peninsula, the flora is much alike throughout. Probably three-fourths of our species are common to all sections, though by no means equally distributed; some being very abundant in one district and rare in another at no great distance. In most cases such change is due to soil rather than to difference in elevation, temperature, or atmospheric moisture.

"The Lower Peninsula is covered with a deep drift of alternating sands, clays, and gravels, and the flora of any section depends chiefly on which of these happens to lie uppermost. With reference to its flora, the penin-

sula may roughly be divided into two great divisions—the hardwood and the softwood lands; one representing the Appalachian flora, and the other, the Canadian.

"The hardwood country lies south of latitude 43°, and consists of very fertile sand, clay, or loam, mostly cleared of the original forest, and largely cultivated.

"The sandy or stony drift of many river valleys in this section supports a heavy growth of oak, frequently interspersed with walnut and hickory, while the margins of the streams, and the neighboring swamps, abound in soft maples, swamp and chestnut oak, white and black ash, elm, hackberry, sycamore, butternut, and similar trees. Willows, dogwoods, viburnums, and buttonbush, are common shrubs in the swamps; and hazel, hawthorn, wild cherry and plum, June berry, witch-hazel, etc., are abundant on the dryer ground.

"On the uplands, and away from streams, clay, loam, and a peculiar black muck soil supersede the sands and gravels of the valleys. The prevailing timber here is beech and maple and oak forest in about equal proportions. Beech and maple generally grow together, forming magnificent forests of great extent. The best wheat farms are usually found on uplands near streams, where the oak timber gradually shades into beech and maple. Plains of fertile sand covered with a low, or scattering growth of oak (oak openings) are frequent, and always very desirable for farming purposes.

"Marshes densely covered with tamarack are common in this part of the State, and nourish in their thick shade such plants as *Drosera rotundifolia, Sarracenia purpurea, Rhus venenata, Ribes rubrum, Chiogenes hispidula, Salix candida, Smilacina trifolia, Pogonia ophioglossoides* and *Calopogon pulchellus.* Arbor-vitæ, red cedar and black spruce are comparatively rare.

"A similar tract of soil and timber occurs in the upper end of the peninsula, north of a line drawn from Thunder bay west to the head of Grand Traverse bay. This is commonly known as the "Traverse region," and has a flora much like that we have just described, with the exception that some of the southern species disappear, and northern ones begin to take their place, or if found growing further south, here first become frequent. Deep forests of hemlock and yellow birch (*B. lutea*) mixed with a fine, tall growth of striped maple (*A. Pennsylvanicum*) are frequent, having underneath a tangled growth of *Taxus baccata,* var. *Canadensis,* and under all a carpet of *Lycopodium annotinum.* Alternating with these are sandy plains covered with a dense growth of *Vacciniums,* yielding a great abundance of fruit. Sugar maples and basswood are also abundant in this region, and reach an immense size. In fact, it would be difficult to find finer groves of maple in any part of the State.

"The pine country proper lies between the two tracts we have described, and embraces about 15,000 square miles. It is composed largely of sand hills and plains, either scantily furnished with vegetation, or densely covered with pine forest. Argillaceous tracts wooded with beech and maple also occur, like oases in a desert; and swamps abound, with the usual lowland timber. Forests of hemlock spruce are frequent, and there are occasional ridges of oak. Birch (*B. lutea*) also begins to be a common forest tree, and attains a large size. The usual timber of the barrens is Jack Pine (*P. Banksiana*). Climatic and other influences have combined to produce groves composed entirely of this species of large size and of great

beauty, for, instead of being 'a straggling shrub, or low tree' (Gray), it rises, often 50-60 feet, straight and symmetrical. All through this region *Pinus Strobus* is the prevailing species and furnishes most of the lumber, but *P. resinosa* is frequent as far south as Clare county, and occurs sparingly in the northern part of Isabella county, which appears to be its southern limit.

"Such is the general character of the sylva down to about latitude 43°, but in the western part of the State, owing perhaps to moister climate, or to favorable soil, hemlock spruce is more abundant, and reaches much farther south, nearly or quite to the Indiana line, and the same is true of white pine.

"The flora of the deep pine woods is interesting, though rather monotonous. Very little undergrowth is found, and their gloomy recesses nourish only such plants as love thick shade. Here the club-mosses (*Lycopodiums*) find a congenial home, and flourish luxuriantly, while *Clintonia borealis* covers the ground. The great round-leaved orchid (*Habenaria orbiculata*), with its tall, greenish spike and twin leaves close to the earth, is also frequent and striking. We shall also meet *Mitchella repens, Maianthemum Canadense, Trillium grandiflorum,* perhaps, and a few ferns, particularly *Asplenium Filix-fœmina* and *Phegopteris Dryopteris.* Other species occur, of course, but not so abundantly. In more open places, and on ridges, we meet *Rhus aromatica* and *Comptonia* along with wintergreen (*Gaultheria*) and trailing arbutus (*Epigœa*), and are often fortunate enough to find the wax-white, fragrant flower of *Moneses uniflora,* or *Polygala paucifolia,* hiding its shining leaves under a wealth of showy pink blossoms.

"The floral treasures of the pine region lie, however, in its swamps and lake borders rather than in the deep woods. Therein grows *Linnœa borealis* in all its delicate beauty, carpeting the ground, and close at hand, the odd, brown-purple flower of *Cypripedium acaule* and the small yellow blossom of its water-loving relative *C. parviflorum.* In such swamps, or within a stone's throw of them, may be found many other plants of equal interest, such as *Medeola Virginica, Ledum latifolium, Andromeda Polifolia, Kalmia glauca, Lonicera oblongifolia, Cardamine pratensis, Gerardia aspera, Mitella nuda, Eriophorum vaginatum, etc.* On lake margins we shall find *Lysimachia* and the blue *Pontederia* and more rarely, *Nesœa* and *Eleocharis quadrangulata.* The lake itself, most likely, will be full of *Nymphaea, Nuphar, Utricularias,* and a world of *Potamogetons* and similar water weeds. Shrubby *Vacciniums* line the bluffs, and here and there gleam the white trunks of paper birches against the dark background of pines.

"In the thick-pine country, where the lumberman's ax has let in the sunlight, new plants spring up freely. Here, *Prunus Pennsylvanica* and poplars are frequent, and the blackberry is omnipresent. *Aralia hispida* and *Physalis lanceolata* are also peculiar to such land, and in August *Gnaphalium decurrens* may be seen whitening thousands of acres.

"One seldom beholds a drearier sight than a dead and deserted lumber region. The valuable trees were all felled years ago, and the lumberman moved on to fresh spoils, leaving behind an inextricably confused mass of tree tops, broken logs, and uprooted trunks. Blackberry canes spring up everywhere, forming a tangled thicket, and a few scattering poplars, birches, and cherries serve for arboreal life, above which tower the dead pines, bleached in the weather and blackened by fire, destitute of

limbs, and looking at a distance not unlike the masts of some great harbor. Thousands of such acres, repellant alike to botanist and settler, can be seen in any of our northern counties.

"In certain districts considerable beech is found associated with the pine. The soil of such tracts is usually of better quality, and can be rendered productive without much labor. It may be noted that in such cases the pine also grows thriftier and makes better lumber."

FLORA OF THE JACK-PINE PLAINS.

The plants of this region are all found in one or more of the regions previously enumerated.

The soil of these plains is mainly sand of considerable depth which dries out quickly after a rain, and is then especially liable to be burned over, the burning often destroying every living plant above the surface of the soil. In this way, by repeated burning, much of the vegetable matter is removed, leaving the surface soil thin.

The following seventy species of plants are almost certain to be found in considerable quantity on any extended area of Jack-pine plains:

a. THOSE MOST COMMON.

Amelanchier Botryapium (L. f.) DC. Shad-bush.
(*Amelanchier Canadensis* var. *oblongifolia* T. & G.)
Andropogon furcatus Muhl. Finger, or Beard-grass.
Andropogon scoparius Michx. Beard-grass.
Arctostaphylos Uva-Ursi (L.) Spreng. Bearberry.
Aster lævis L. Smooth Aster.
Carex Pennsylvanica Lam. Pennsylvania Sedge.
Comptonia peregrina (L.) Coulter.
(*Myrica asplenifolia* L.)
Danthonia spicata (L.) Beauv. Wild Oat-grass.
Epigæa repens L. Trailing Arbutus.
Gaultheria procumbens L. Wintergreen.
Leptilon Canadense (L.) Britton. Horse-weed.
(*Erigeron Canadensis* L.)
Oryzopsis juncea (Michx.) B. S. P. Mountain Rice.
(*Oryzopsis Canadensis* Torr.)
Pinus divaricata (Ait.) Gord. Gray Pine. Jack Pine.
(*Pinus Banksiana* Lambert.)
Populus tremuloides Michx. Aspen.
Prunus Pennsylvanica L. f. Wild Red, or Pin Cherry.
Prunus pumila L. Sand Cherry.
Prunus Virginiana L. Choke-Cherry.
Pteridium aquilinum (L.) Kuntze. Eagle Fern.
(*Pteris aquilina* L.)
Quercus coccinea Wang. Scarlet Oak.
Quercus velutina Lam. Black Oak.
(*Quercus tinctoria* Bartram.)
Rumex Acetosella L. Sheep Sorrel.
Salix humilis Marsh. Low willow.
Solidago nemoralis Ait. Golden Rod.
Vaccinium Canadense Richards. Low Blueberry.
Vaccinium Pennsylvanicum Lam. Dwarf Blueberry.
Vaccinium vacillans Kalm. Low Blueberry.

b. THOSE LESS FREQUENT.

Adopogon Virginicum (L.) Kuntze. Dwarf Dandelion. Virginia Goatsbeard.
(*Krigia amplexicaulis* Nutt.)
Agrostis hyemalis (Walt.) B. S. P.
(*Agrostis scabra* Willd.) Hair-Grass.

Antennaria plantaginifolia (L.) Richards. Plaintain-leaved Everlasting.
Apocynum androsæmifolium L. Dogbane.
Aralia hispida Vent. Bristly Sarsaparilla.
Campanula rotundifolia L. Bluebell. Harebell.
Carduus odoratus (Muhl.) Porter.
(*Cnicus pumilus* Torr.)
Ceanothus Americanus L. New Jersey Tea.
Comandra umbellata (L.) Nutt. Bastard Toad-flax.
Convolvulus spithamæus L. Low Bindweed.
Diervilla Diervilla (L.) MacM. Bush Honeysuckle.
(*Diervilla trifida* Moench.
Erigeron ramosus (Walt.) B. S. P. Daisy Fleabane.
(*Erigeron strigosus* Muhl.)
Festuca ovina L. Sheep's Fescue.
Fragaria Virginiana Duchesne. Strawberry.
(*Fragaria Virginiana Illinoensis* A. Gray.)
Gaylussacia resinosa (Ait.) T. & G. Black Huckleberry.
Gnaphalium decurrens Ives. Everlasting.
Helianthemum Canadense (L.) Michx. Frost-wort.
Helianthus divaricatus L. Wild Sunflower.
Helianthus occidentalis Riddell. Wild Sunflower.
Hieracium venosum L. Rattlesnake-weed.
Houstonia longifolia Gaertn. Houstonia.
(*Houstonia purpurea longifolia* A. Gray.)
Koeleria cristata (L.) Pers. Koeleria.
Lacinaria cylindracea (Michx.) Kuntze. Blazing Star.
(*Liatris cylindracea* Michx.)
Lithospermum Gmelini (Michx.) A. S. Hitchcock. Hairy Puccoon.
(*Lithospermum hirtum* Lehm.)
Lycopodium complanatum L. Ground-pine.
Melampyrum lineare Lam. Cow-wheat.
(*Melampyrum Americanum* Michx.)
Monarda fistulosa L. Wild Bergamot.
Onagra biennis (L.) Scop. Evening primrose.
(*Œnothera biennis* L.)
Panicum depauperatum Muhl. Panic-Grass.
Panicum dichotomum L. Panic-Grass.
Pinus resinosa Ait. Norway or Red Pine.
Pinus Strobus L. White Pine.
Polygala polygama Walt. Pink polygala.
Populus grandidentata Michx. Large-toothed Aspen.
Potentilla Canadensis L. Five-finger. Cinque-foil.
Quercus alba L. White Oak.
Rubus Canadensis L. Dewberry.
Rubus hispidus L. Dewberry.
Rudbeckia hirta L. Cone-flower.
Sibbaldiopsis tridentatata (Soland.) Rydb. Three-toothed Cinque-foil.
(*Potentilla tridentata* Ait.)
Solidago juncea Ait. Golden Rod.
Unifolium Canadense (Desf.) Greene.
(*Maianthemum Canadense* Desf.)
Viola arenaria DC. Sand Violet.
(*Viola canina puberula* S. Wats.)
Viola pedata L. Bird-foot Violet.

The above list consists of representatives of thirty families, of fifty-four genera, and of seventy species. The families of plants best represented on the plains are the *Rosaceæ* by six species, *Compositæ* by thirteen species, *Gramineæ* by nine species, *Vacciniaceæ* by four.

The following large and prominent families of the State are not represented in the list given above: *Ranunculaceæ*, *Cruciferæ*, *Caryophyllaceæ*, *Saxifragaceæ*, *Umbelliferæ*, *Orchidaceæ*.

Most remarkable of all is the absence of any *Papilionaceæ* though the family is second in size only to the *Compositæ*. The *Papilionaceæ* is represented in the State by 43 native species and varieties.

The number of biennials given in this list is remarkably small, only two, and there are no annuals in it. Sixty-eight out of seventy are perennials and most persistent plants well adapted by long, deep roots and root-stocks to live in poor soil which is subject to severe droughts. Most of them are admirably adapted to survival after a severe fire has burned over the ground and killed the tops of the plants.

PLANTS INDICATING A FERTILE SOIL.

Acer Saccharum Marsh. Sugar Maple, when the wood is solid and of fine quality.
(*Acer saccharinum* Wang.)
Acer nigrum Michx. Black sugar maple.
(*Acer saccharinum nigrum* Torr. & Gray.)
Adiantum pedatum L. Maidenhair Fern.
Angelica villosa (Walt.) B. S. P. Angelica.
(*Angelica hirsuta* Muhl.)
Apios Apios (L.) Mac M. Ground-nut.
(*Apios tuberosa* Moench.)
Arisæma triphyllum (L.) Torr. Indian Turnip.
Asplenium angustifolium Michx. Spleenwort.
Asplenium acrostichoides Sw. Spleenwort.
(*Asplenium thelypteroides* Michx.)
Cassia Marylandica L. Wild Senna.
Collinsonia Canadensis L. Rich-weed. Stone-root.
Crataegus tomentosa L. Hawthorn.
Dentaria diphylla Michx. Toothwort. Pepper-root.
Dentaria laciniata Muhl. Toothwort. Pepper-root.
Bicuculla Canadensis (Goldie) Millsp. Squirrel Corn.
(*Dicentra Canadensis* DC.)
Bicuculla Cucullara (L.) Millsp. Dutchman's Breeches.
(*Dicentra Cucullaria* DC.)
Fraxinus Americana L. White Ash.
Fraxinus quadrangulata Michx. Blue Ash.
Hydrastis Canadensis L. Golden Seal.
Hydrophyllum Canadense L. Waterleaf.
Hydrophyllum Virginicum L. Waterleaf.
Juglans cinerea L. Butternut.
Juglans nigra L. Black Walnut.
Menispermum Canadense L. Moonseed.
Orchis spectabilis L. Showy Orchis.
Podophyllum peltatum L. May-Apple. Mandrake.
Quercus alba L. White Oak. When well grown.
Quercus macrocarpa Michx. Bur-Oak.
Ribes Cynosbati L. Prickly Gooseberry.
Rubus occidentalis L. Black Raspberry.
Scrophularia Marylandica L. Figwort.
(*Scrophularia nodosa Marylandica* A. Gray.)
Taraxacum Taraxacum (L.) Karst. Dandelion.
(*Taraxacum officinale* Weber.)
Tilia Americana L. Basswood.
Ulmus Americana L. American Elm.
Ulmus racemosa Thomas. Rock Elm.
Uvularia grandiflora J. E. Smith. Bellwort.
Verbena hastata L. Blue Vervain.

When well grown and of good size, several other trees are indications of good soil.

PLANTS PECULIAR TO THE PRAIRIES.

The following plants are peculiar to the prairie region of the southwestern portion of the State:

Amorpha canescens Pursh. Lead-Plant.
Asclepias verticillata L. Milkweed.
Aster sericeus Vent. Aster.
Atheropogon curtipendulus (Michx.) Fourn.
(*Bouteloua curtipendula* (Michx.) Torr.)
Baptisia leucantha Torr. & Gray. False Indigo.
Brauneria purpurea (L.) Britton. Cone-Flower.
(*Echinacea purpurea* Moench.)
Coreopsis palmata Nutt.
Helianthus scaberrimus Ell. Sunflower.
(*Helianthus rigidus* Desf.)
Phlox bifida Beck. Phlox.
Silphium integrifolium Michx. Rosin-weed.
Silphium laciniatum L. Compass-plant.
Silphium perfoliatum L. Cup-plant.

OVERLAPPING OF NORTHERN AND SOUTHERN SPECIES IN THE GRAND RIVER VALLEY.

Northern Species.

Carex Magellanica Lam. Sedge.
Carex pauciflora Lightf. Sedge.
Carex tenuiflora Wahl. Sedge.
Dracocephalum parviflorum Nutt. Dragon-head.
Eriophorum vaginatum L. Cotton-grass.
Lonicera oblongifolia (Goldie) Hook. Swamp Fly-Honeysuckle.
Mimulus Jamesii T. & G. Monkey-flower.
Primula Mistassinica Michx. Primrose.
Symphoricarpos pauciflorus (Robbins) Britton. Snowberry.
(*Symphoricarpos racemosus pauciflorus* Robbins.)
Taxus Canadensis Willd. American Yew. Ground Hemlock.

Southern Species.

Asimina triloba (L.) Dunal. Papaw.
Bidens trichosperma tenuiloba (A. Gray) Britton. Tick-seed. Sunflower.
(*Coreopsis trichosperma* var. *tenuiloba* A. Gray.)
Cassia Marylandica L. Wild Senna.
Cercis Canadensis L. Red-bud. Judas-tree.
Chaerophyllum procumbens (L.) Crantz.
Collinsia verna Nutt. Blue-eyed Mary.
Eleocharis interstincta (Vahl.) R. & S. Spike-rush.
(*Eleocharis equisetoides* Torr.)
Eleocharis mutata (L.) R. & S. Spike-rush.
(*Eleocharis quadrangulata* R. & S.)
Eleocharis olivacea Torr. Spike-rush.
Eleocharis Robbinsii Oakes. Spike-rush.
Eleocharis rostellata Torr. Spike-rush.
Erigenia bulbosa (Michx.) Nutt. Harbinger of Spring.
Gymnocladus dioica (L.) Koch. Ky. Coffee-tree.
(*Gymnocladus Canadensis* Lam.)
Hicoria laciniosa (Michx. f.) Sarg. King-nut.
(*Carya sulcata* Nutt.)
Hemicarpha micrantha (Vahl.) Britton.
(*Hemicarpha subsquarrosa* Nees.)
Liriodendron Tulipifera L. White-wood. Tulip-tree.
Meibomia Marylandica (L.) Kuntze. Tick-Trefoil.
(*Desmodium Marilandicum* Boot.)
Morus rubra L. Red Mulberry.

Silphium terebinthinaceum Jacq. Prairie Dock.
Tradescantia Virginica L. Common Spiderwort.
Utricularia resupinata B. D. Greene. Bladderwort.

COMPARISON OF THE FLORA OF THE EASTERN AND THE WESTERN SIDES OF THE STATE IN THE LATITUDE OF 44°40′.

On the east side, the latitude in question is near Harrisville in Alcona county. On the west side it is near Frankfort in Benzie county.

It has long been known that the climate of the west shore where the wind sweeps across Lake Michigan was milder in winter, and throughout the year less variable than it is on the east side of the State. So far as observed, the plants of the State which are only found in the vicinity of the great lakes are more abundant in individuals on the west shore.

A. NORTHERN PLANTS FOUND ON THE EAST SIDE OF THE STATE AND NOT ON THE WEST.

Botrychium Lunaria (L.) Swartz. Moonwort.
Botrychium simplex Hitchcock. Moonwort.
Carex capillaris L. Sedge.
Carex durifolia Bailey. Back's Sedge.
(*Carex Backii* Boott.)
Carex Houghtonii Torr. Sedge.
Dracocephalum parviflorum Nutt. Dragon-head.
Kalmia angustifolia L. Sheep Laurel. Lambkill.
Kalmia glauca Ait. Swamp Laurel. Pale Laurel.
Picea Canadensis (Mill) B. S. P. White Spruce.
(*Picea alba* Link.)
Ribes lacustre (Pers.) Poir. Swamp Gooseberry.
Sparganium simplex Huds.

B. SOUTHERN PLANTS FOUND ON THE WEST SIDE OF THE STATE AND NOT ON THE EAST.

Adiantum pedatum L. Maiden Hair Fern.
Acer saccharinum L. Silver Maple.
(*Acer dasycarpum* Ehrh.)
Rubus occidentalis L. Black Raspberry.
Sambucus Canadensis L. Common Elder.
Sassafras Sassafras (L.) Karst.
(*Sassafras officinale* Nees.)
Ulmus fulva Michx. Red Elm.
Ulmus racemosa Thomas. Rock Elm.

This list is doubtless incomplete, but so far as it goes it sustains the prevailing notion that the west side of the State has the milder climate. We might be able to see why silver maple, sassafras, black raspberry, red elm and rock elm thrive on the west shore and not on the east, but we are unable to see why the northern plants found on the east shore should not be found on the west shore. Perhaps there is some other reason than the difference of climate of the present day.

PLANTS SUPPOSED TO HAVE IMMIGRATED FROM THE NORTHEAST.

Calypso bulbosa (L.) Oakes.
(*Calypso borealis* Salisb.)
Clintonia borealis (Ait.) Raf.
Equisetum littorale Kuehl.
Eriocaulon septangulare Withering. Pipewort.
Gyrostachys stricta Rydb. Hooded Ladies' Tresses.
(*Gyrostachys Romanzffiana* (Cham.) MacM.)
Selaginella selaginoides (L.) Link.
(*Selaginella spinosa* Beauv.)

Trillium erectum L. Wake Robin.
Trillium undulatum Willd. Painted Wake-Robin.
(*Trillium erythrocarpum* Michx.)

PLANTS SUPPOSED TO HAVE IMMIGRATED FROM THE NORTH AND WEST.

Adenocaulon bicolor Hook.
Anemone parviflora Michx. Anemone.
Artemisia gnaphalodes Nutt. Mugwort.
(*Artemisia Ludoviciana gnaphalodes* T. & G.)
Brauneria pallida (Nutt.) Britton. Purple Cone-flower.
(*Echinacea angustifolia* DC.)
Bromus breviaristatus (Hook.) Buckl. Brome Grass.
Castilleja acuminata (Pursh) Spreng. Painted-cup.
(*Castilleja pallida septentrionalis* A. Gray.)
Drosera linearis Goldie. Sundew.
Euphorbia serpyllifolia Pers. Spurge.
Iva xanthiifolia (Fresen.) Nutt. Marsh Elder.
Lonicera involucrata (Richards) Banks. Honeysuckle.
Mertensia paniculata (Ait.) G. Don. Lungwort.
Mimulus Jamesii T. & G. Monkey-flnower.
Mimulus moschatus Dougl. Musk-flower.
Panicularia pallida (Torr.) Kuntze.
(*Glyceria pallida* Trin.)
Roripa obtusa (Nutt.) Britton.
(*Nasturtium obtusum* Nutt.)
Parnassia palustris L. Grass of Parnassus.
Parnassia parviflora DC. Grass of Parnassus.
Phacelia Franklinii (R. Br.) Gray.
Polygonum lapathifolium incanum (Schmidt) Koch. Knotweed.
Potentilla Robbinsiana Oakes.
(*Potentilla frigida* A. Gray.)
Ranunculus reptans intermedius (Hook.) T. & G. Creeping Spearwort.
(*Ranunculus Flammula intermedius* Hook.)
Rosa Engelmanni Watson. Rose.
Rosa Sayi Schwein. Rose.
Rumex salicifolius Weinm. White Dock.
Sorbus sambucifolia (C. & S.) Roem. Western Mountain Ash.
(*Pyrus sambucifolia* Cham. & Schlecht.)
Symphoricarpos occidentalis Hook. Wolfberry.

TREES OF MICHIGAN COMPARED WITH THOSE OF EUROPE.

Michigan is very rich in trees. If we have counted correctly there are 90 species of indigenous trees and three exotics which have escaped from cultivation.

To comprehend the relative importance of our trees, let us glance at the forests of Great Britain. Great Britain and Ireland contain 121,260 square miles of land, Michigan 60,000, a little less than one-half as much as Great Britain. She has one species of basswood not so good as ours; one maple not over twenty feet high; one cherry from ten to twenty feet high; one small ash, two elms, two poplars, one beech, which grows very large but not very high; one small white birch, one species of pine, by no means a match for our white pine; a species of oak which sometimes grows to a great size.

Great Britain has about ten species of trees native to her soil. Michigan, with half the territory, has eighty-five species. Great Britain has no whitewood, no white or red cedar, no walnuts or hickories. Michigan has six species of maple of tree size, a basswood, a whitewood, honey locust, Kentucky coffee tree. three cherries, a pepperidge, five species of ash, a

sassafras, three elms, a hackberry, a mulberry, a buttonwood, black walnut, butternut, eight hickories, thirteen oaks, a chestnut, a beech, four tree birches, three willows of tree size, five poplars, three pines, three spruces, one hemlock, a balsam fir, one larch, one arbor-vitæ and a red cedar.

In all Europe there are only 85 species of trees.

WHY HAS MICHIGAN SO MANY TREES AND GREAT BRITAIN SO FEW.

This question now very naturally arises: Why has the Atlantic region, including Michigan, so many species of trees and why has Europe so few? Certainly we cannot attribute this difference to a defective soil and climate of Europe, as they now exist, for Europe can grow all sorts of trees now found in the temperate zone, while "Great Britain alone can grow double or treble the number of trees that the Atlantic States can."

The former geological conditions of their continents help to explain all this difference in the distribution of trees to the entire satisfaction of scientists.

Away back in the Tertiary Period the trees of the regions now possessing an arctic climate were such as now thrive in a warm temperate zone like that of Georgia and California. This is well illustrated by the abundant fossil remains of trees. Following this, came a long time when extreme cold prevailed, known as the Glacial Epoch, when snow and ice for most or all of the year extended to the Ohio river. At the approach of cold, the trees slowly retreated southward, as generation followed generation. The plants such as now thrive in southern Michigan, perhaps then extended to what now forms the State of Alabama, while the arctic plants reached Ohio.

As the climate again gradually grew warmer, the trees and other plants slowly migrated northward. Some arctic plants were stranded on the White mountains and in Labrador, where they still remain; others went farther north.

Plants of the cool temperate zone reached Michigan. In a similar manner, during the Glacial Epoch the plants of Europe were driven southward. The Alps, the Pyrenees, the Appenines, the Caucasus, still contain some of these arctic plants which retreated there at the close of the Glacial Epoch. Most of the plants of the warm temperate region had perished and therefore were unable to retreat when the continent became warmer.

I quote the words of Dr. A. Gray, from whom other hints are taken, as found in the American Journal of Science, page 194, 1878. "I conceive that three things have conspired to this loss. First, Europe hardly extending south of latitude 40°, is all within the limits generally assigned to severe glacial action. Second, its mountains trend east and west, from the Pyrenees to the Carpathians and the Caucasus beyond, near its southern border; and they had glaciers of their own, which must have begun operations, and poured down the northward flanks, while the plains were still covered with forest on the retreat from the great ice wave coming from the north. Attacked both on front and rear, much of the forest must have perished then and there. Third, across the line of retreat of those which may have flanked the mountain ranges, or were stationed south of them, stretched the Mediterranean, an impassable barrier.

"Greenland may be referred to, by way of comparison, as a country

which, having undergone extreme glaciation, bears the marks of it in the extreme poverty of its flora, and in the absence of the plants to which its southern portion, extending six degrees below the arctic circle, might be entitled. It ought to have trees, and might support them. But since destruction by glaciation, no way has been open for their return.

"In the American continent the mountains run north and south. The trees, when touched on the north by the on-coming refrigeration, had only to move their southern border southward, along an open way, and there was no impediment to their due return. So our lines have been cast in pleasant places, and the goodly heritage of forest trees is one of the consequences."

NATIVE FORAGE PLANTS.

The native grasses while young, from early spring to winter, without exception, furnish a tender bite which is very acceptable to all sorts of horses, cattle, sheep, and even swine and poultry. Most of the native grasses (about 120 in number) grow in isolated bunches, more or less frequent, and furnish but little food, while a considerable portion soon become unpalatable as the summer approaches. The most important native grasses for pasture or meadow are found in open places and are: June grass (not native), fowl meadow-grass, and several other species of *Poa* which have no well-known common name; several manna-grasses (*Paniculariа*); a few small fescues, two or three Eatonias; blue-joint, two or three small redtops, and several species of Muhlenberg's grass, two or three kinds of mountain rice, reed-grass, two beard-grasses, and several panic grasses. The following, though quickly and thoroughly disseminated, are not native to our State: Barn-yard grass, crab-grass, foxtail or pigeon-grass, sweet vernal-grass, Timothy, meadow foxtail, the larger redtop so popular for lowlands, tall oat-grass, orchard grass, tall fescue, perennial rye-grass, quick-grass.

Besides the true grasses, there are about one hundred and eighty species of grass-like plants mostly found on marshes, which are called *sedges*. They are mostly confounded with the grasses except by the botanist. In addition to these are some twenty-five other narrow-leaved and grass-like plants known as *rushes* which furnish more or less herbage. Many of the sedges and rushes make hay of fair quality when cut early before they become harsh and woody.

On the Jack-pine plains and in many other portions of the sandy stump-lands in the northern counties are two species of sedges found abundantly on dry land. These are *Carex Pennsylvanica* Lam., and *C. umbellata vicina* Dewey; they are popularly called grasses by the residents of those counties. Inquiries have frequently been made concerning the name, and statements made to the effect that they were very nutritious and that cattle got fat on them early in the spring. So far as I have observed, cattle prefer something else as it starts in spring, thus leaving the sedges untouched to grow all summer and remain green under the snow till the following spring. In spring the cattle eagerly devour the last year's growth of the sedges and thrive on it. This is but another evidence that the quality of the fodder is not of so great importance, provided animals can get enough of it, and can be induced to eat it.

Besides the native grasses, sedges and rushes, there are great numbers of other herbaceous plants, usually one here and there which are eaten by

hungry stock. Leaves and the tender growth of shrubs and young trees are often eaten in great quantities, and in times of scarcity cattle get astride of young trees bending down the tops so as to reach the leaves and twigs. Sheep eat the greatest variety of plants and thus where they feed closely are valuable aids in the clearing of a new country. They eat briars, elders and sprouts of stumps, but will leave some things which they do not like. The weeds that are thus left have an excellent chance to spread and in time occupy much of the ground once covered by the nutritious and edible plants which have been removed. Here we see one of the most excellent reasons for the interference of man in removing the weeds and in encouraging the introduction of the more valuable fodder plants. While these early native pastures and meadows are of incalculable value to the pioneer, without some care they soon deteriorate.

In times of great scarcity of feed in a new country, farmers used to fell trees that cattle might eat the twigs, or "browse," which contain much starch and protoplasm stored away by the trees ready to be used in giving the young growth a start on the approach of warm weather.

Michigan has no native species of clover.

NATIVE BEE PLANTS.

In Michigan there is a very large number of plants which furnish a good quality of honey. If the species is abundant in any region, it usually becomes known to the apiarist as a good bee plant; if not abundant it very likely fails to attract attention. A plant may be rare or important in one region and abundant in another. In autumn, asters and golden rods are known as excellent bee plants, because some few of the many species in the State are plentiful in nearly every neighborhood, but the same sorts of asters or golden rods do not everywhere throughout the State furnish a great amount of the honey. As a rule those plants which produce odorous or showy flowers afford honey and will be visited by honey bees unless the flower is of a shape which makes it impossible for the bee to reach the food.

Probably in the State there are of native plants, introduced weeds and field crops, a thousand species which furnish excellent food for bees. This is nearly one hundred times as many as the bee keeper has in mind, unless he has given unusual attention to the subject.

Our open low lands furnish a large proportion of the bee pasture; the forest some; the weeds and some of the field, garden and orchard crops a fair amount. Extremely dry or very wet weather are both unfavorable to the yield of honey. Drainage of the swamps and the clearing of waste places are unfavorable to the interests of the bee keeper.

As the botanist now looks at the subject, colors and odors are mere advertisements to call the attention of insects to the rich supplies of food in store for them. It may be said that the honey is there for the bees, but primarily it is there for the good of the plant, secondarily for the good of the insect. Had good old Dr. Watts lived in our day, he would have no doubt written his familiar verse in this way:

How doth the little busy bee
Improve each shining hour!
By carrying pollen day by day
To fertilize each flower.

WEEDS, NATIVE AND INTRODUCED.

A new country is comparatively free from weeds, but as the years pass by, one after another, weeds are introduced. Some of them are natives of the neighborhoods in which they are found. but most of them have been introduced from other portions of our own country or from foreign countries. The farmer is not inclined to adopt Emerson's notion of a weed as "a plant whose virtues have not yet been discovered," at least he doesn't purpose trying to find a use for them. The seeds of most weeds find their way onto a farm nicely mixed with seeds of grasses, grains and clovers, which are drilled in or sowed broadcast on fertile soil, where they are afforded an excellent opportunity to grow and multiply. In some instances weeds are introduced as a part of the packing or straw employed to protect castings, marble, crockery or fruit trees. Such foreign packing should always be burned at once.

By these processes the older the country the more troublesome weeds it will have, as every new intruder usually comes to stay. In most cases a weed becomes well established before it is discovered and the inquiry comes: "What is it, and how can I get rid of it?"

Weeds are likely to become most troublesome, where farmers are slovenly and where they are not thorough in cultivating throughout the season the so-called "hoed crops." In regions where "hoed crops" are seldom raised or only raised in small quantity, certain weeds are likely to prove very annoying. The majority of our weeds have been introduced from Europe, as will be seen by a comparison of the two lists which follow:

LIST OF WEEDS INTRODUCED FROM EUROPE AND ASIA.

The List includes at least seventy-eight species, and very likely more.
Abutilon Abutilon (L.) Rusby. Velvet Leaf.
(*Abutilon Avicennæ* Gaertn.)
Agrostemma Githago L. Corn Cockle.
(*Lychnis Githago* Scop.)
Alsine media L. Common Clickweed.
(*Stellaria media* Cyr.)
Alyssum alyssoides (L.) Gouan. Yellow Alyssum.
(*Alyssum calycinum* L.)
Amaranthus hybridus L. Slender Pigweed.
(*Amaranthus chlorostachys* Willd.)
Amaranthus retroflexus L. Rough Pigweed.
Anthemis Cotula L. May-weed.
Arctium Lappa L. Burdock.
Brassica arvensis (L.) B. S. P. Charlock.
(*Brassica Sinapistrum* Boiss.)
Brassica juncea (L.) Cosson. Indian Mustard.
Brassica nigra (L.) Koch. Black Mustard.
Bromus hordeaceus L. Soft-Chess.
(*Bromus mollis* L.)
Bromus racemosus L. Upright-Chess.
Bromus secalinus L. Cheat. Chess.
Bursa Bursa-pastoris (L.) Britton. Shepherd's Purse.
(*Capsella Bursa-pastoris* Medic.)
Camelina microcarpa Andrz. Small-fruited False-flax.
Camelina sativa (L.) Crantz. False-flax.
Carduus arvensis (L.) Robs. Canada Thistle.
(*Cnicus arvensis* Hoffm.)
Carduus lanceolatus L. Common or Bull Thistle.
(*Cnicus lanceolatus* Willd.)

Chaetochloa glauca (L.) Scribn. Yellow Fox-tail.
(*Setaria glauca* Beauv.)
Chaetochloa viridis (L.) Scribn. Green Fox-tail.
(*Setaria viridis* Beauv.)
Chenopodium album L. Lamb's Quarters. Pigweed.
Chenopedium glaucum L. Oak-leaved Goosefoot.
Chrysanthemum Leucanthemum L. White Daisy. Ox-Eye Daisy.
Cichorium Intybus L. Chicory.
Conium maculatum L. Poison Hemlock.
Convolvulus arvensis L. Bindweed.
Cynoglossum officinale L. Hound's-Tongue.
Datura Stramonium L. Stramonium. Thorn-apple.
Datura Tatula L. Purple Thorn-apple.
Daucus Carota L. Carrot.
Diplotaxis muralis (L.) DC. Sand Rocket.
(*Sisymbrium murale* L.)
Dipsacus sylvestris Huds. Wild Teasel.
Echinops sphaerocephalus L.
Eragrostis major Host. Stink grass.
Euphorbia Cyparissias L. Cypress Spurge.
Hypericum perforatum L. St. John's wort.
Inula Helenium L. Elecampane.
Lactuca Scariola L. Prickly Lettuce.
Lappula Lappula (L.) Karst. Stickseed.
(*Echinospermum Lappula* Lehm.)
Leonurus Cardiaca L. Motherwort.
Lepidium apetalum Willd. Apetalous Pepper-grass.
(*Lepidium intermedium* A. Gray.)
Lepidium sativum L. Pepper-grass.
Linaria Linaria (L.) Karst. Butter-and-eggs. Toad-flax.
(*Linaria vulgaris* Mill.)
Lithospermum arvense L. Red root. Corn Gromwell. Wheat Thief.
Malva rotundifolia L. Common Mallow.
Medicago lupulina L. Black Medic.
Melilotus alba Desf. White Melilot. Sweet Clover.
Nepeta Cataria L. Catnep. Catmint.
Panicum capillare L. Witch Grass.
Panicum Crus-galli L. Barnyard-Grass.
Plantago lanceoata L. Ribgrass. English Plantain.
Polygonum Convolvulus L. Black Bindweed.
Portulaca oleracea L. Purslane.
Ranunculus acris L. Buttercup.
Ranunculus bulbosus L. Buttercup.
Ranunculus repens L. Creeping Buttercup.
Roripa sylvestris (L.) Bess. Yellow Water-cress.
(*Nasturtium sylvestre* R. Br.)
Rumex Acetosella L. Field or Sheep Sorrel.
Rumex crispus L. Curled Dock.
Rumex obtusifolius L. Bitter Dock.
Salsola Tragus L. Russian Thistle.
Saponaria officinalis L. Soapwort. Bouncing Bet.
Silene noctiflora L. Night-flowering Catchfly.
Silene vulgaris (Moench) Garcke. Bladder Campion.
(*Silene Cucubalus* Wibel.)
Sisymbrium altissimum L. Tall Sisymbrium.
Sisymbrium officinale (L.) Scop. Hedge Mustard.
Sonchus arvensis L. Sow-Thistle.
Sonchus asper (L.) All. Spiny-leaved Sow-Thistle.
Sonchus oleraceus L. Snow-Thistle.
Syntherisma linearis (Krock.) Nash. Small Grab-grass.
(*Panicum glabrum* Gaud.)
Syntherisma sanguinalis (L.) Dulac. Crab-grass.
(*Panicum sanguinale* L.
Tanacetum vulgare L. Common Tansy.
Taraxacum Taraxacum (L.) Karst. Dandelion.

(*Taraxacum officinale* Weber.)
Tragopogon pratensis L. Goat's-beard.
Verbascum Blattaria L. Moth Mullen.
Verbascum Thapsus L. Mullen.
Xanthium spinosum L. Spiny Clot-bur.

LIST OF INDIGENOUS WEEDS.

This includes thirty-two species and very likely many more.
Acnida tamariscina tuberculata (Moq.) Uline & Bray. Water-hemp.
Acnida tamariscina prostrata Uline & Bray. Prostrate water-hemp.
Amaranthus blitoides S. Wats. Prostrate Amaranth.
Amaranthus graecizans L. Tumble-weed.
(*Amaranthus albus* L.) Tumble-weed.
Ambrosia artemisiaefolia L. Roman Wormwood. Hogweed. Ragweed.
Artemisia biennis Willd. Biennial Wormwood.
Asclepias Syriaca L. Common Milkweed.
(*Asclepias Cornuti* Decaisne.)
Bidens frondosa L. Beggar-ticks. Stick-tight.
Cenchrus tribuloides L. Hedge-hog or Bur-grass.
Chenopodium hybridum L. Maple-leaved Goosefoot.
Cyperus rotundus L. Nut-grass.
Cyperus esculentus L. Nut-Grass.
Erigeron annuus (L.) Pers. Fleabane.
Erigeron ramosus (Walt.) B. S. P. Daisy Fleabane.
(*Erigeron strigosus* Muhl.)
Euphorbia maculata L. Spotted Spurge.
Euphorbia nutans Lag. Spurge.
(*Euphorbia Preslii* Guss.)
Hedeoma pulegioides (L.) Pers. Pennyroyal.
Helianthus tuberosus L. Jerusalem Artichoke.
Iva xanthiifolia (Fresen.) Nutt. Marsh Elder.
Lepidium Virginicum L. Peppergrass.
Leptilon Canadense (L.) Britton. Horse-weed. Fleabane.
(*Erigeron Canadensis* L.)
Onagra biennis (L.) Scop. Evening-primrose.
(*Œnothera biennis* L.)
Panicum capillare L. Old-witch Grass.
Plantago major L. Plantain.
Plantago Rugelii Decaisne. Plantain.
Polygonum aviculare L. Knotgrass.
Polygonum erectum L. Erect Knotgrass.
Polygonum littorale Link. Shore Knotweed.
Prunella vulgaris L. Heal-all.
(*Brunella vulgaris* L.)
Rudbeckia hirta L. Yellow Daisy. Black-eyed Susan.
Urtica gracilis Ait. Tall Wild Nettle.
Xanthium Canadense Mill. Cocklebur. Clotbur.

NATIVE POISONOUS PLANTS.

Besides two species of nettles, *Urtica gracilis* Ait., and *Urticastrum divaricatum* (L.) Kuntze, there are only two species of native plants which are of any prominence as poisonous to the touch, and often these two are not poisonous to all persons. The latter plants or shrubs are known as poison sumach or poison dogwood (*Rhus Vernix* L.), and poison ivy, or poison oak (*Rhus radicans* L.). The last named plant is the most troublesome, as it is often common on dry land, while the one previously noticed is confined to swamps which are not so often frequented. Poison ivy has three leaflets to each leaf, while the handsome Virginian Creeper, often confounded with it, usually has five leaflets to the leaf, and is harmless. The roots, leaves or flowers of many of the

medicinal plants are more or less poisonous when eaten in sufficient quantity, but fortunately, in nearly all such cases, there is something repulsive to the taste or to the smell. Unless one is a botanist or takes the advice of a good botanist, it is safest to let strange plants alone.

NATIVE PLANTS FAST DISAPPEARING.

The fathers and grandfathers of many of us spent a considerable portion of their energies in clearing away trees, shrubs, and breaking up the land that they might have fertile fields. The newer portions of our State are still rapidly undergoing this same transition.

As the country becomes older and more thickly settled almost everything seems to conspire against the trees and smaller plants. Proprietors are still making extensions to their clearings. The "tidy" farmer ditches the cat-holes and marshes, clears out the elders and viburnums to make more room for turnips and better grasses. He turns stock into the wood lot and the flowers of spring and summer retreat to the brush-heaps and a few places inaccessible to the cattle. Fire burns out the dried-up swamps. The officers of the railroad see that the strips alongside the track are often mowed. The highways are attacked and the larger vegetation removed. In places, nurserymen or their agents collect large numbers of the choicer wild plants, as prairie roses and lady's slippers. Near high schools and colleges, the student collectors exterminate many choice plants, root and branch. Many wild plants vigorously protest against these attempts toward their extermination, and start again and again to recover the lost ground, but with the hand of a thorough farmer against them, sooner or later they succumb, the scattered remnants only surviving in the few remaining swamps, along railroads, on a few ragged hills and out of the way places.

Let me utter a vigorous protest against the practice of collecting great quantities of flowers, just to carry home and throw away. And may I hope that every teacher who chances to read this paragraph will also utter a protest against this practice?

LIST OF TREES INDIGENOUS TO MICHIGAN.

The distinction between a tree and a shrub is a purely arbitrary one. If the trunk attains a diameter of one foot the species ranks as a tree, if less than a foot it ranks as a shrub. The papaw and the witch-hazel rank as trees in some regions of the country, but in Michigan I have not included them in the list of trees of the State. I enumerate ninety trees for Michigan.

Abies balsamea (L.) Miller. Balsam Fir.
Acer Negundo L. Box Elder. Ash-leaved Maple. Box Elder.
(*Negundo aceroides* Moench.)
Acer nigrum Michx. Black Sugar Maple.
(*Acer saccharinum nigrum* T. & G.)
Acer Pennsylvanicum L. Striped maple.
Acer rubrum L. Red Maple. Soft Maple.
Acer saccharinum L. Silver Maple. Soft Maple.
(*Acer dasycarpum* Ehrh.)
Acer Saccharum Marsh. Sugar Maple.
(*Acer saccharinum* Wang.)
Æsculus glabra Willd. Fetid or Ohio Buckeye.
Amelanchier Botryapium (L. f.) DC. Shad-bush.

Amelanchier Canadensis (L.) Medic. Shad-bush. June-berry.
Betula lenta L. Cherry Birch. Sweet B. Black B.
Betula lutea Michx. f. Yellow or Gray Birch.
Betula papyrifera Marshall. Paper or Canoe Birch.
Carpinus Caroliniana Walt. Blue Beech.
Castanea dentata (Marsh.) Borkh. Chestnut.
(*Castanea sativa Americana* A. Gray.)
Celtis occidentalis L. Hackberry.
Cercis Canadensis L. Red Bud. Judas tree.
Cornus florida L. Flowering Dogwood.
Crataegus Crus-galli L. Cockspur Thorn.
Crataegus gemmosa Sargent.
Crataegus macracantha Lodd. Long-spined Thorn.
(*Crataegus coccinea macracantha* Dudley.)
Crataegus mollis (T. & G.) Scheele. Hawthorn.
(*Crataegus coccinea mollis* Torr & Gray.)
Crataegus punctata Jacq. Hawthorn.
Crataegus tomentosa L. Hawthorn.
Fagus Americana Sweet. Beech.
(*Fagus ferruginea* Ait.)
Fraxinus Americana L. White Ash.
Fraxinus lanceolata Borck. Green Ash.
(*Fraxinus viridis* Michx.)
Fraxinus nigra Marsh. Black Ash.
(*Fraxinus sambucifolia* Lam.)
Fraxinus Pennsylvanica Marsh. Red Ash.
(*Fraxinus pubescens* Lam.)
Fraxinus quadrangulata Michx. Blue Ash.
Gleditsia triacanthos L. Honey-Locust.
Gymnocladus dioica (L.) Koch. Kentucky Coffee-tree.
(*Gymnocladus Canadensis* Lam.)
Hicoria alba (L.) Britton. Mocker-nut.
(*Carya tomentosa* Nutt.)
Hicoria borealis Ashe. Northern Hickory.
Hicoria glabra (Mill.) Britton. Pig-nut Hickory.
(*Carya porcina* Nutt.)
Hicoria laciniosa (Michx. f.) Sargent. King-nut, Big Shell-bark Hickory.
(*Carya sulcata* Nutt.)
Hicoria microcarpa (Nutt.) Britton. Small Pig-nut Hickory.
(*Carya microcarpa* Nutt.)
Hicoria minima (Marsh.) Britton. Bitter-nut Hickory.
(*Carya amara* Nutt.)
Hicoria ovata (Mill.) Britton. Shag-bark Hickory.
(*Carya alba* Nutt.)
Hicoria villosa Ashe.
Juglans cinerea L. Butternut.
Juglans nigra L. Black Walnut.
Juniperus Virginiana L. Red Cedar.
Larix laricina (Du Roi.) Koch. Tamarack. American Larch.
(*Larix Americana* Michx.)
Liriodendron Tulipifera L. Tulip-tree. Whitewood.
Morus rubra L. Red Mulberry.
Nyssa sylvatica Marsh. Pepperidge. Tupelo. Black or Sour Gum.
Ostrya Virginiana (Mill.) Willd. Ironwood. Hop-hornbeam.
Picea brevifolia Peck. Swamp Spruce.
Picea Canadensis (Mill.) B. S. P.
(*Picea alba* Link.)
Picea Mariana (Mill.) B. S. P. Black Spruce.
(*Picea nigra* Link.)
Pinus divaricata (Ait.) Gord. Jack or Scrub Pine.
(*Pinus Banksiana* Lambert.)
Pinus resinosa Ait. Norway or Red Pine.
Pinus Strobus L. White Pine.
Platanus occidentalis L. Buttonwood. Sycamore.
Populus balsamifera L. Balsam Poplar.

Populus candicans Ait. Balm of Gilead.
(*P. balsamifera candicans* A. Gray.)
Populus deltoides Marsh. Cottonwood.
(*Populus monilifera* Ait.)
Populus grandidentata Michx. Large-toothed Aspen.
Populus heterophylla L. Downy Poplar.
Populus tremuloides Michx. American Aspen.
Prunus nigra Ait. Canada Plum. Horse Plum.
Prunus Pennsylvanica L. f. Wild Red Cherry. Pin Cherry.
Prunus serotina Ehrh. Black Cherry.
Prunus Virginiana L. Choke Cherry.
Quercus acuminata (Michx.) Houda. Chestnut Oak.
(*Quercus Muhlenbergii* Engelm.)
Quercus alba L. White Oak.
Quercus Alexanderi Britton. Alexander's Oak.
Quercus ellipsoidalis E. J. Hill. Hill's Oak.
Quercus coccinea Wang. Scarlet Oak.
Quercus imbricaria Michx. Shingle Oak.
Quercus macrocarpa Michx. Bur Oak.
Quercus minor (Marsh.) Sargent. Post Oak.
(*Quercus obtusiloba* Michx.)
Quercus palustris DuRoi. Pin Oak.
Quercus platanoides (Lam.) Sudw. Swamp White Oak.
(*Quercus bicolor* Willd.)
Quercus rubra L. Red Oak.
Quercus Schneckii Britton. Schneck's red oak.
(*Q. Texana* Sargent.)
Quercus velutina Lam. Black Oak.
(*Quercus tinctoria* Bartram.)
Salix amygdaloides Anders. Willow.
Salix nigra Marsh. Black Willow.
Sassafras Sassafras (L.) Karst. Sassafras.
(*Sassafras officinale* Nees.)
Sorbus Americana Marsh. American Mountain Ash.
(*Pyrus Americana* DC.)
Sorbus sambucifolia (C. & S.) Roem. Western Mountain Ash.
(*Pyrus sambucifolia* C. & S.)
Thuja occidentalis L. White Cedar. Arbor Vitae.
Tilia Americana L. Basswood. Linden.
Tsuga Canadensis (L.) Carr. Hemlock.
Ulmus Americana L. American Elm.
Ulmus fulva Michx. Slippery or Red Elm.
Ulmus racemosa Thomas. Rock Elm.
Viburnum Lentago L. Sheep-berry.

Four or more hybrid oaks have been found, but these were not enumerated above. They are: Quercus alba× macrocarpa, Quercus Leana,= Q. velutina× imbricaria, Quercus platanoides× macrocarpa, Quercus imbricaria× rubra.

LIST OF SHRUBS INDIGENOUS TO MICHIGAN.

There are two hundred and nine species and varieties.

Acer Pennsylvanicum L. Striped Maple.
Acer spicatum Lam. Mountain Maple.
Alnus Alnobetula (Ehrh.) K. Koch. Green Alder.
(*Alnus viridis* DC.)
Alnus incana (L.) Willd. Speckled Alder.
Alnus rugosa (DuRoi.) K. Koch. Smooth Alder.
Amelanchier alnifolia Nutt. June or Service Berry.
Amelanchier rotundifolia (Michx.) Roem. June or Service Berry.
Amorpha canescens Pursh. Lead-plant.
Andromeda Polifolia L. Wild Rosemary.
Aralia hispida Vent. Bristly Sarsaparilla.
Arctostaphylos Uva-Ursi (L.) Spreng. Bearberry.
Aronia arbutifolia (L.) Medic. Red Chokeberry.
Aronia nigra (Willd.) Britton. Black Chokeberry.

(*Pyrus arbutifolia* L. f.)
Aronia nigra (Willd.) Britton.
(*Pyrus arbutifolia melanocarpa* Hook.)
Asimina triloba (L.) Dunal. Papaw.
Benzoin Benzoin (L.) Coulter. Spice-bush.
(*Lindera Benzoin* Blume.)
Betula glandulosa Michx. Glandular Birch.
Betula pumila L. Low Birch.
Ceanothus Americanus L. New Jersey Tea. Red-root.
Ceanothus ovatus Desf. Smaller Red-root.
Celastrus scandens L. Shrubby or Climbing Bittersweet.
Cephalanthus occidentalis L. Button-bush.
Chamaedaphne calyculata (L.) Moench. Leather-leaf.
(*Cassandra calyculata* D. Don.)
Comptonia peregrinia (L.) Coulter. Sweet-fern.
(*Myrica asplenifolia* L.)
Cornus alternifolia L. f. Alternate-leaved Cornel.
Cornus Amonum Mill. Kinnikinnik.
(*Cornus sericea* L.)
Cornus Baileyi Coult. & Evans. Bailey's Cornel.
Cornus candidissima Marsh. Panicled Cornel.
(*Cornus paniculata* L. Her.)
Cornus circinata L. Her. Round-leaved Cornel.
Cornus stolonifera Michx. Red-osier Cornel.
Cornus stricta Lam. Stiff Cornel.
Corylus Americana Walt. Hazel-nut.
Corylus rostrata Ait. Beaked Hazel-nut.
Crataegus acutiloba Sargent.
Crataegus albicans Ashe.
Crataegus altrix Ashe.
Crataegus ater Ashe.
Crataegus attenuata Ashe.
Crataegus borealis Ashe.
Crataegus brevispina (Dougl.) Farwell.
(*Crataegus punctata brevispina* Dougl.)
Crataegus caesa Ashe.
Crataegus coccinea L.
Crataegus decans Ashe.
Crataegus Dodgei Ashe.
Crataegus fallax Ashe.
Crataegus filipes Ashe.
Crataegus glareola Ashe.
Crataegus immanis Ashe.
Crataegus latisepala Ashe.
Crataegus lanta Ashe.
Crataegus lumaria Ashe.
Crataegus Michiganensis, Ashe.
Crataegus nuperia Ashe.
Crataegus obtecta Ashe.
Crataegus onusta Ashe.
Crataegus pascens Ashe.
Crataegus pastora Sargent.
Crataegus prona Ashe.
Crataegus prunifolia (Marsh.) Pers.
Crataegus pubifolia Ashe.
Crataegus pubipes Ashe.
Crataegus rotundifolia (Ehrh.) Borck.
Crataegus redolans Ashe.
Crataegus sera Sargent.
Crataegus structilis Ashe.
Crataegus tenax Ashe.
Crataegus virella Ashe.
Dasiphora fruticosa (L.) Rybd. Shrubby Cinquefoil.
(*Potentilla fruticosa* L.)
Decodon verticillatus (L.) Ell. Willow Herb.

Diervilla Diervilla (L.) Mac M. Bush Honeysuckle.
(*Diervilla trifida*, Moench.)
Empetrum nigrum L. Black Crowberry.
Epigaea repens L. Trailing Arbutus. Mayflower.
Euonymus atropurpureus Jacq. Burning Bush. Wahoo.
Euonymus obovatus Nutt. Running Strawberry Bush.
(*Enonymus Americanus obovatus* T. & G.)
Gaylussacia resinosa (Ait.) T. & G. Black Huckleberry.
Hypericum Kalmianum L. Kalm's St. John's-wort.
Hypericum prolificum L. Shrubby St. John's-wort.
Ilex verticillata (L.) A. Gray. Winter-berry.
Ilicioides mucronata (L.) Britton. Mountain Holly.
(*Nemopanthes Canadensis* DC.)
Kalmia angustifolia L. Sheep-laurel.
Kalmia glauca Ait. Swamp-laurel.
Ledum Groenlandicum Œlder. Labrador Tea.
(*Ledum latifolium* Ait.)
Lepargyraea argentea (Nutt.) Greene. Buffalo-berry.
(*Shepherdia argenta* Nutt.)
Lonicera Caprifolium L. Perfoliate Honeysuckle.
(*Lonicera grata* Ait.)
Lonicera ciliata Muhl. Fly Honeysuckle.
Lonicera coerulea L. Mountain Fly-Honeysuckle.
Lonicera dioica L. Glaucous Honeysuckle.
(*Lonicera glauca* Hill.)
Lonicera glaucescens Rydb. Douglas' Honeysuckle.
Lonicera hirsuta Eaton. Hairy Honeysuckle.
Lonicera involucrata (Richards) Banks.
Lonicera oblongifolia (Goldie) Hook. Swamp Fly-Honeysuckle.
Malus coronaria (L.) Mill. American Crab Apple.
(*Pyrus coronaria* L.)
Menispermum Canadense L. Moonseed.
Myrica cerifera L. Bayberry. Wax-myrtle.
Myrica Gale L. Sweet Gale.
Opulaster opulifolius (L.) Kuntze. Ninebark.
(*Physocarpus opulifolius* Maxim.)
Oxycoccus macrocarpus (Ait.) Pers. Large Cranberry.
(*Vaccinium macrocarpon* Ait.)
Oxycoccus Oxycoccus (L.) MacM. Small Cranberry.
(*Vaccinium Oxycoccus* L.)
Parthenocissus quinquefolia (L.) Planch. Virginia Creeper.
(*Ampelopsis quinquefolia* Michx.)
Parthenocissus quinquefolia laciniata Planch.
(*Parthenocissus vitacea* Planch.)
Parthenocissus quinquefolia hirsuta (Knerr.) T. & G.
Polycodium stamineum (L.) Greene. Deerberry.
(*Vaccinium staminium* L.)
Prunus Americana Marsh. Wild Yellow or Red Plum.
Prunus pumila L. Sand Cherry. Dwarf Cherry.
Ptelea trifoliata L. Hop-tree. Wafer-ash.
Quercus prinoides Willd. Dwarf Chestnut Oak.
Rhamnus alnifolia L'Her. Dwarf Alder.
Rhus aromatica Ait. Fragrant Sumach.
(*Rhus Canadensis* Marsh.)
Rhus copalina L. Dwarf Sumach.
Rhus glabra L. Smooth Sumach.
Rhus hirta (L.) Sudw. Staghorn Sumach.
(*Rhus typhina* L.)
Rhus radicans L. Poison Ivy.
(*Rhus Toxicodendron radicans* Marsh.)
Rhus Vernix L. Poison Sumach.
(*Rhus venenata* DC.)
Ribes Cynosbati L. Prickly Gooseberry.
Ribes floridum L'Her. Wild Black Currant.
Ribes gracile Michx. Missouri Gooseberry.

Ribes lacustre (Pers.) Poir. Swamp Gooseberry.
Ribes oxyacanthoides lacustre Pers.
Ribes oxyacanthoides L. Swamp Gooseberry.
Ribes prostratum L'Her. Fetid Currant.
Ribes rotundifolium Michx. Round-leaved Gooseberry.
Ribes rubrum L. Red Currant.
(*Ribes rubrum subglandulosum* Maxim.)
Rosa Arkansana Porter.
Rosa blanda Ait.
Rosa Carolina L. Swamp Rose.
Rosa Carolina× humilis C. F. Wheeler. A hybrid.
Rosa Engelmanni S. Wats.
Rosa humilis Marsh. Low Rose.
Rosa Sayi Schwein. Say's Rose.
Rosa setigera Michx. Prairie Rose.
Rubus Americanus (Pers.) Britton.
(*Rubus triflorus* Richards.)
Rubus Canadensis L. Low Dewberry.
(*Rubus Millspaughii* Britton.)
Rubus hispidus L. Running Swamp Dewberry.
Rubus nigrobaccus Bailey. Blackberry.
(*Rubus villosus* Ait.)
Rubus occidentalis L. Black Raspberry.
Rubus odoratus L. Purple Flowering-raspberry.
Rubus parviflorus Nutt. Salmon-Berry.
(*Rubus Nutkanus* Mocino.)
Rubus setosus Bigel. Bristly Blackberry.
Rubus strigosus Michx. Wild Red Raspberry.
Salix adenophylla Hook. Tomentose Willow.
Salix amygdaloides Anders. Peach-leaved Willow.
Salix balsamifera (Hook) Barratt. Balsam Willow.
Salix balsamifera lanceolata Bebb.
Salix balsamifera vegeta Bebb.
Salix Bebbiana Sargent. Bebb's Willow.
(*Salix rostrata* Richards.)
Salix candida Fluegge. Hoary Willow.
Salix candida× cordata Bebb. A hybrid.
Salix cordata Muhl. Heart-leaved Willow.
Salix cordata× sericea Bebb.
Salix discolor Muhl. Glaucous Willow.
Salix discolor× eriocephala Anders.
Salix discolor× prinoides (Pursh) Anders.
Salix fluviatilis Nutt. Sandbar Willow.
(*Salix longifolia* Muhl.)
Salix fragilis L.
Salix fragilis× alba Wimmer.
Salix glaucophylla Bebb. Broad-leaved Willow.
Salix glaucophylla× angustifolia Bebb.
Salix glaucophylla× brevifolia Bebb.
Salix humilis Marshall. Prairie Willow.
Salix humilis× discolor Bebb.
Salix interior× Wheeleri Rowlee.
Salix lucida Muhl. Shining Willow.
Salix myrtilloides L. Myrtle Willow.
Salix myrtilloides× pedicellaris Anders.
Salix nigra Marshall. Black Willow.
Salix nigra falcata (Pursh) Torr.
Salix petiolaris J. E. Smith. Petioled Willow.
Salix petiolaris gracilis Anders.
Salix petiolaris× candida Bebb.
Salix sericea Marsh. Silky Willow.
Salix sericea× candida Bebb.
Salix tristis Ait. Dwarf Gray Willow.
Salix viminalis L. Basket Osier.
Sambucus Canadensis L. Purple-berried Elder.

Sambucus pubens Michx. Red-berried Elder.
Smilax hispida Muhl. Green Briar.
Smilax rotundifolia L. Green Briar.
Spiraea salicifolia L. Willow-leaved Meadow-sweet.
Spiraea tomentosa L. Hardhack.
Staphylea trifolia L. Bladdernut.
Symphoricarpos occidentalis Hook. Wolfberry.
Symphoricarpos pauciflorus (Robbins) Britton. Low Snowberry.
Symphoricarpos racemosus Michx. Snowberry.
Symphoricarpos Symphoricarpos (L.) MacM. Coral-berry.
(*Symphoricarpos vulgaris* Michx.)
Vaccinium caespitosum Michx. Dwarf Bilberry.
Vaccinium Canadense Richards. Canada Blueberry.
Vaccinium corymbosum L. Swamp-blueberry.
Vaccinium corymbosum amoenum A. Gray.
Vaccinium membranaceum Dougl. Thin-leaved Bilberry.
(*Vaccinium myrtilloides* Hook.)
Vaccinium nigrum (Wood) Britton. Black Blueberry.
Vaccinium ovalifolium J. E. Smith. Oval-leaved Bilberry.
Vaccinium Pennsylvanicum Lam. Dwarf or Low-bush Blueberry.
Vaccinium Pennsylvanicum angustifolium (Ait.) A. Gray.
Vaccinium uliginosum L. Great Bilberry.
Vaccinium vacillans Kalm. Low Blueberry.
Viburnum acerifolium L. Maple-leaved Arrow-wood.
Viburnum alnifolium Marsh. Hobble-bush.
(*Viburnum lantanoides* Michx.)
Viburnum cassinoides L. Withe-rod.
Viburnum dentatum L. Arrow-wood.
Viburnum Opulus L. Cranberry-tree.
Viburnum pauciflorum Pylaie. Few-flowered Cranberry-tree.
Viburnum prunifolium L. Black Haw.
Viburnum pubescens (Ait.) Pursh.
Vitis aestivalis Michx. Summer Grape.
Vitis bicolor LeConte. Blue or Winter Grape.
Vitis cordifolia Michx. Forest Grape.
Vitis vulpina L. Riverside Grape.
(*Vitis riparia* Michx.)
Xanthoxylum Americanum Mill. Prickly Ash.
Xolisma ligustrina (L.) Britton. Privet Andromeda.

CATALOGUE.

Range of species north or south has been indicated by the following abbreviations at the right: S.—1st, 2d and 3d tiers of counties. C.—From 3d tier of counties northward to Houghton Lake region. N.—Remainder of the Lower Peninsula. L. P.—Lower Peninsula. U. P.—Upper Peninsula. Th.—Whole State so far as known. In a few instances S. E. and S. W. have been used to indicate the southeastern and southwestern parts of the Lower Peninsula. In case of rare or local species, I have given all the localities known, but for more common ones, have usually indicated only the range and relative frequency. Throughout the catalogue B. & K. refer to notes of H. C. Beardslee and Chas. A. Kofoid. W. J. B.—W. J. Beal. All plants which have been collected within ten miles of Lansing are indicated by an asterisk.*

OPHIOGLOSSACEÆ Presl. Adder's-Tongue Family.

OPHIOGLOSSUM L.

1. O. Engelmanni Prantl. Keweenaw Co., O. A. Farwell.
*2. O. vulgatum L. Adder's tongue. Keweenaw Co., O. A. Farwell; St. Clair Co., C. K. Dodge; Manistee, F. P. Daniels; Ann Arbor, C. A. Davis; due north of the Agricultural College on the Zac. Chandler farm; very likely overlooked in most places where it might be found. Rare.

BOTRYCHIUM Swartz.

2a. B. boreale (Fries.) Milde. Keweenaw Co., O. A. Farwell.
3. B. dissectum Spreng. *B. ternatum dissectum* D. C. Eaton. South Haven, Bailey; Detroit, Foerste; Alma, C. A. Davis; St. Clair Co., J. W. Stacey.
4. B. lanceolatum (S. G. Gmel.) Angstroem. Lance-leaved Grape-fern. Damp, mossy places. Lake Superior, H. Gillman; Keweenaw Co., O. A. Farwell.
5. B. Lunaria (L.) Swartz. Moonwort. "Lake Superior (Lesquereux.) and sparingly northward," Eaton in Gray's Manual; Keweenaw Co., O. A. Farwell; Harrisville, W. J. B. Rare.
6. B. matricariae (Schrank.) Spreng. Keweenaw Co., O. A. Farwell.
7. B. matricariaefolium A. Braun. Dark, wet woods. Lake Superior, D. C. Eaton; Keweenaw Co., O. A. Farwell.
8. B. obliquum Muhl. *B. ternatum obliquum* D. C. Eaton. Pastures and meadows. Fort Gratiot and S. Mich., Winch. Cat.; Flint; S. Haven, Bailey; Ann Arbor, Allmendinger Cat.; Gratiot Co.; Keweenaw Co., Detroit, O. A. Farwell; Berrien Co., H. S. Pepoon. Infrequent.
8a. B. obliquum intermedium (D. C. Eaton) Underw. *B. ternanatum australe* D. C. Eaton. Keweenaw Co., O. A. Farwell.
9. B. Onondagense Underw. Keweenaw Co., O. A. Farwell.
10. B. simplex Hitchcock. Little Grape-fern. Hillsides. Westward to Lake Superior, D. C. Eaton, Ferns of N. Am.; Keweenaw Co., O. A. Farwell; near Oscoda in 1888, the only station known in the Lower Peninsula. Rare.
11. B. tenebrosum A. A. Eaton. Keweenaw Co., O. A. Farwell.
*12. B. Virginianum (L.) Swartz. Virginia Grape-Fern. Rich woods. Variable. Common.
13. B. Virginianum gracile Pursh. With type.

OSMUNDACEÆ R. Br. Royal Fern Family.

OSMUNDA L.

*18. O. cinnamomea L. Cinnamon Fern. Swamps. Common.
*19. O. Claytoniana L. Clayton's Fern. Moist grounds, common. Th.
*20. O. regalis L. Flowering Fern. Swamps, common.

POLYPODIACEÆ R. Br. Fern Family.

ONOCLEA L.

*21. O. sensibilis L. Sensitive Fern. Wet places. Abundant. Th.

MATTEUCCIA Todaro. ONOCLEA L. in part.

*22. M. Struthiopteris (L.) Todaro. Ostrich Fern. *Onoclea Struthiopteris* (L.) Hoffm. Alluvial soil. Infrequent. Th.

WOODSIA R. Br.

23. W. alpina (Bolton) S. F. Gray. Alpine Woodsia. *W. hyperborea* R. Br. U. P., O. A. Farwell.
24. W. Ilvensis (L.) R. Br. Rusty Woodsia. N. E., Winch. Cat.; Lake Superior, Whitney Cat.; Lower Falls of the Quinnesec, E. J. Hill; Norway, C .F. Wheeler.
25. W. obtusa (Spreng.) Torr. Blunt-lobed Woodsia. Rocky places. Huron Co., C. A. Davis. U. P. Not rare.
26. W. Oregana D. C. Eaton. Oregon Woodsia. Crevices of rocks south shore of Lake Superior, Dr. Robbins in Gray's Manual.
27. W. scopulina D. C. Eaton. Rocky Mountain Woodsia. Rocks of the lower falls of the Menominee river, C. F. Wheeler.

DENNSTAEDTIA Bernh. DICKSONIA L'Her.

28. D. punctilobula (Michx.) Moore. *Dicksonia punctilobula* (Michx.) A. Gray. Petoskey, Emmet Co., Winch. Cat.; Keweenaw Co., O. A. Farwell.

FILIX Adans. CYSTOPTERIS Bernh.

*29. **F. bulbifera** (L.) Underw. *Cystopteris bulbifera* (L.) Bernh. Bulblet Fern. Moist shaded hillsides. Locally frequent. Th.

*30. **F. fragilis** (L.) Underw. *Cystopteris fragilis* (L.) Bernh. Brittle Fern. Woods and river banks. Ann Arbor, Allmend. Cat.; Flint; Hubbardston; Alma, C. A. Davis; northward to Lake Superior, Whitney. Th.

31. **F. fragilis tenue** (Michx.) *Cystopteris fragilis dentata* Hook. Keweenaw Co., and Ypsilanti, O. A. Farwell.

32. **F. montana** (Lam.) Underw. Mountain Cystopteris. *Cystopteris montana* (Lam.) Bernh. North shore of Lake Superior, Britton & Brown.

POLYSTICHUM Roth. ASPIDIUM in part.

33. **P. acrostichoides** (Michx.) Schott. Christmas Fern. *Aspidium acrostichoides* Sw. Shady ground. Th.

34. **P. Braunii** (Spenner) Fée. Braun's Holly-fern. *Aspidium aculeatum Braunii* Doell. Ontonagon Peninsula, Eaton's Ferns of N. A.; Keweenaw Co., O. A. Farwell.

35. **P. Lonchitis** (L.) Roth. Holly-Fern. *Aspidium Lonchitis* Sw. Woods south shore of Lake Superior, Prof. Whitney; Keweenaw Co., O. A. Farwell.

DRYOPTERIS Adans. ASPIDIUM in part.

36. **D. Boottii** (Tuckerm.) Underw. Boott's Shield-fern. *Aspidium Boottii* Tuckerm. Alder thickets. Hubbardston; Ann Arbor, Winch. Cat.; Norway, C. F. Wheeler; Keweenaw Co., O. A. Farwell.

*37. **D. cristata** (L.) A. Gray. Crested Shield-fern. *Aspidium cristatum* Sw. Swamps. Frequent. Th.

38. **D. cristata Clintoniana** (D. C. Eaton) Underw. *Aspidium cristatum Clintonianum* D. C. Eaton. Ann Arbor, Allmendinger Cat.; Hubbardston; found at Lake Nipigon, Ont., Macoun; Alma, C. A. Davis. Infrequent. Th.

39. **D. Filix-Mas** (L.) Schott. Male Fern. *Aspidium Filix-mas* Sw. Rocky woods. Keweenaw Peninsula, Lake Superior, A. Gray.

40. **D. fragrans** (L.) Schott. Fragrant Shield-fern. *Aspidium fragrans* Sw. Crevices of shaded cliffs. Isle Royale, and Keweenaw Peninsula, Dr. Lyons.

*41. **D. Goldieana** (Hook.) A. Gray. Goldie's Fern. *Aspidium Goldieanum* Hook. Moist woods. Flint; Hubbardston, C. F. Wheeler; Owosso, G. H. Hicks; Black River, St. Clair Co., C. K. Dodge; Alma, C. A. Davis.

*42. **D. marginalis** (L.) A. Gray. Evergreen Wood-fern. *Aspidium marginale* Sw. Hillsides in rich woods. S. Haven, Bailey; Flint; Hubbardston and northward. Frequent. Th.

*43. **D. Noveboracensis** (L.) A. Gray. New York Fern. *Aspidium Noveboracense* Sw. Swamps. Ann Arbor, Allmendinger Cat.; Macomb Co.; Hubbardston; Gore Bay, Manitoulin Island, J. Bell. Canadian Cat. L. P.

44. **D. simulata** Davenport. Belle Isle, O. A. Farwell.

*45. **D. spinulosa** (Retz) Kuntze. Spinulose Shield-fern. *Aspidium spinulosum* Sw. Shady woods. Flint; Hubbardston; L. Superior, D. C. Eaton, Ferns of N. A. Th.

46. **D. spinulosa dilatata** (Hoffm.) Underw. *Aspidium spinulosum* var. *dilatatum* Hook. Woods. Ann Arbor, Allmendinger Cat.; Macomb Co.; Flint; Hubbardston; Petoskey, and northward. Th.

*47. **D. spinulosa intermedia** (Muhl.) Underw. *Aspidium spinulosum* var. *intermedium* D. C. Eaton. Woods. Common. Th.

*48. **D. Thelypteris** (L.) A. Gray. Marsh Shield-fern. *Aspidium Thelypteris* Sw. Swamps. Frequent. Th.

PHEGOPTERIS Fée.

*49. **P. Dryopteris** (L.) Fée. Oak-fern. Open woods. Frequent north of lat. 43°, especially under pines. Th.

*50. **P. hexagonoptera** (Michx.) Fée. Broad Beech-fern. Beech woods. Flint; Hubbardston; Ann Arbor, Allmendinger Cat.; Keweenaw Co., O. A. Farwell. Frequent. Th.

*51. P. Phegopteris (L.) Underw. Long Beech-fern. *P. polypodioides* Fée. Isle Royale, Dr. A. B. Lyons; Keweenaw Co., O. A. Farwell; Frankfort; Inland, Grand Traverse Co.; rocks, Grand Ledge. Abundant northward. Th.

WOODWARDIA J. E. Smith.

52. W. areolata (L.) Moore. Net-veined Chain-fern. *W. angustifolia* J. E. Smith. South Haven, L. H. Bailey in 1880; near Black River, St. Clair Co., C. K. Dodge.
*53. W. Virginica (L.) J. E. Smith. Virginia Chain-fern. Marshes, Marquette Co., Burt. MS. Cat.; Westville, Montcalm Co.; Flint; So. Haven, Bailey; Hubbardston, C. F. Wheeler; Ann Arbor, Allmendinger Cat.; near Port Huron, C. K. Dodge. Th.

CAMPTOSORUS Link.

54. C. rhizophyllus (L.) Link. Walking-fern. Keweenaw Co., O. A. Farwell; Alpena, V. M. Spalding, C. F. Wheeler; Norway on the northeast exposure of quartz rock, C. F. Wheeler. Rare.

ASPLENIUM L.

*55. A. acrostichoides Sw. Silvery Spleenwort. *A. thelypteroides* Michx. Rich woods. Frequent. Th.
*56. A. angustifolium Michx. Narrow-leaved Spleenwort. Rich woods, S. W., Wright Cat.; Ann Arbor, rare, Allmendinger Cat.; Flint; S. Haven, Bailey; Ionia Co. Infrequent. C. & S.
*57. A. Filix-foemina (L.) Bernh. Lady Fern. Moist woods. Exceedingly variable. Common. Th.
58. A. montanum Willd. Mountain Spleenwort. Near Clifton, Keweenaw Co., O. A. Farwell.
59. A. platyneuron (L.) Oakes. Ebony Spleenwort. *A. ebeneum* Ait. Allegan, Miss Josie A. Williams. Only station known in the State.
60. A. Rutra-muraria L. Wall Rue Spleenwoort. N. E., Winch. Cat.; Keweenaw Co., O. A. Farwell. Rare.
61. A. Trichomanes L. Maidenhair Spleenwort. N. E., Winch. Cat.; Keweenaw Point; Isle Royale, Porter. Infrequent. N. & U. P.

ADIANTUM L.

*62. A. pedatum L. Maiden-hair Fern. Rich, moist woods. One of the most beautiful ferns. Common. Th.

PTERIDIUM Scop. PETRIS L. in part.

*63. Pteridium aquilinum (L.) Kuhn. Common Brake. *Pteris aquilina* L. Open ground. Common. Th.

CRYPTOGRAMMA R. Brown.

64. C. acrostichoides R. Br. American Rock-brake. Isle Royale, Gray's Manual; Caribou Island, Porter. U. P.
65. C. Stelleri (Gmel.) Prantl. Slender Cliff-brake. *Pellaea Stelleri* (Gmel.) Walt. Louse Isle, Winch. Cat.; Pictured Rocks, G. H. Hicks.

PELLAEA Link.

66. P. atropurpurea (L.) Link. Purple Cliff-brake. Norway, C. F. Wheeler. Shaded limestone rocks. N. E. Winch. Cat.

POLYPODIUM L.

*67. P. vulgare L. Common Polypody. Dry banks and rocks. Grand Ledge, W. J. B.; Hubbardston, C. F. Wheeler. Rarely seen in L. P., but very common in U. P.

EQUISETACEÆ Michx. Horsetail Family.

EQUISETUM L.

*68. E. arvense L. Common Horsetail. Damp sandy grounds. Common. Th.
69. E. arvense campestre Schultz. Keweenaw Co., O. A. Farwell.
70. E. arvense riparium Milde. Keweenaw Co., O. A. Farwell.
*71. E. fluviatile L. Swamp Horsetail. *E. limosum* L. In shallow water. Common. Th.
72. E. fluviatile polystachyon (Bruckner) Prayer. Keweenaw Co., O. A. Farwell.
*73. E. hyemale L. Scouring-Rush. Wet banks, common. Th.
*74. E. hyemale ramigerum A. Br. Keweenaw Co., O. A. Farwell.
75. E. laevigatum A. Br. Smooth Scouring-rush. Port Huron, C. K. Dodge; Manistee, F. P. Daniels.
76. E. littorale Kuehl. Shore Horsetail. New Buffalo; Oscoda; Keweenaw Co., O. A. Farwell.
77. E. palustre L. Marsh Horsetail. Lake Huron to the Arctic Sea. (Hook, Fl.) U. P.
*78. E. pratense Ehrh. Thicket Horsetail. Macomb Co.; Flint; to Pine Lake, Emmet Co., Winch. Cat.; Keweenaw Co., O. A. Farwell. Th.
*79. E. robustum A. Br. Stout Scouring-rush. Port Huron, C. K. Dodge; Morenci, W. J. B.; Manistee, F. P. Daniels.
80. E. scirpoides Michx. Sedge-like Equisetum. Macomb Co.; Flint; Hubbardston; Petoskey; to L. Superior, Whitney Cat., O. A. Farwell. Th.
*81. E. sylvaticum L. Wood Horse-tail. Ann Arbor, rare, Allmend. Cat.; Clinton Co.; Montcalm Co.; Flint; and northward where it is common. Th.
82. E. variegatum Schleich. Variegated Equisetum. S. E., Winch. Cat.; Hubbardston; Muir; Macomb Co.; Flint; sand dunes at the head of Little Traverse Bay; Drummond's I., Port Huron, C. K. Dodge; Detroit and Keweenaw Co., O. A. Farwell. Infrequent. Th.

LYCOPODIACEÆ Michx. Club-Moss Family.

LYCOPODIUM L.

83. L. alpinum L. Alpine Club-moss. Keweenaw Co., O. A. Farwell.
84. L. annotinum L. Stiff Club-moss. Woods. Petoskey to L. Superior; St. Clair Co., C. K. Dodge. Common.
85. L. Chamaecyparissus A. B. *L. complanatum Chamaecyparissus* D. C. Eaton. Tp. of Arthur, Clare Co., May 15, 1849, G. H. Cannon; Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels.
*86. L. clavatum L. Common Club-moss. Dry woods. Common. Th. N. & U. P.
87. L. clavatum monostachyon Hook. Keweenaw Co., O. A. Farwell.
*88. L. complanatum L. Ground-Pine. Bangor, Van Buren Co., Bailey; shore of Pine Lake; Keweenaw Co., O. A. Farwell; Woodward Lake, Ionia Co.; Flint; Macomb Co., Stanton, and northward. Th.
89. L. inundatum L. Bog Club-moss. Drummond's I.; Willow River, Huron Co.; Sugar Is., Winch. Cat.; Keweenaw Co., O. A. Farwell. Th.
*90. L. lucidulum Michx. Shining Club-moss. Moist woods. Frequent. Th.
*91. L. obscurum L. Ground Pine. Moist woods.
92. L. obscurum dendroideum D. C. Eaton. Macomb Co., Cooley; Keweenaw Co., O. A. Farwell, near Port Huron, C. K. Dodge; Oscoda.
93. L. porophilum Lloyd & Underw. Keweenaw Co., O. A. Farwell.
94. L. sabinaefolium Willd. Cedar-like Club-Moss. Keweenaw Co., O. A. Farwell. Rare.
95. L. Selago L. Fir Club-moss. Lake Superior, Gray's Manual; Keweenaw Co., O. A. Farwell. Infrequent U. P.

SELAGINELLACEÆ Underw.

SELAGINELLA Beauv.

*96. S. apus (L.) Spring. Creeping Selaginella. Ann Arbor, Allmendinger Cat.; Hubbardston, Grand Ledge, C. F. Wheeler; Orion, O. A. Farwell; Port Huron, W. S. Cooper and C. K. Dodge; Alma and Ann Arbor, C. A. Davis. Infrequent.

*97. S. rupestris (L.) Spring. Rock Selaginella. Flint; Woodward Lake; rare in L. P.; but common in U. P. Park Lake, Clinton Co., C. F. Wheeler; Oscoda, Grayling, Harrison, W. J. B.; Keweenaw Co., O. A. Farwell. Th.

98. S. selaginoides (L.) Link. Low Selaginella. *S. spinosa* Beauv. Isle Royale, Dr. A. B. Lyons; Keweenaw Co., O. A. Farwell. U. P.

ISOETACEÆ Underw. Quillwort Family.

ISOETES L.

99. L. echinospora Braunii (Durieu) Engelm. Braun's Quillwort. Gray's Manual.

100. L. lacustris L. Lake Quillwort. River St. Marie, head of ship canal, T. C. Porter; St. Clair River, W. S. Cooper.

PINACEÆ Lindl. Coniferæ in part. Pine Family.

PINUS L.

101. P. divaricata (Ait.) Gord. Gray Pine. Jack Pine. *P. Banksiana* Lambert In the western part of the state noticed as far south as Grand Haven. Occurs at south end of Lake Michigan in Indiana; Sand Pt., Port Austin, Saginaw Bay, and northward along the shore of L. Huron, not common, Winch. Cat. In the central part of the State this pine is first seen in the northern part of Clare Co., where it is common in groves on sandy barrens. Trees sometimes grow 50 to 70 feet in height and rarely have been measured with a diameter of two feet.

102. P. resinosa Ait. Red Pine. Norway Pine. Dry woods. First noticed in Isabella Co., in center of the L. P.; very abundant in Clare Co., and northward. "Frequent on low, sandy plains in U. P., where it forms orchard-like groves," Whitney. Usually 100 to 110 feet high. On the east side of the State it extends southward to Port Huron, C. K. Dodge; Gratiot Co., C. A. Davis.

*103. P. Strobus L. White Pine. Weymouth Pine. C. & N. W.

LARIX Adans.

*104. P. laricina (DuRoi) Koch. American Larch. Tamarack. *L. Americana* Michx. Swamps. A slender tree southward, but sometimes reaching 100 feet in height northward, where it is abundant. Th.

PICEA Link.

105. P. brevifolia Peck. Keweenaw Co., O. A. Farwell.

106. P. Canadensis (Mill.) B. S. P. White Spruce. *P. alba* Link. From Ludington to Alcona Co., W. J. B.; Bay Co., G. M. Bradford; Petoskey and northward. N. & U. P. Common.

*107. P. Mariana (Mill.) B. S. P. Black Spruce. *P. nigra* Link. Frequent northward, and occasional south—a small tree in spagnous swamps. Th.

TSUGA Carriere.

*108. T. Canadensis (L.) Carriere. Hemlock. Grand Ledge. Rarely seen south of latitude 43° except in the west, and scarce on the Huron shore, but common on the east shore of Lake Michigan; Berrien Co., H. S. Pepoon, and from the central part of the State northward. Very abundant and of great size in Emmet Co. Th. except in S. E.

ABIES Juss.

109. A. balsamea (L.) Miller. Balsam or Balm-of-Gilead Fir. Frequent in the northern part of Clare Co.; Bay Co., G. M. Bradford; abundant at Petoskey, and northward. N. C. & U. P.

THUJA L. THUYA.

110. T. occidentalis L. Arbor Vitae. White Cedar. Lake Orion, Whitmore Lake. C. A. Davis; Montcalm Co., and northward.

JUNIPERUS L.

*111. J. communis L. Common Juniper. Occasional throughout, along the highest grounds.

112. J. nana Willd. *J. Sibirica* Burgsd. Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels; common in Jackson, Livingston and Washtenaw Counties, C. A. Davis.

113. J. Sabina L. Shrubby Red Cedar. *J. Sabina procumbens* Pursh. Little Traverse Bay, W. J. B.; Keweenaw Co., and Rochester, O. A. Farwell. N. & U. P.

*114. J. Virginiana L. Red Cedar. Savin. Bluffs and sterile soil. Throughout, but scarce.

TAXACEÆ Lindl. Yew Family.

TAXUS L.

115. T. Canadensis Marsh. American Yew. Ground Hemlock. *T. minor* (Michx.) Britton. Moist woods in the shade of evergreens. Along the east shore of Lake Michigan as far south as South Haven, L. H. Bailey; Macomb Co., where it is rare. St. Clair Co., C. K. Dodge. Common northward.

TYPHACEÆ J. St. Hil. Cat-tail Family.

TYPHA L.

116. T. angustifolia L. Narrow-leaved Cat-tail Flag. Detroit, Dr. A. B. Lyons; Port Huron, C. K. Dodge; Ann Arbor and Tuscola Co., C. A. Davis; Bay Co., G. M. Bradford; S. E. Rare.

*117. T. latifolia L. Common Cat-tail. Borders of streams and ponds. Common. Th.

SPARGANIACEÆ Agardh. Bur-Reed Family.

SPARGANIUM L.

118. S. androcladum (Engelm.) Morong. Branching Bur-reed. *S. simplex androcladum* Engelm. Ann Arbor, Allmendinger Cat.; Flint, Dr. Clark; Macomb Co.; St. Clair Co., C. K. Dodge; Bay Co., G. M. Bradford.

*119. S. eurycarpum Engelm. Broad Fruited Bur-reed. Borders of ponds. Frequent. C. & S.

120. S. minimum Fries. Small Bur-reed. Hubbardston; Manistee, E. J. Hill; Macomb Co.; Homestead, Benzie Co.; Keweenaw Co., O. A. Farwell. Rare.

*121. S. simplex Hudson. Simple-stemmed Bur-reed. Fruitport, E. J. Hill; Oscoda; Keweenaw Co., O. A. Farwell; Black River, B. & K. Common. Th.

122. S. simplex angustifolium (Michx.) Engelm. Isle Royale, Whitney Cat.; Keweenaw Co., O. A. Farwell; N. shore of Lake Superior, Agassiz.

NAIADACEÆ Lindl. Pondweed Family.

POTAMOGETON L.

123. P. alpinus Balbis. Northern Pondweed. *P. rufescens* Schrad. N. & U. P.

*124. P. amplifolius Tuckerman. Large-leaved Pondweed. Maple River; Macomb Co.; Pere Marquette River at Ludington, E. J. Hill; St. Clair Co., C. K. Dodge.

125. P. diversifolius Raf. Rafinesque's Pondweed. *P. hybridus* Michx. Dr. A. B. Lyons; Keweenaw Co., O. A. Farwell.

126. P. filiformis Pers. Filiform Pondweed. Belle Isle, O. A. Farwell.

*127. P. foliosus Raf. Leafy Pondweed. *P. pauciflorus* Pursh. Hubbardston, Grand Rapids, Flint, Macomb Co., and northward. Typical forms occur at Manistee and Frankfort, E. J. Hill. Th.

128. P. foliosus Niagarensis. (Tuckerman) Morong. In running water, Hubbardston; Macomb Co., and along the great lakes.

*129. P. Friesii Ruprecht. Fries' Pondweed. *P. mucronatus* Schrad. Crooked River, Cheboygan Co., and common at Manistee and Frankfort, E. J. Hill; Oscoda. St. Mary's River, Sault de Ste Marie; Detroit, O. A. Farwell.

*130. P. heterophyllus Schreb. Various-leafed Pondweed. Th.

131. P. heterophyllus graminifolius (Fries) Morong. *P. gramineus graminifolius* Flint, Macomb Co.; St. Clair Co., C. K. Dodge, and northward.

132. P. heterophyllus longipedunculatus Morong. Woodward Lake, Ionia Co., Fries. Detroit, O. A. Farwell; Bear Lake, Van Buren Co., E. J. Hill; Black Lake, Cheboygan Co., B. & K.

133. P. Hillii Morong. Hill's Pondweed. Marquette Co., E. J. Hill; St. Clair Co., C. K. Dodge; Sturgis, F. P. Daniels.

134. P. Illinoensis Morong. Manistee, F. P. Daniels.

135. P. interruptus Kitaibel. Interrupted Pondweed. Manistee, E. J. Hill; Port Huron, C. K. Dodge.

136. P. lateralis Morong. Opposite-leafed Pondweed. Bear Lake, Van Buren Co., E. J. Hill; St. Clair Lake & River, C. K. Dodge. Rare.

*137. P. lonchites Tuckerman. Long-leafed Pondweed. *P. fluitans* Roth. Hubbardston; Macomb Co.; Detroit River, D. H. Campbell; Portage Lake, Crawford Co.; Manistee, F. P. Daniels. Infrequent. L. P.

*138. P. lucens L. Shining Pondweed. Muskegon River, near Houghton Lake; Flint; S. Mich., Wright Cat.; Detroit River, D. H. Campbell; Lake Superior, Agassiz. Th.

139. P. marinus L. Crystal Lake near Frankfort, Benzie Co., E. J. Hill; St. Clair Co., C. K. Dodge.

*140. P. natans L. Common Floating Pondweed. Ponds. Frequent. Th.

141. P. Nuttallii Cham. & Sch. *P. Pennsylvanicus* Cham. Ionia Co.; Macomb Co., Dr. D. Cooley; Fruitport, E. J. Hill; Keweenaw Co., O. A. Farwell. Th.

142. P. obtusifolius Mertens & Koch. Blunt-leafed Pondweed. "Floating in Gratiot Lake, N. Mich." Gray's Manual. St. Clair Co., J. W. Stacey.

*143. P. pectinatus L. Fennel-leafed Pondweed. Frequent. Th.

*144. P. perfoliatus L. Clasping-leafed Pondweed. Detroit, O. A. Farwell; Ingham Co., C. F. Wheeler, and northward. Common.

146. P. perfoliatus Richardsonii A. Bennett. *P. perfolatus lanceolatus* Robbins. Detroit, O. A. Farwell; Bay Co., G. M. Bradford.

*147. P. praelongus Wulfen. White-stemmed Pondweed. Maple River; Pine Lake, Ingham Co.; Black Lake, Cheboygan Co. Common. Th.

148. P. pusillus L. Small Pondweed. Manistee Lake, E. J. Hill; Detroit River, D. H. Campbell; Manistee, F. P. Daniels; St. Clair Lake and River, C. K. Dodge.

*149. P. Robbinsii Oakes. Robbins' Pondweed. North shore of Lake Superior; Pine Lake, seven miles N. E. of the Agricultural College, the only station known in the L. P., C. F. Wheeler.

151. P. rutilis Wolfg. Slender Pondweed. Detroit River, between Peche Isle and Belle Isle, C. F. Wheeler; Round Lake, near Jackson; St. Clair Lake & River, C. K. Dodge.

152. P. spathulaeformis (Robbins) Morong. Spatulate-leafed Pondweed. New Baltimore, Lake St. Clair, A. J. Pieters, J. W. Stacey.

153. P. Spirillus Tuckerman. Spiral Pondweed. "Lake Superior," A. Gray's Manual; Keweenaw Co., O. A. Farwell; St. Clair Lake and River, C. K. Dodge.

154. P. Vaseyi Robbins. St. Clair Lake and River, C. K. Dodge; Algonac, W. S. Cooper.

*155. P. Zizii Roth. Ziz's Pondweed. *P. angustifolius* Berch & Presl. Crystal Lake, near Frankfort; Bear Lake, Manistee Co., E. J. Hill; Crystal Lake, Montcalm Co.; Woodward Lake, Ionia Co.; Detroit River, D. H. Campbell. Infrequent.

*156. P. zosteraefolius Schum. Eel-grass. Pondweed. Common. Th.

ZANNICHELLIA L.

157. Z. palustris L. Zannichellia. Dr. A. B. Lyons, Detroit; Bay City, G. M. Bradford. Rare.

NAIAS L.

*158. N. flexilis (Willd.) Rost. & Schmidt. Slender Naias. Ponds. Frequent. Th.

*159. N. flexilis robusta Morong. South Haven, Park Lake, Clinton Co., Bailey; St. Clair Co., C. K. Dodge.

160. N. marina L. Large Naias. Flint, Dr. D. Clark.

SCHEUCHZERIACEÆ Agardh. Arrow-Grass Family.

TRIGLOCHIN L.

161. T. maritima L. Seaside Arrow-grass. Shores of Great Lakes; borders of deer licks in interior; Macomb Co.; S. Mich., Wright Cat.; Hubbardston, Petoskey, C. F. Wheeler; Algonac, W. S. Cooper.
162. T. palustris L. Marsh Arrow-grass. Marshes. Frequent. Th.

SCHEUCHZERIA L.

163. S. palustris L. S. W. Wright Cat.; Macomb Co., Cooley; Hubbardston; Montcalm Co.; Keweenaw Co., O. A. Farwell. Infrequent.

ALISMACEÆ D C. Water-Plantain Family.

ALISMA L. Water-Plantain.

*164. A. plantago-aquatica L. Water Plantain. Shallow water. Common. Th.
165. A. tenellum Mart. Dwarf Water Plantain. *Echinodorus parvulus* Engelm. A. B. Lyons; Gray's Manual, 6th edition.

SAGITTARIA L.

*166. S. arifolia Nutt. Arum-leaved Arrow-head. Alma, C. A. Davis; St. Clair Co., C. K. Dodge; Belle Isle, O. A. Farwell.
167. S. cristata Engelm. Crested Arrow-head. St. Clair Co., and Sarnia, C. K. Dodge.
*168. S. graminea Michx. Grass-leaved Sagittaria. Dr. A. B. Lyons, Detroit; Park Lake, Clinton Co., L. H. Bailey; Belle Isle, O. A. Farwell. S.
*169. S. heterophylla Pursh. Elk Rapids, Winch. Cat.; Detroit, D. H. Campbell; Alma, Chas. A. Davis. Th.
*170. S. latifolia Willd. Broad-leaved Arrow-head. Wet places. Th. *S. variabilis* Engelm.
171. S. rigida Pursh. *S. heterophylla rigida* Engelm. Fruitport, E. J. Hill; Great Lakes.

VALLISNERIACEÆ Dumort. Tape-Grass Family

PHILOTRIA Raf. (Elodea Michx.)

*172. P. Canadensis (Michx.) Britton. Water-weed. *Elodea Canadensis* Michx. Slow streams. Common. Th.

VALLISNERIA L.

*173. V. spiralis L. Tape-grass. Eel-grass. Ann Arbor, Winch. Cat.; S. Haven, Bailey; Hubbardston, to Petoskey; Lake Superior, Agassiz. Slow streams. Common in Grand River and tributaries. Th.

GRAMINEÆ Juss. Grass Family.

ANDROPOGON L.

*174. A. furcatus Muhl. Forked Beard-grass. Light soil. Common. Th.
*175. A. scoparius Michx. Broom Beard-grass. Sterile soil. Frequent, L. P.
176. A. scoparius multiramea Hack. Manistee, F. P. Daniels.

SORGHASTRUM Nash. CRYSOPOGON Benth. Not Trin.

*177. S. avenaceum (Michx.) Nash. Indian Grass. *Chrysopogon avenaceus* Benth. Dry banks and sandy fields. Variable. Infrequent. C. & S.

PASPALUM L.

178. P. laeve Michx. S. W. H. S. Pepoon. Rare.
179. P. Muhlenbergii Nash. Detroit, O. A. Farwell.
*180. P. setaceum Michx. Detroit, O. A. Farwell.

SYNTHERISMA Walt. PANICUM L. in part.

*181. S. linearis (Krock.) Nash. Small Crab-grass. *Panicum glabrum* Gaudin. Waste places. Common.
*182. S. sanguinalis (L.) Dulac. Large Crab-grass. *Panicum sanguinale* L. Waste places, gardens and fields. A bad weed, difficult to dig up or pull out; grows quickly and is perhaps the worst weed we have in gardens, at least in some localities. S. linearis grows in the same situations but is smaller and easily eradicated. Th.

ECHINOCHLOA Beauv. PANICUM L. in part.

*183. E. Crus-galli (L.) Beauv. Barnyard-grass. *Panicum Crus-galli* L. Low grounds. Variable. Common.
*184. E. Walteri (Pursh.) Nash. Salt-marsh Cockspur Grass. *P. Crus-galli hispidum* Ell. Low river bank near Hubbardston where it appears to be indigenous, C. F. Wheeler. Common along streams. L. P.

PANICUM L.

185. P. boreale Nash. Northern Panicum. Grand Rapids, Miss E. J. Cole; Traverse City, C. F. Wheeler; Keweenaw Co., O. A. Farwell.
*186. P. capillare L. Old-witch Grass. Sandy soil, fields. The spreading panicle is easily broken off and blown about by the wind. Common. Th.
*187. P. clandestinum L. Hispid Panicum. Low grounds. Macomb Co.; Flint, Hubbardston; Bay Co., G. M. Bradford. Rare.
*188. P. Columbianum Scribn. Belle Isle, O. A. Farwell.
*189. P. commutatum Schultes. Variable Panicum. Detroit, O. A. Farwell; St. Clair Co., J. W. Stacey. Dry woods.
*190. P. depauperatum Muhl. Capac, W. S. Cooper. Starved Panicum. Dry woods and barrens. Common. Th.
*191. P. dichotomum L. Forked Panicum. Dry or low grounds. A very common and exceedingly variable grass. Some forms seem distinct enough to be considered species. The most common forms are:
*192. P. dichotomum commune S. Wats.
*193. P. dichotomum fasciculatum S. Wats.
*194. P. dichotomum gracile S. Wats. Th.
195. P. agrostoides Muhl. Long Panicum. Specimen in the College Herb., collected by Dr. Cooley, at Washington, Macomb Co. Only locality known in the State.
196. P. flexile (Gattinger) Scrib. Wiry Panicum. Detroit, O. A. Farwell; Port Huron, C. K. Dodge; Manchester, C. F. Wheeler.
*196a. P. macrocarpon Le Conte. Detroit, O. A. Farwell.
*197. P. implicatum Scribn. Keweenaw Co., O. A. Farwell.
*198. P. maculatum Ashe. "Formerly included in P. dichotomum." St. Clair Co., C. K. Dodge.
199. P. polyanthes Schul. Small-fruited Panicum. *P. microcarpon* Muhl. Washington, Macomb Co., Dr. Cooley.
200. P. miliaceum L. Millet. Detroit and Keweenaw Co., O. A. Farwell.
201. P. nitidum Lam. Shining Panicum. Washington, Macomb Co., Dr. Cooley; Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels.
202. P. proliferum Lam. Manistee, F. P. Daniels.
*203. P. Porterianum Nash. Porters Panicum. *P. latifolium* Walt. Rich woods. Northward to Oscoda Co. Common. L. P.
*204. P. pubescens Lam. Hairy Panicum. Common. C. & S.
*205. P. Scribnerianum Nash. Scribner's Panicum. *P. scoparium* S. Wats. Not Lam. Hubbardston. Along the R. R. between St. Johns and Muir, northward to Baldwin; Bay Co., G. M. Bradford; Detroit, O. A. Farwell. Frequent in light sandy soil. Th.
*206. P. sphaerocarpon Ell. Round-fruited Panicum. *P. microcarpon sphaero-*

carpon (Ell.) Beal. Jackson, S. H. Camp; Detroit, O. A. Farwell; St. Clair Co., J. W. Stacey; Bay Co., G. M. Bradford.

*207. P. virgatum L. Tall Smooth Panicum. S. Mich., Wright Cat.; Ionia, Flint; Sturgis, F. P. Daniels. A tall coarse grass along rivers. Infrequent. Th.

208. P. xanthophysum A. Gray. Slender Panicum. N. & U. P. Grand Traverse Co., Indian River and Black Lake; Cheboygan Co., C. F. Wheeler; Alma, C. A. Davis; Manistee, F. P. Daniels; Northward to Keweenaw Co., O. A. Farwell.

CHAETOCHLOA Scribn. SETARIA Beauv.

*209. C. glauca (L.) Scrib. Yellow Foxtail. *Setaria glauca* Beauv. Cultivated Fields. Common. Th.

*210. C. Italica (L.) Scrib. Hungarian Grass, Italian or German Millet. *Setaria Italica* R. & S. Persisting after cultivation.

211. C. verticillata (L.) Scribn. *Setaria verticillata* Beauv. Ypsilanti, O. A. Farwell.

*212. C. viridis (L.) Scrib. Green Foxtail. *Setaria viridis* Beauv. Cultivated fields. Common. Th.

CENCHRUS L.

*213. C. tribuloides L. Sand-bur. Bur-grass. This bad weed has begun to occupy the S. part of the State, as far north as Bay City. Not as yet troublesome in fields, but likely to become so on sandy farms.

ZIZANIA L.

214. Z. aquatica L. Indian Rice. Water Oats. Borders of large streams and lakes. Yields an edible grain, gathered by the Indians, and greedily eaten by wild ducks which haunt lakes and rivers during its ripening in innumerable numbers. Th.

HOMALOCENCHRUS Mieg. (LEERSIA Sw.)

*215. H. oryzoides (L.) Poll. Rice Cut-grass. *L. oryzoides* Sm. Ditches and wet lands. Frequent. Th.

*216. H. Virginicus (Willd.) Britton. White Grass. *L. Virginica* Willd. Wet woods and river banks. Th.

PHALARIS L.

*217. P. arundinacea L. Reed Canary-grass. Borders of streams. "var. picta, the leaves striped with white, is the familiar ribbon-grass of the garden," Gray's Manual. Ann Arbor, Prof. M. W. Harrington; S. Mich., Winch. Cat.; Macomb Co.; Flint; Hubbardston; northward to L. Superior, Whitney Cat. Th.

218. P. Canariensis L. Canary-grass. Occasionally found in waste places.

ANTHOXANTHUM L.

*219. A. odoratum L. Sweet Vernal-grass. Ionia; Grand Rapids, Coleman Cat.; well established in the College lawn. Rare.

SAVASTANA Schrank. HIEROCHLOE Gmel.

220. S. odorata (L.) Scribn. Holy Grass. *Hierochloe borealis* R. & S. Not confined to the shores of the Great Lakes; rarely found in the central part of the State. Ionia, Hubbardston, C. F. Wheeler; Macomb Co.; Flint; S. E., Winch. Cat.; Ann Arbor, C. A. Davis. More common northward. Th.

ARISTIDA L.

221. A. gracilis Ell. Slender Aristida. Port Huron, C. K. Dodge; near Ann Arbor, C. A. Davis.

222. A. purpurascens Poir. Sturgis, F. P. Daniels.

STIPA L. Feather-grass.

223. S. avenacea L. Black Oat-grass. S. W., Wright Cat.; South Haven, Bailey; Baldwin; Dundee; Cass Co.; Point Aux Pins, Macoun Can. Cat. Infrequent. Th.
*224. S. spartea Trin. Porcupine-grass. Dry plains. S. Mich., Wright Cat.; Macomb Co.; shore of Woodward Lake, Ionia Co.; hills along Grand River near Ionia; Ann Arbor, Tuscola Co., C. A. Davis.

ORYZOPSIS Michx.

*225. O. asperifolia Michx. White-grained Mountain Rice. Hillsides. Common.
226. O. juncea (Michx.) B. S. P. Slender Mountain Rice. *O. Canadensis* Torr. Sterile soil. S. E., Wright Cat.; Hubbardston; Montcalm Co., C. A. Davis; Macomb Co.; common in Clare Co.; frequent from Ionia northward. Th.
*227. O. melanocarpa Muhl. Black-fruited Mountain Rice. Woods. S. Mich., Wright Cat.; S. Haven; Macomb Co.; Flint; Hubbardston. Infrequent. C. & S.

MILIUM L.

*228. M. effusum L. Tall Millet-grass. Woods. Frequent. Th.

MUHLENBERGIA Schreber.

*229. M. diffusa Willd. Nimble Will. Open woods, becoming frequent around dwellings. S. Mich., Wright Cat.; So. Haven, Bailey; Flint; Hubbardston; C. F. Wheeler; Ann Arbor, C. A. Davis. C. & S.
*230. M. Mexicana (L.) Trin. Meadow Muhlenbergia. Low grounds. Variable. Frequent. Th.
*231. M. racemosa (Michx.) B. S. P. Marsh Muhlenbergia. *M. glomerata* Trin. Marshes. Infrequent. Th.
*232. M. sylvatica Torr. Wood Muhlenbergia. Hubbardston; Flint; Lake Superior, Agassiz. Frequent. Th.
*233. M. tenuiflora (Willd.) B. S. P. Slender Muhlenbergia. *M. Willdenovii* Trin. Flint; Macomb Co.; S. Mich., Wright Cat.; Muskegon, C. F. Wheeler. Rare. C. S.

BRACHYELYTRUM Beauv.

*234. B. erectum (Schreb.) Beauv. *B. aristatum* Beauv. Woods. Frequent.

PHLEUM L.

235. P. alpinum L. Mountain Timothy. L. Superior, Whitney Cat.; Keweenaw Co., O. A. Farwell.
*236. P. pratense L. Timothy. Herd's-Grass. Meadows everywhere.

ALOPECURUS L.

*237. A. geniculatus L. Marsh Foxtail. *A. geniculatus aristulatus* Torr. S. Mich., Wright Cat.; Macomb Co.; Bay Co., G. M. Bradford. Th.
*238. A. pratensis L. Meadow Foxtail. Cultivated from Europe and escaped.

SPOROBOLUS R. Br.

239. S. brevifolius (Nutt.) Nash. Keweenaw Co., O. A. Farwell.
240. S. cryptandrus (Torr.) A. Gray. Sand Dropseed. S. Mich., Winch. Cat.; Point Edward, River St. Clair, Macoun; Bay City, W. J. B.; shore of Gull Lake, Augusta, Kalamazoo Co. Infrequent. C. & S.
*241. S. neglectus Nash. Small Rush-grass. Frequent. Lansing, Saugatuck, C. F. Wheeler; Detroit, O. A. Farwell.
242. S. serotinus (Torr.) A. Gray. Late-flowered Dropseed. Sandy wet places, Gray's Manual; Keweenaw Co., O. A. Farwell. Rare.
*243. S. vaginaeflorus (Torr.) Wood. Sheathed Rush-grass. Flint; Detroit, Dr. A. B. Lyons; Ann Arbor, C. A. Davis. Rare. C. & S.

CINNA L.

*244. C. arundinacea L. Low grounds. Frequent.
*245. C. latifolia (Trev.) Griseb. Slender Wood Reed-grass. *C. pendula* Trin. Wet woods throughout.

LIBRARY OF THE UNIVERSITY OF CALIFORNIA

AGROSTIS L.

*246. A. alba L. Fiorin or White Bent-grass. Red Top. *A. alba vulgaris* Thurber. Naturalized from Europe. A common pasture and meadow grass.
*247. A. alba stolonifera (L.) Vasey. Detroit, O. A. Farwell.
*248. A. canina L. Brown Bent-grass. Ann Arbor, Winch. Cat., Ronald, Ionia Co. Infrequent.
*249. A. coarctata Ehrh. Detroit, O. A. Farwell.
*250. A. hyemalis (Walt.) B. S. P. Rough Hair-grass. *A. scabra* Willd. Marshes and sterile soil. Common. Th.
*251. A. perennans (Walt.) Tuckerman. Thin-grass. Swamps and damp woods. S. Mich., Wright; Macomb Co.; Hubbardston. Infrequent.
*252. A. pseudo-intermedia O. A. Farwell. New name for *A. intermedia* Scribn. Detroit, O. A. Farwell.

CALAMAGROSTIS Adans.

*253. C. Canadensis (Michx.) Beauv. Blue-Joint. Common.
254. C. confinis (Willd.) Nutt. Bog Reed-grass. *C. Lapponica* Trin. Isle Royale, Lake Superior, T. C. Porter; Sturgis, F. P. Daniels.
255. C. Langsdorfii (Link.) Trin. Langsdorf's Reed-grass. Isle Royale, T. C. Porter.
256. C. neglecta (Ehrh.) Gaertn. Narrow Reed-grass. *C. stricta* Beauv. Flint, D. Clark; S. W., H. S. Pepoon; and northward to Lake Superior, A. Gray. Rare.

AMMOPHILA Host.

257. A. arenaria (L.) Link. Sea Sand-reed. N. & U. P. *A. arundinacea* Host. This is one of the few plants found both by the ocean and the shores of the Great Lakes, Petoskey; Point au Chene, Winch. Cat.; shores of Lake Huron, C. K. Dodge. Infrequent.

CALAMOVILFA Hack.

258. C. longifolia (Hook.) Hack. Long-leaved Reed-grass. *Calamagrostis longifolia* Hook. Sand dunes along L. Michigan. S. Haven, Bailey; Petoskey; Point au Chene, Winch. Cat.; Oscoda. Th.

APERA Adans.

*259. A. Spica-venti L. Beauv. College lawn.

DESCHAMPSIA Beauv.

*260. D. caespitosa (L.) Beauv. Tufted Hair-grass. In bogs. Frequent. Th.
261. D. flexuosa (L.) Trin. Common Hair-grass. Oscoda; Baldwin, to Lake Superior; Keweenaw Co., O. A. Farwell.

TRISETUM Pers.

262. T. subspicatum (L.) Beauv. Narrow False-oat. *T. subspicatum molle* A. Gray. Marquette, E. J. Hill; Whitney's Cat. Common. U. P.

AVENA L.

263. A. Smithii Porter. Smith's Oat. *Melica Smithii* Vasey. Keweenaw Point; Isle Royale, Dr. Robbins, in Gray's Man.; woods near Sault Ste. Marie, C. E. Smith; shore of Crystal Lake, Benzie Co., L. H. Dewey; Vanderbilt, Otsego Co.; Keweenaw Co., O. A. Farwell.
**264. A. striata Michx. Oat Grass. Woods. Infrequent.

ARRHENATHERUM Beauv.

*265. A. elatius (L.) Beauv. Tall Oat-grass. *A. avenaceum* Beauv. Bay Co., Macomb Co. Escaped from cultivation.

DANTHONIA DC.

266. D. intermedia Vasey. Keweenaw Co., O. A. Farwell.
*267. D. spicata (L.) Beauv. Wild Oat-grass. Sterile soil. Frequent. Th.

CAPRIOLA Adans. CYNODON Rich.

*268. C. Dactylon (L.) Kuntze. Bermuda-grass. Near the depot of the Chicago and Grand Trunk R. R. in Lansing.

SPARTINA Schreb.

*269. S. cynosuroides (L.) Willd. Fresh-water Cord-grass. Banks of rivers, to N. Shore L. Superior, Agassiz. Tuscola and Washtenaw. Th.

ATHEROPOGON Muhl. BOUTELOUA Lagasca. in part.

270. A. curtipendulus (Michx.) Fourn. *Bouteloua racemosa* Lag. Dry plains. S. Mich., Wright Cat.; Macomb Co., on farm of L. D. Watkins, Manchester; Sturgis, F. P. Daniels.

ELEUSINE Gaertn.

*271. E. Indica (L.) Gaertn. Dog's-tail or Wire-grass. S. & C. Ionia Co.; South Haven, Bailey; S. Mich., Wright Cat. Infrequent. S. & C.

PHRAGMITES Trin.

*272. P. Phragmites (L.) Karst. Reed. *P. communis* Trin. Swamps. Infrequent. Th.

TRICUSPIS Beauv. TRIODIA R. Br., in part.

273. T. seslerioides (Michx.) Torr. Tall Red-top. *Triodia cuprea Jacq.* S. Mich., Wright's Cat. S. E.

TRIPLASIS Beauv. TRIODIA R. Br., in part.

274. T. purpurea (Walt.) Chapm. Sand-grass. Shore of Lake Erie, Monroe Co., C. F. Wheeler; shore of Lake Huron, C. K. Dodge.

ERAGROSTIS Beauv.

275. E. capillaris (L.) Nees. Capillary Eragrostis. Coleman's Cat.; S. Mich., Wright's Cat.; Ann Arbor, M. W. Harrington. Rare.

*276. E. Eragrostis (L.) Karst. Low Eragrostis. *Eragrostis minor* Host.

*277. E. Frankii Steud. Frank's Eragrostis. Roadsides in low ground. Hubbardston; Gratiot Co.; Grand Rapids; Detroit, O. A. Farwell. Frequent. C. & S.

*278. E. hypnoides (Lam.) B. S. P. Creeping Eragrostis. *E. reptans* Nees. Gravelly borders of streams. Common.

*279. E. major Host. Stink Grass. Waste places and gardens, a common weed. Th.

280. E. pectinacea (Michx.) Steud. Purple Eragrostis. *E. pectinacea spectabilis* A. Gray. Dry sandy ground. S. Mich., Wright Cat. Reaches its northern limits at Howard City. Infrequent. C. & S.

281. E. pilosa (L.) Beauv. Tufted Eragrostis. Grand Rapids, Coleman's Cat.; Macomb Co.; Detroit, O. A. Farwell; S. W., H. S. Pepoon.

282. E. Purshii Schrad. Pursh's Eragrostis. Norway, C. F. Wheeler; Rochester, W. A. Brotherton and J. W. Stacey; Detroit, O. A. Farwell; Bay Co., G. M. Bradford. I see no way of distinguishing this species from E. pilosa (L.) Beauv. and believe them to be the same.

EATONIA Raf.

*283. E. nitida (Spreng.) Nash. Slender Eatonia. *E. Dudleyi* Vasey. Hubbardston, C. F. Wheeler; Orion, O. A. Farwell. Rare.

*284. E. obtusata (Michx.) A. Gray. Blunt-scaled Eatonia. S. E., Wright Cat.; Flint; Hubbardston, northward. Rare.

*285. E. Pennsylvanica (D C.) A. Gray. Moist woods. Common. Th.

KOELERIA Pers.

*286. K. cristata (L.) Pers. Dry hills. Frequent. Th.

KORYCARPUS Zea. DIARRHENA Raf.

*287. K. diandrus (Michx.) Kuntze. *Diarrhena American* Beauv. S. Mich., Wright Cat.; Flint; Hubbardston. Reaches its northern limits in Ionia Co. Rare.

BRIZA L.

288. B. media L. Quaking Grass. Bay City, G. M. Bradford, C. A. Davis.

CYNOSURUS L.

289. Cynosurus cristatus L. Dog's-tail Grass. Scarce.

DACTYLIS L.

*290. D. glomerata L. Cock's Foot. Orchard Grass. Escaped from cultivation and becoming frequent.

POA L.

291. P. alpina L. Alpine Spear-grass. U. P. Isle Royale, C. G. Loring, Jr., in Gray's Manual; Keweenaw Co., O. A. Farwell. U. P.

*292. P. alsodes A. Gray. Grove Meadow-grass. Banks of brooks, flowering early. Flint; Hubbardston; Inland, Grand Traverse Co. Infrequent. L. P.

*293. P. annua L. Low Spear-grass. Low grounds. Common. Th.

294. P. autumnalis Muhl. Flexuous Spear-grass. *P. flexuosa* Muhl. Swamps. Hubbardston; Constantine, C. F. Wheeler; Flint, Dr. Clark; Rochester, O. A. Farwell.

*295. P. compressa L. Wire-grass. Blue Grass (of N. Eng. and N. Y.). Dry fields. Common. Th.

*296. P. debilis Torr. Weak Spear-grass. Hillsides. S. Mich., Winch. Cat.; Macomb Co.; Hubbardston to N. Shore of Lake Superior, Macoun. Infrequent. Th.

*297. P. flava L. Fowl Meadow-grass. *P. serotina* Ehrh. Wet meadows, where it is a valuable grass, sometimes mistaken for Red-top, Agrostis aba. Th.

298. P. glauca Vahl. Glaucous Spear-grass. Near Grayling; Isle Royale; Keweenaw Co., O. A. Farwell.

300. P. nemoralis L. Wood Meadow-grass. *P. caesia strictoir* A. Gray. Throughout N. Mich. Infrequent except in the north.

*301. P. pratensis L. Kentucky Blue Grass. June Grass. Pastures and meadows.

302. P. sylvestris A. Gray. Sylvan Spear-grass. Low woods. S. Mich., Winch. Cat.; Flint; Hubbardston. Rare. C. & S.

303. P. trivialis L. Roughish Meadow-Grass. S. Mich., Wright Cat.; Flint. Sparingly in cultivation.

GRAPHEPHORUM Desv.

*304. G. melicoideum (Michx.) Beauv. River banks. Macomb Co.; Flint; Hubbardston; Petoskey; Grand Detour, Upper Michigan, Prof. Porter. Rare South, abundant at Pic River, Macoun. Th.

305. G. melicoides major A. Gray. Swamps. Hubbardston; Washington, Macomb Co.

PANICULARIA Fabr. GLYCERIA R. Br.

*306. P. Americana (Torr.) MacM. Tall Manna-grass. *Glyceria grandis* S. Wats. Common. Th.

*307. P. Canadensis (Michx.) Kuntze. Rattlesnake-grass. *Glyceria Canadensis* Trin. Borders of marshes. S. W., Winch. Cat.; Flint; Ionia Co. Infrequent. Th.

*308. P. fluitans (L.) Kuntze. Floating Manna-grass. *Glyceria fluitans* R. Br. Shallow water. Common. Th.

308a. P. laxa Scribn. Northern Manna-grass. Muskegon, W. J. B.; Keweenaw Co., O. A. Farwell.

*309. P. nervata (Willd.) Kuntze. Nerved Manna-grass. *Glyceria nervata* Trin. Very variable. Woods and wet meadows. Th.

*310. P. pallida (Torr.) Kuntze. Pale Manna-grass. *Glyceria pallida* Trin. Constantine, Wheeler; Keweenaw Co., O. A. Farwell; Manistee and Sturgis, F. P. Daniels. Wet places. Rare. S.

PUCCINELIA Parl.

311. P. airoides (Nutt.) S. Wats. Slender Meadow-grass. Bay City, G. M. Bradford. Near salt works.

FESTUCA L.

*312. F. elatior L. Taller Meadow Fescue. *F. elatior pratensis* A. Gray. Meadows and roadsides.

*313. F. nutans Willd. Nodding Fescue-grass. Woods. Frequent. L. P.

*314. F. octoflora Walt. Slender Fescue-grass. *F. tenella* Willd. Dry grounds. Infrequent. Th.

*315. F. ovina L. Sheep's Fescue. Dry grounds. Common. Very variable. Th.

*316. F. ovina capillata (Lam.) Hack. Hair-leaved Fescue. In lawns. Not uncommon.

317. F. ovina duriuscula (L.) Hack. Hard Fescue. S. Mich., Wright Cat.; Petoskey; Bay Co., G. M. Bradford; Keweenaw Co., Robbins; Isle Royale, Gillman. Sparingly in cultivation. Th.

*318. F. ovina marginata Hack. In lawns. Common.

319. F. rubra heterophylla Hack. Variable-leaved Fescue. Clifton, O. A. Farwell; Frankfort, W. J. B.; shores of Traverse Bay; Muskegon, C. F. Wheeler.

BROMUS L.

*320. B. asper Murr. Hairy Brome-grass. Gray's Manual.

321. B. breviaristatus Thurb. Short-awned Chess. First collected on low sand dunes near Bay View in August, 1879, by C. F. Wheeler. In June, 1881, Prof. V. M. Spalding also collected this species near Charlevoix. The farthest station east known for this grass.

322. B. brizaeformis Fisch & Mey. Quake-grass. Manistee, F. P. Daniels.

*323. B. ciliatus L. Fringed Brome-grass. *B. purgans* L. Frequent. Variable.

*324. B. hordeaceus L. Soft Chess. *B. mollis* L. Agricultural College, W. J. B.; Detroit, O. A. Farwell.

*325. B. inermis Leyss. Awnless Brome-grass. Sandy meadows. Spreading.

*326. B. Kalmii A. Gray. Wild Chess. Dry open woods. S. E., Wright's Cat.; Flint; Macomb Co.; Hubbardston; northward to Quinnessec, E. J. Hill. Infrequent.

327. B. racemosus L. Smooth Brome-grass. Detroit, O. A. Farwell; Alma, C. A. Davis. Th.

*328. B. secalinus L. Chess. Cheat. Too common in wheat fields.

328a. B. sterilis L. Barren Brome-grass. Bay City, G. M. Bradford.

*329. B. tectorum L. Downy Brome-grass. Grand Rapids. Spreading rapidly; Detroit, O. A. Farwell; Bay Co., G. M. Bradford.

LOLIUM L.

331. L. perenne L. Common Darnel. Ray or Rye-grass. Macomb Co., Dr. Cooley; Ionia Co.; Bay City, G. M. Bradford. Scarcely naturalized.

332. L. temulentum L. Bearded Darnel. Scarcely naturalized. Flint; Macomb Co.; S. Mich., Wright Cat.

AGROPYRON Gaertn.

*333. A. caninoides (Ramalay) Beal. Lansing, escaped from cultivation.

334. A. caninum (L.) R. & S. Awned Wheat-grass. Th. S. Mich., Winch. Cat.; Macomb Co.; Hubbardston; northward.

335. A. dasystachyum (Hook.) Vasey. Northern Wheat-grass. Common at Petoskey; N. W., Winch. Cat.; Oscoda and the sandy shores of the great lakes; Keweenaw Co., O. A. Farwell. N. & U. P.

336. A. occidentale Scribn. Occasional. Port Huron, C. K. Dodge.

337. A. pseudorepens S. and S. Shores of Grand Traverse Bay, C. F. Wheeler; St. Clair Co., C. K. Dodge; Keweenaw and Marquette counties, O. A. Farwell.

*338. A. repens (L.) Beauv. Quick-grass or Quack-grass. Cultivated grounds. Varies greatly. Sometimes grows to the exclusion of everything else, and is widespread enough to be considered a pest. Th.

339. A. Richardsoni Shreb. Bay View, W. J. B.; shore of Grand Traverse Bay, C. F. Wheeler; Keweenaw Co., O. A. Farwell.

340. A. spicatum (Pursh.) Scribn. & Small. Western Wheat-grass. Introduced into Bay City, G. M. Bradford; Keweenaw Co., Detroit, O. A. Farwell.

341. A. tenerum Vasey. Port Huron; along railroads; an advent from the west, C. K. Dodge.

342. A. violaceum (Hornem.) Vasey. Purplish Wheat-grass. Keweenaw Peninsula, Farwell; Lake Superior, Porter.

SECALE L.

*343. S. cereale L. Rye. Escaped from cultivation.

HORDEUM L.

*344. H. jubatum L. Squirrel-tail Grass. Sands. N. shore of Lake Superior, Agassiz; Sault de Ste. Marie, R. Bell; Bay City. Infrequent. Th.

345. H. nodosum L. Detroit, O. A. Farwell. Occasional.

ELYMUS L.

*346. E. Canadensis L. Nodding Wild Rye. *E. Canadensis glaucifolius* A. Gray. River banks. Common. Th.

347. E. glaucus Buckl. Smooth Wild Rye. S. W., H. S. Pepoon. Britton and Brown. *E. Sibiricus Americanus* S. Wats. & Coul. Marquette, Porter in Gray's Manual; Keweenaw Co., O. A. Farwell.

348. E. mollis Trin. Smooth Wild Rye. Shores of the Great Lakes, Gray's Manual; north shore of Lake Superior, Agassiz.

*349. E. robustus Scribn. & Sm. Muskegon, C. D. McLouth; Lansing, W. J. B.

*351. E. striatus Willd. Slender Wild Rye. *E. striatus villosus* A. Gray. S. Mich., Wright Cat.; Flint; Hubbardston; Manistee, F. P. Daniels; northward to L. Superior. Th.

*352. E. Virginicus L. Virginia Wild Rye. River banks. Common. Th.

*353. E. Virginicus glaucus Beal. Grand River Valley, with the species. W. J. B.

HYSTRIX Moench. ASPRELLA Willd.

*354. Hystrix Hystrix (L.) Millsp. Bottle-Brush grass. Th. *Asprella Hystrix* Willd. Moist woods. A variety of this grass is found in one locality near Hubbardston with smooth and very glaucous culms, leaves rough, hairy.

CYPERACEÆ J. St. Hil. Sedge Family.

CYPERUS L.

*356. C. diandrus Torr. Low Cyperus. Low grounds. Common. C. & S.

358. C. Engelmanni Steud. Englemann's Cyperus. Fruitport, E. J. Hill; shore of Park Lake, Clinton Co., C. F. Wheeler; St. Clair Co., C. K. Dodge. Rare.

*359. C. erythororhizos Muhl. Red-rooted Cyperus. Macomb Co., D. Cooley; Bay City, G. M. Bradford. Infrequent.

*360. C. esculentus L. Yellow Nutt-grass. A troublesome weed on low grounds, spreading rapidly by means of its nut-like tubers. Hard to eradicate. Muir; Flint; Grand Rapids; Detroit; north to Oscoda. Frequent. L. P.

361. C. esculentus angustispicatus Britton. Detroit, A. O. Farwell.

*362. C. filiculmis Vahl. Slender Cyperus. Sterile soil. Common. C. & S.

363. C. flavescens L. Yellow Cyperus. Grand Rapids, Coleman Cat.; S. Mich., Wright Cat. Rare. S.

364. C. Houghtonii Torr. Houghton's Cyperus. Hilltops near Indian River, Wheeler; Long Lake, Cheboygan Co., B. & K. St. Clair., C. K. Dodge.

*365. C. inflexus Muhl. Awned Cyperus. *C. aristatus* Boeckl. Grand Rapids, Coleman; Ionia; banks of Cedar River, Agricultural College. Not common.

366. C. refractus Engelm. Detroit, O. A. Farwell.

*367. C. rivularis Kunth. Shining Cyperus. *C. diandrus castaneus* Torr. Frequent.

368. C. Schweinitzii Torr. Schweinitz's Cyperus. Lake Michigan shore at South Haven, L. H. Bailey; Kalamazoo, Tuthill; Manistee, F. P. Daniels, S. West, H. S. Pepoon.

369. C. speciosus Vahl. Michaux's Cyperus. Low grounds. Hubbardston; Flint.
*370. C. strigosus L. Straw-colored Cyperus. Low grounds. Common. C. & S.
371. C. strigosus capitatus Boeckl. Detroit, O. A. Farwell.
372. C. strigosus robustior Kunth. Detroit, O. A. Farwell.

KYLLINGA Rott.

374. K. pumila Michx. Low Kyllingia. Dr. Lyons. S.

DULICHIUM L. C. Richard.

*375. D. arundinaceum (L.) Britton. *D. spathaceum* Pers. Borders of swamps and ponds. Common. Th.

ELEOCHARIS R. Br.

*376. E. acicularis (L.) R. & S. Spike-rush. Alma, Washtenaw Co., C. A. Davis. Th.
377. E. acuminata (Muhl.) Nees. Flat-stemmed Spike-rush. Shores of lakes.
378. E. Engelmanni Steud. Engelmann's Spike-rush. Jackson, S. H. Camp; Detroit, O. A. Farwell.
*379. E. intermedia (Muhl.) Schultes. Matted Spike-rush. Macomb Co., Dr. D. Cooley; Grand Traverse Bay, Winch. Cat.; Hubbardston, C. F. Wheeler; Alma, C. A. Davis. Low river banks.
380. E. interstincta (Vahl.) R. & S. Knotted Spike-rush. *E. equisetoides* Torr. Wright Cat., Jackson Co., 1838.
*381. E. mutata (L.) R. & S. Quadrangular Spike-rush. *E. quadrangulata* R. & S. S. Mich., A. Gray; Flint; borders of Crystal Lake, Montcalm Co.; shores of Park Lake and Pine Lake; Algonac, W. S. Cooper. Infrequent. C. & S.
381a. E. obtusa Schults. Ovoid Spike-rush. Wet grounds. N. shore of L. Superior, Agassiz; southward; St. Clair Co., C. K. Dodge; Alma, Ann Arbor, C. A. Davis. Th.
*382. E. olivacea Torr. Bright-green Spike-rush. Shores of Park Lake, Clinton Co., Wheeler; Vestaburg, C. A. Davis.
*383. E. ovata (Roth.) R. & S. Keweenaw Co., O. A. Farwell.
*384. E. palustris (L.) R. & S. Creeking Spike-rush. Wet places. Alma, Ann Arbor, C. A. Davis; Bay Co., G. M. Bradford.
385. E. palustris calva A. Gray. Lake Antoine, E. J. Hill.
*386. E. palustris glaucescens (Willd.) A. Gray. Alma, C. A. Davis. Frequent.
*387. E. Robbinsii Oakes. Robbins' Spike-rush. Shallow water. Park Lake, Clinton Co. The only station known in the State, C. F. Wheeler.
*388. E. palustris vigens L. H. Bailey. Indian River, Cheboygan Co.; along the Great Lakes, Gray's Man. Keweenaw Co., O. A. Farwell.
389. E. rostellata Torr. Beaked Spike-rush. Marshes. Hubbardston; Macomb Co., Drummond's I., Winch. Cat. Tuscola and Washtenaw Counties, C. A. Davis; Bay Co., G. M. Bradford. Rare.
390. E. tenuis (Willd.) Schultes. Slender Spike-rush. Oscoda; east coast of L. Superior, Canadian Cat.; Keweenaw Co., O. A. Farwell; Ann Arbor, C. A. Davis. Th.

PSILOCARYA Torr.

391. P. scirpoides Torr. Long-beaked Bald-rush. Shore of Crooked Lake, near Grand Rapids, Miss E. J. Colé.

STENOPHYLLUS Raf.

392. S. capillaris (L.) Britton. Hair-like Stenophyllus. *Fimbristylis capillaris* A. Gray. S. Mich., Wright; Detroit; St. Clair Co., C. K. Dodge; Algonac, W. S. Cooper. Rare.

FIMBRISTYLIS Vahl.

*393. F. autumnalis (L.) R. & S. Slender Fimbristylis. S. Mich., Wright; Pine Lake, Ingham Co., Manistee, F. P. Daniels; Vestaburg, C. A. Davis.
394. F. castanea (Michx.) Vahl. Marsh Fimbristylis. Port Huron, C. K. Dodge.

SCIRPUS L.

*395. S. Americanus Pers. Three-square. *S. pungens* Vahl. Borders of ponds, Alma, Ann Arbor, C. A. Davis. Common. Th.

*396. S. atrovirens Muhl. Dark green Bulrush. Alma, Ann Arbor, C. A. Davis. Wet meadows. Common.

397. S. atrovirens pallidus Britton. Introduced from the west into Bay Co., G. M. Bradford; Keweenaw Co., Detroit, O. A. Farwell.

398. S. caespitosus L. Tufted Club-rush. Dr. A. B. Lyons; Marquette, E. J. Hill; north shore of Lake Superior, Agassiz. U. P.

*399. S. Clintonii A. Gray. Clinton's Club-rush. Bluffs along Fish Creek, Hubbardston; Bath, L. H. Bailey; St. Clair Co., C. K. Dodge; Bay City, C. A. Davis. Rare. C.

*400. S. cyperinus (L.) Kunth. Wool-grass. Wats. & Coult. Alma, Ann Arbor, C. A. Davis. Frequent and variable.

401. S. cyperinus Eriophorum (Michx.) Britton. *Eriphorum cyperinum laxum.* Keweenaw Co., O. A. Farwell; Bay City, G. M. Bradford.

*402. S. debilis Pursh. Weak-stalked Club-rush. Macomb Co., Cooley; Park Lake, Clinton Co., Wheeler; St. Clair Co., C. K. Dodge; Manistee, F. P. Daniels; Alma, C. A. Davis. Rare.

403. S. Hallii A. Gray. Muskegon, C. D. McLouth.

404. S. fluviatilis (Torr.) A. Gray. River Club-rush. Margins of rivers, S. Mich., Winch. Cat.; Macomb Co.; Hubbardston; Detroit; abundant along Maple River; west of Lake Superior, Macoun. Th.

*405. S. lacustris L. Great Bulrush. In still water. Common. Th.

*406. S. lineatus Michx. Reddish Bulrush. Common in Bay Co., G. M. Bradford; S. W., H. S. Pepoon. River banks, S. Mich., Winch Cat.; Flint; Macomb Co.; Hubbardston; St. Clair Co., C. K. Dodge; Keweenaw Co.; Alma, Ann Arbor, C. A. Davis.

407. S. microcarpus Presl. Small-fruited Bulrush. *S. sylvaticus digynus* Boeckl. Lake Nipigon, Ont., Macoun; Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels.

408. S. nanus Spreng. Dwarf Club-rush. Hubbardston; Clinton Co., E. F. Smith. *Eleocharis pygmaea* Torr.

409. S. Olneyi A. Gray. Olney's Bulrush. Border of deer lick near Hubbardston, Wheeler.

410. S. pauciflorus Lightf. Few-flowered Club-rush. Grand Rapids, Miss E. J. Cole; Port Austin, C. A. Davis. Frequent northward.

411. S. polyphyllus Vahl. Leafy Bulrush. S. Mich., Wright Cat.

*412. S. Smithii A. Gray. Smith's Club-rush. Shore of Park Lake, C. F. Wheeler, 1891; Bay Co., G. M. Bradford; Manistee, F. P. Daniels.

*413. S. subterminalis Torr. Water Club-rush. Houghton Lake; Woodward Lake, Ionia Co.; Flint; Macomb Co.; S. Mich., Wright Cat.; north of Lake Superior. Can. Cat.; Alma, C. A. Davis. Infrequent.

413a. S. sylvaticus L. Wood Bulrush. Keweenaw Co., O. A. Farwell.

*414. S. Torreyi Olney. Torrey's Bulrush. Border of Pine Lake, Ingham Co., Bailey.

ERIOPHORUM L.

415. E. alpinum L. Alpine Cotton-grass. Mud Lake; Petoskey, E. J. Hill; Macomb Co., Cooley; Elk Rapids, W. S. Cooper; Keweenaw Co. Infrequent. Th.

416. E. gracile Koch. Slender Cotton-grass. S. Mich. Wright Cat.; Flint; Hubbardston; Montcalm Co., and northward; Kalamazoo, Tuthill; Ann Arbor, C. A. Davis. Rare. Th.

*418. E. polystachyon L. Tall Cotton-grass. Swamps. Common. Th.

*419. E. vaginatum L. Sheathed Cotton-grass. Sphagnous swamps. S. Mich., Winch. Cat.; Macomb Co.; St. Clair Co., C. K. Dodge; near Ann Arbor, C. A. Davis; Hubbardston; Bay Co., and northward. Th.

*420. E. Virginicum L. Virginia Cotton-grass. Sphagnous swamps. Ann Arbor, Allmend. Cat.; Flint; Bay Co.; Hubbardston, Vestaburg, C. A. Davis, northward. Infrequent. Th.

421. E. Virginicum album A. Gray. Keweenaw Co., O. A. Farwell.

FUIRENA Rottb.

422. F. squarrosa Michx. Squarrose Fuirena. *F. squarrosa pumila* Torr. Macomb Co., Dr. D. Cooley. Rare. C.

HEMICARPHA Nees and Arn.

*423. H. micrantha (Vahl.) Britton. C. & S. *H. subsquarrosa* Nees. S. W., Wright Cat.; Pine Lake, Ingham Co. Rare.

RHYNCHOSPORA Vahl.

424. R. alba (L.) Vahl. White Beaked-rush. Bogs. Ann Arbor, Allmend. Cat.; Macomb Co.; Hubbardston; N. E., Winch. Cat.; Keweenaw Co., F.; Mackinaw City, Wheeler; Alma, Ann Arbor, C. A. Davis. Th.

425. R. capillacea Torr. Capillary Beaked-rush. Bogs and sandy lake shores. Hubbardston; Flint; Macomb Co.; Petoskey; St. Clair Co.; Jackson Co., C. A. Davis. Infrequent. Th.

426. R. capillacea laeviseta Hill. C. K. Dodge. Shore Grand Traverse Bay, near Torch Lake, E. J. Hill; Saginaw and Tuscola Counties, C. A. Davis; Orion, O. A. Farwell.

427. R. cymosa Ell. Sturgis, F. P. Daniels.

428. R. fusca R. & S. Escanaba, E. J. Hill.

*429. R. glomerata (L.) Vahl. Clustered Beaked-rush. S. Mich., Wright Cat. Macomb Co.; Hubbardston; Pine Lake, Ingham Co.; St. Clair Co., C. K. Dodge; Keweenaw Co., O. A. Farwell; Vestaburg, C. A. Davis.

CLADIUM P. Browne.

*430. C. mariscoides Torr. Twig-rush. Bogs. S. Mich., Winch. Cat.; Macomb Co.; Flint; Hubbardston; Cheboygan Co., B. & K.; Tuscola and Washtenaw Counties, C. A. Davis. L. P.

SCLERIA Berg.

431. S. triglomerata Michx. Tall Nut-rush. S. Mich., Wright Cat., Macomb Co.; Flint; St. Clair Co., C. K. Dodge; Livingston Co. Rare. S.

432. S. verticillata Muhl. Low Nut-rush. S. Macomb Co., D. Cooley; St. Clair Co., C. K. Dodge; Orion, O. A. Farwell; Sebewaing, C. A. Davis. Rare.

CAREX L.

433. C. abacta Bailey. Yellowish Sedge. *C.Michauxiana* Boeckl. Keweenaw Co., O. A. Farwell; near Portage River, T. C. Porter.

434. C. adusta Boott. Browned Sedge. Crawford Co., Bailey; Grayling, G. H. Hicks. Rare.

*435. C. alata Torr. Broad-winged Sedge. *C. straminea alata* Bailey. Keweenaw Co., O. A. Farwell; Crystal Lake, Montcalm Co., Wheeler; South Haven; Niles, Dr. G. L. Ames, in University Herb.

436. C. albolutescens cumulata Bailey. Bay Co., G. M. Bradford.

*437. C. Albursina Sheldon. White Bear Sedge. *C. laxiflora latifolia* Boott. Fox-tail Sedge. Macomb Co.; Gray's Man.; Agricultural College grounds; Ypsilanti, O. A. Farwell. Frequent. S.

*438. C. alopecoidea Tuckerman. Fox-tail Sedge. Macomb Co.; Gray's Man.; Agricultural College grounds; Ypsilanti, O. A. Farwell.

439. C. alopecoidea sparsispicata Dewey. Flint, Clark; Macomb Co., Cooley; Detroit, O. A. Farwell.

440. C. alpina Swartz. Alpine Sedge. Isle Royale, Whitney Cat. U. P.

441. C. altocaulis (Dewey) Britton. Sheathed Sedge. *C. Saltuensis* Bailey. Keweenaw Co., O. A. Farwell; Oscoda Co., L. H. Bailey. N. & U. P.

*442. C. aquatilis Wahl. Water Sedge. Margins of streams. S. Mich., Wright Cat.; Flint; Macomb Co., Hubbardston; Sitting Rabbit, Winch. Cat.; Alma, Ann Arbor, C. A. Davis. Infrequent. Th.

443. C. aquatilis elatior Bab. Bay Co., G. M. Bradford. Belle Isle, O. A. Farwell.

444. C. arcta Boott. Northern Clustered Sedge. *C. canescens polystachya* Boott. Keweenaw Co., O. A. Farwell.

445. C. arctata Boott. Drooping Wood Sedge. Woods. S. Mich., Winch. Cat.; Petoskey; Hubbardston; Grand Ledge; Van Buren Co., Bailey. Th.

446. C. arctata Faxoni Bailey. Isle Royale, Dr. Sandberg; Keweenaw Co., O. A. Farwell.

447. C. arctata x castanea Bailey. Keweenaw Co., O. A. Farwell.

*448. C. aristata R. Br. Awned Sedge. *C. trichocarpa aristata* Bailey. Agricultural College grounds; northward to Lake Superior.

*449. C. Asa-Grayi Bailey. Gray's Sedge. *C. Grayi* Carey. Low grounds. Macomb Co.; Flint; Hubbardston; Lenawee Co., Detroit, O. A. Farwell. Rare.

*450. C. aurea Nutt. Golden-fruited Sedge. Borders of cool springs. S. E. Wright Cat.; Ann Arbor, Allmend. Cat.; Macomb Co.; Pine Lake, Ingham Co.; Hubbardston; Petoskey and northward; St. Clair Co., C. K. Dodge. Infrequent. Th.

451. C. Bicknellii Britton. Bicknell's Sedge. *C. straminea Crawei* Boott. Ann Arbor, Dr. D. Clark; Grass Lake, C. F. Wheeler.

*452. C. bromoides Schk. Brome-like Sedge. Low grounds. Common. Th.

453. C. brunnescens (Pers.) Poir. Brownish Sedge. *C. canescens alpicola* Wahl. Keweenaw Co., O. A. Farwell.

*454. C. brunnescens gracilior Britton. *C. canescens vulgaris* Bailey. Common.

*455. C. canescens L. Silvery Sedge. In swamps. Th.

456. C. capillaris L. Hair-like Sedge. Point de Tour, Lake Michigan, A. Gray; Sturgeon Point, Alcona Co.; Mackinaw, G. H. Hicks. Common in the Lake Superior region. N. & U. P.

*457. C. Careyana Torr. Cary's Sedge. Rich woods. Macomb Co.; Flint; Hubbardston; southward. Alma, C. A. Davis. Rare. S.

458. C Caroliniana Schwein. Carolina Sedge. Lapeer, Miss M. Owen.

459. C. castanea Wahl. Chestnut Sedge. Keweenaw Co., O. A. Farwell. Th.

*460. C. cephaloidea Dewey. Thin-leaved Sedge. Fields. Macomb Co., Cooley; Ann Arbor, Allmend. Cat.; St. Clair Co., C. K. Dodge; Alma, C. A. Davis; Detroit, O. A. Farwell. Rare. C. & S.

*461. C. cephalophora Muhl. Oval-headed Sedge. Open woods. Common. C. & S.

*462. C. chordorhiza L. f. Creeping Sedge. Th. Upland swamps. Macomb Co.; Hubbardston. Rare. Th.

*463. C. comosa Boott. Bay Co., G. M. Bradford; C. & S.

464. C. conjuncta Boott. Manistee, F. P. Daniels.

465. C conoidea Schk. Field Sedge. S. E. Wright Cat.; Flint; Manistee, F. P. Daniels; Ann Arbor, C. A. Davis. Infrequent. C. & S.

466. C. costellata Britton. Ribbed Sedge. Th. *C. virescens costata* Dew. S. Mich., Winch. Cat.; Detroit; Constantine, Dundee, Wheeler; So. Haven; Keweenaw Co., O. A. Farwell.

467. C. Crawei Dewey. Crawe's Sedge. Macomb Co., Cooley; Keweenaw Co., O. A. Farwell; St. Clair Co., C. K. Dodge; Tuscola Co., C. A. Davis; Bay Co., G. M. Bradford. Rare.

468. C Crawfordii Fernald. Keweenaw Co., O. A. Farwell.

*469. C. crinita Lam. Low grounds. Common. Th.

*470. C. cristatella Britton. Crested Sedge. Th. *C. tribuloides cristata* Bailey. Hubbardston, Wheeler. Rare.

*471. C. Crus-Corvi Shuttlew. Raven's-foot Sedge. Valley of River Raisin near Dundee; Reform school marsh, Lansing, C. F. Wheeler. The only localities known in the State.

*472. C. Davisii Schwein & Torr. Davis' Sedge. Keweenaw Co., Farwell. Rare.

*473. C. decomposita Muhl. Large-panicled Sedge. Hubbardston; Ann Arbor, Allmendinger Cat.; S. Mich., Wright Cat.; Lansing, Bailey. Very rare.

474. C. deflexa Hornem. Northern Sedge. Keweenaw Co., O. A. Farwell.

475. C. deflexa Deanei Bailey. Keweenaw Co., O. A. Farwell; Mackinaw, G. H. Hicks.

476. C. deflexa Farwellii Britton. *C. deflexa meadia* Bailey. Keweenaw Co., O. A. Farwell.

*477. C. Deweyana Schwein. Dewey's Sedge. Woods. Frequent. Th.

*478. C. digitalis Willd. Slender Wood Sedge. Macomb Co.; Flint; Hubbardston. Frequent. S.

*479. C. digitalis copulata Bailey. Hubbardston; southward.

480. C. durifolia Bailey. *C. Backii* Boott. Orion, Keweenaw Co., O. A. Farwell; Ontonagon River, Whitney Cat.; West Harrisville, Alcona Co. Rare.

482. C. exilis Dewey. Coast Sedge. Keweenaw Peninsula, O. A. Farwell. The only stations known in the State for this rare species.

*483. C. festucacea Willd. Fescue Sedge. *C. straminea brevior* Dewey. Common. Th.

*484. C. filiformis L. Slender Sedge. Bogs. Alma, Ann Arbor, C. A. Davis; S.

Mich., Wright Cat.; Hubbardston; Macomb Co.; Flint; common at Pine Lake, Ingham Co.; Keweenaw Co., O. A. Farwell; Towar's Swamp, Lansing.

486. C. flacca Shreb. *C. glauca* Scop. South shore of Belle Isle, O. A. Farwell.

487. C. flava L. Yellow Sedge. Sphagnous swamps. Frequent. Th.

488. C. flava graminis Bailey. Crawford Co., Bailey; Keweenaw Co., O. A. Farwell. Rare.

489. C. foenea Willd. Hay Sedge. Alger's Camp, Alcona Co.; Potts; Oscoda Co.; Crawford Co.; Isle Royale, Dr. J. H. Sandberg.

490. C. foenea perplexa Bailey. Port Huron, C. K. Dodge; Muskegon, Mrs. H. W. Fallars. Northward.

491. C. folliculata L. Long Sedge. S. Mich., Wright Cat.; So. Haven; Flint; Macomb Co., to L. Superior. Rare. Th.

*492. C. formosa Dewey. Handsome Sedge. Macomb Co.; St. Clair Co., C. K. Dodge; Bay Co., G. M. Bradford. Rare. Th.

*493. C. fusca All. Brown Sedge. Bogs. S. E., Winch. Cat.; Macomb Co.; Hubbardston; northward to L. Superior, Whitney Cat.; Alma, C. A. Davis. Irefrequent. Th.

*494. C. gracillima Schwein. Graceful Sedge. Wet meadows. Th.

*495. C. granularis Muhl. Meadow Sedge. Low grounds. Common. Th.

496. C. granularis Shriveri Britton. *C. Haleana* Olney. Frequent in low, clay ground, C. F. Wheeler.

*497. C. grisea Wahl. Gray Sedge. Moist woods. Variable. Th.

*498. C. gynandra Schwein. Nodding Sedge. *C. crinita gynandra* Schwein. Low grounds. Common. Th.

*499. C. Hartii Dewey. Hart-Wright's Sedge. *C. retrorsa Hartii* A. Gray. Hubbardston; Agricultural College Farm; Keweenaw Co., O. A. Farwell. Th.

500. C. Haydeni Dewey. Hayden's Sedge. *C. stricta decora* Bailey. Peche Isle and Belle Isle, C. F. Wheeler.

*501. C. Hitchcockiana Dewey. Hitchcock's Sedge. Woods. Flint; Hubbardston and southward. S.

502. C. Houghtonii Torr. Houghton's Sedge. North part of Clare Co.; Keweenaw Co.; Cheboygan Co.; Oscoda northward. N. & U. P.

*503. C. hystricina Muhl. Porcupine Sedge. Wet meadows. Common. Th.

504. C. hystricina Dudleyi Bailey. Owosso, G. H. Hicks; Manistee, F. P. Daniels; Bay Co., G. M. Bradford.

*505. C. interior Bailey. Keweenaw Co., Orion, Detroit, O. A. Farwell. Common. Th.

*506. C. intumescens Rudge. Bladder Sedge. Swamps. South Haven and northward. Common. Th.

507. C. intumescens Fernaldi Bailey. Detroit, O. A. Farwell.

*508. C. Jamesii Schwein. James' Sedge. Keweenaw Co., O. A. Farwell; Cassopolis; Dundee. Infrequent. Th.

*509. C. lanuginosa Michx. Woolly Sedge. *C. filiformis latifolia* Boeckl. Keweenaw Co. and Detroit, O. A. Farwell; Bay Co., G. M. Bradford; Flint; Macomb Co.; Hubbardston; Alma; Ann Arbor, C. A. Davis.

510. C. laxiculmis Schwein. Spreading Sedge. Washington, Macomb Co., Dr. Cooley; St. Clair Co., C. K. Dodge. Rare.

*511. C. laxflora Lam. Loose-flowered Sedge. Beech and maple woods. Exceedingly variable. Common. Th.

*512. C. laxiflora blanda (Dewey) Boott. *C. laxiflora striatula* Carey. Ypsilanti, Detroit, O. A. Farwell; Bay Co., G. M. Bradford.

513. C. laxiflora patulifolia (Dewey) Carey. Frequent in center and south.

*515. C. laxiflora varians Bailey. Keweenaw Co., O. A. Farwell; southward. Common. Th.

516. C. lenticularis Michx. Lenticular Sedge. "Upper Michigan," Gray; Laughing Fish R., L. Superior, Henry Gillman. U. P.

*517. C. leptalea Wahl. Bristle-stalked Sedge. *C. polytrichoides* Willd. Low grounds. Common. Th.

*518. C. limosa L. Mud Sedge. Bogs. Ann Arbor, Allmend. Cat.; Macomb Co.; Hubbardston and northward. Irefrequent. Th.

519. C. livida (Wahl) Willd. U. P. Lake Superior, Gray's Manual; St. Clair Co., C. K. Dodge. Rare.

*520. C. longirostris Torr. Long-beaked Sedge. Woods. Flint; Hubbardston and northward; St. Clair Co., C. K. Dodge. Rare.

*521. C. lupulina Muhl. Hop Sedge. Low grounds. Common. Th.

522. C. lupulina x lupuliformis C. A. Davis. New hybrid. Alma, C. A. Davis.

*523. C. **lupulina pedunculata** Dew. "With the species, but not common," L. H. Bailey in Gray's Manual, 6th ed. Frequent at Lansing.

*524. C. **luplina x retrorsa** Dudley. Lansing, Bailey; Alma, C. A. Davis.

*525. C **lupuliformis** Sartwell. Hop-like Sedge. Washington, Dr. D. Cooley; Lansing, C. F. Wheeler; Alma, C. A. Davis.

526. C. **lurida** Wahl. Sallow Sedge. Antrim Co.; S. Mich., Wright Cat.; So. Haven; Hubbardston; Keweenaw Co., O. A. Farwell. Th.

*527. C. **lurida flaccida** Bailey. Detroit, O. A. Farwell.

*528. C. **lurida subglobosa** Fernald. Near the Agricultural College, C. F. Wheeler.

*529. C. **Magellanica** Lam. Magellan Sedge. Sphagnous swamps. Local. Th.

*530. C. **Meadii** Dewey. Mead's Sedge. *C. tetanica Meadii* Bailey. Sphagnous swamps. Macomb Co.; Hubbardston; Constantine. Infrequent.

*532. C. **monile** Tuckerman. Necklace Sedge. Th. Mud Lake, Petoskey, E. J. Hill; Keweenaw Co., O. A. Farwell; Colon, Wheeler; Reform School marsh, Lansing; Alma, Ann Arbor, C. A. Davis. Th.

533. C. **monile x utriculata,** O. A. Farwell. Sphagnum swamps. Keweenaw Co., O. A. Farwell.

*534. C. **Muhlenbergii** Schk. Muhlenberg's Sedge. Fields. Hubbardston; Bay City; Macomb Co.; southward. Infrequent. C. & S.

*535. C. **Muskingumensis** Schwein. Muskingum Sedge. Hubbardston; Flint; near Owosso, G. H. Hicks; St. Clair Co., C. K. Dodge; Ann Arbor and Alma, C. A. Davis; Ypsilanti, Detroit, O. A. Farwell; Manistee, F. P. Daniels. C. & S.

536. C. **oligocarpa** Schk. Few-fruited Sedge. Flint; Macomb Co., Dr. D. Cooley; Cassopolis, Wheeler; Detroit, Dr. Lyons.

537. C. **oligosperma** Michx. Few-seeded Sedge. Borders of swamps and lakes. Hubbardston; Woodward Lake; Houghton Lake; Crawford Co.; Keweenaw Co., O. A. Farwell; Ann Arbor, C. A. Davis.

538. C. **pallescens** L. Pale Sedge. Dr. A. B. Lyons; Keweenaw Co., O. A. Farwell; Sault de Ste. Marie, Macoun.

*539. C. **pauciflora** Lightfoot. Few-flowered Sedge. U. P., Dr. A. B. Lyons; Chocolate R., L. Superior, Henry Gillman; Towar's swamp near Agricultural College; Alma, Ann Arbor, C. A. Davis.

*540. C. **pedicellata** (Dewey) Britton. Fibrous-rooted Sedge. Th. *C. communis* Bailey.

*541. C. **pedicellata Wheeleri** (Bailey) Britton. *C. communis Wheeleri* L. H. Bailey. Hubbardston; Grand Ledge; Alcona Co.; Mackinac, G. H. Hicks; Keweenaw Co., O. A. Farwell. Infrequent.

*542. C. **pedunculata** Muhl. Long-stalked Sedge. Hillsides. Alma, C. A. Davis; Bay Co., G. M. Bradford. Infrequent. Th.

*543. C. **Pennsylvanica** Lam. Pennsylvania Sedge. Dry woods. Our commonest Sedge. Th.

*544. C. **plantaginea** Lam. Plantain-leaved Sedge. Hillsides. Alma, C. A. Davis. Scarce. Th.

*545. C. **platyphylla** Carey. Broad-leaved Sedge. Grand Rapids, Coleman's Cat.; Hubbardston; St. Clair Co., C. K. Dodge. Infrequent. S.

546. C. **prasina** Wahl. Drooping Sedge. Wet meadows. Hubbardston; S. Mich., Winch. Cat.; Flint; Port Huron, C. K. Dodge; Alma, C. A. Davis. Rare. S.

547. C **praticola** Rydb. *C. pratensis* Drej. not Hose. Northern Meadow Sedge. Pic River, Lake Superior, C. Loring.

548. C. **Pseudo-Cyperus** L. Cyperus-like Sedge. Margins of streams. S. Mich., Wright Cat.; Macomb Co.; Hubbardston; Sturgis, F. N. Daniels; Alma, Ann Arbor, C. A. Davis; northward. Infrequent. Th.

*549. C. **pubescens** Muhl. Pubescent Sedge. S. Mich., Wright Cat.; Macomb Co.; Hubbardston to Lake Superior, Whitney Cat. Frequent. Th.

550. C. **Redowskyana** C. A. Myer. Redowsky's Sedge. *C. gynocrates* Wormskiold. N. E. and N. W., Winch. Cat.; Sturgeon Point, Alcona Co.; Mio, Oscoda Co.; Manistee, F. P. Daniels.

551. C. **retroflexa** Muhl. Reflexed Sedge. *C. rosea retroflexa* Torr. Frequent. S.

*552. C. **retrorsa** Schwein. Retrorse Sedge. River banks. Frequent. Th.

553. C. **Richardsonii** R. Br. Richardson's Sedge. Barrens. Ionia Co.; Macomb Co.; Gaylord, G. E. Hancorne; Orion, O. A. Farwell. Scarce.

555. C. **riparia** W. Curtis. River-bank Sedge. Wet places. Common. Th.

*556. C. **rosea** Schk. Stellate Sedge. Moist woods. Frequent. S.

*557. C. **rosea radiata** Dewey. Dry woods. Frequent. Th.

*558. C. Sartwellii Dewey. S. Mich., Winch. Cat.; Macomb Co.; Flint; Hubbardston; Orion, O. A. Farwell; and northward. Th.

*559. C. scabrata Schwein. Rough Sedge. Banks of streams. Flint; Grand Rapids, Coleman Cat.; Hubbardston to Lake Superior, Whitney Cat. Infrequent.

560. C. Schweinitzii Dew. Schweinitz's Sedge. Near Fraser's, Crawford Co., Prof. L. H. Bailey. Only station known in State.

561. C. scirpoidea Michx. Scirpus-like Sedge. Winch. Cat.; Drummond's Isle; Keweenaw Co., O. A. Farwell. N. E.

*562. C. scoparia Schk. Pointed Broom Sedge. Th. *C. scoparia minor* Boott. Keweenaw Co., O. A. Farwell; Bay Co., G. M. Bradford. Low grounds. Common.

563. C. setifolia (Dewey) Britton. Brittle-leaved Sedgw. Th. *C. Eburnea* Boott. Hillside. South Haven on sand dunes and northward. Infrequent.

*564. C. siccata Dewey. Dry-spiked Sedge. Barrens. Macomb Co.; Flint; Hubbardston; near Park Lake, Clinton Co.; northward to Lake Superior. Infrequent.

*565. C. sparganioides Muhl. Burr-reed Sedge. Low, rich grounds, St. Clair Co., C. K. Dodge. Infrequent. C. & S.

*566. C. squarrosa L. Squarrose Sedge. S. Mich., Wright Cat.; Hubbardston; Lansing, Bailey; Keweenaw Co., O. A. Farwell; Ann Arbor, C. A. Davis. Rare.

567. C. sterilis Willd. Little Prickly Sedge. *C. echinata microstachys* Boeckl. S. Mich., Winch. Cat.; Hubbardston; Flint; Macomb Co.; Orion and Keweenaw Co., O. A. Farwell; and northward. Common.

568. C. sterilis cephalantha Bailey. *C. echinata cephalantha* Bailey. Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels.

*569. C. stipata Muhl. Awl-fruited Sedge. Low grounds. Common. Th.

*570. C. straminea Willd. Straw Sedge. Ann Arbor, Allmendinger Cat.; Macomb Co.; Flint; Hubbardston; South Haven; northward to Keweenaw Co., O. A. Farwell. Frequent. Th.

571. C. straminea ferruginea (A. Gray.) Bailey. Detroit, O. A. Farwell; Port Huron, Algonac, C. K. Dodge; Bay Co., G. M. Bradford.

*572. C. straminea mirabilis (Dewey) Tuck. *C. mirabilis* Dewey. Alma, Ann Arbor, C. A. Davis; Hubbardston and northward. Infrequent. Th.

*573. C. stricta Lam. Tussock Sedge. Low grounds. Common. Th.

574. C. stricta angustata (Boot) Bailey. Less common than the species. Ann Arbor, C. A. Davis; Hubbardston, Wheeler; Keweenaw Co., O. A. Farwell.

575. C stricta x filiformis Bailey. Keweenaw Co., O. A. Farwell.

576. C. sychnocephala Carey. Dense Long-beaked Sedge. Crystal, Montcalm Co. Only station known in the State, C. F. Wheeler. S.

*577. C. tenella Schk. Soft-leaved Sedge. Sphagnous swamps. Common.

*578. C. tenuiflora Wahl. Sparse-flowered Sedge. U. P., Whitney Cat.; Oscoda Co.; Towar's swamp, near Lansing, Wheeler, 1890; Keweenaw Co., O. A. Farwell. Th.

579. C. tenuis Rudge. Slender-stalked Sedge. *C. debilis Rudgei* Bailey. Bear Lake, Van Buren Co., E. J. Hill; shores of Barron Lake, Cass Co.; Gaylord; Alcona Co.

580. C. tenuis interjecta (Bailey) Britton. *C. debilis interjecta* Bailey. Grayling, W. J. B.

*581. C. teretiuscula Good. Lesser-panicled Sedge. Swamps. Common. Th.

*582. C teretiuscula prairea (Dewey) Britton. *C. teretiuscula ramosa* Boott. Orion, O. A. Farwell. Frequent.

*583. C. tetanica Schkuhr. Woods Sedge. S. Mich., Winch. Cat.; Grand Ledge, C. F. Wheeler; near Pine Lake, Ingham Co., L. H. Bailey; Macomb Co.; Flint, and northward.

*584. C. tetanica Woodii Bailey. Central part of the State. Infrequent.

*585. C. tribuloides Wahl. Blunt-broom Sedge. Th.

*586. C. tribuloides Bebbii Bailey. Bay Co., G. M. Bradford; Detroit, O. A. Farwell. Th.

587. C. tribuloides moniliformis (Tuck.) Britton. *C. tribuloides reducta* Bailey. Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels; Alma, C. A. Davis; Bay Co., G. M. Bradford.

588. C. tribuloides turbata Bailey. Swales. Keweenaw, O. A. Farwell. Frequent.

589. C. triceps hirsuta Bailey. In an oak wood on the farm of Hon. N. B. Hayes, in North Plains township, Ionia Co. Not known to occur elsewhere in the State.

*590. C. trichocarpa Muhl. Hairy-fruited Sedge. S. Mich., Wright Cat.; Macomb Co. Infrequent. C. & S.

*591. C. trisperma Dewey. Keweenaw Co., O. A. Farwell; Bay Co., G. M. Bradford.

*592. C. Tuckermani Dewey. Tuckerman's Sedge. Swamps. S. Mich., Wright Cat.;

Hubbardston; Flint; Macomb Co., and northward; Alma, Ann Arbor, C. A. Davis. Frequent.

*593. **C. umbellata** Schk. Umbel-like Sedge. Grand Ledge, C. F. Wheeler; Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels. Rare.

594. **C. umbellata vicina** Dewey. Alcona Co.; Keweenaw Co., O. A. Farwell; Grand Ledge, C. F. Wheeler; St. Clair Co., C. K. Dodge.

*595. **C. utriculata** Boott. Bottle Sedge. Swamps. Common. Th.

576. **C. varia** Muhl. Emmons' Sedge. Barron Lake; New Buffalo; Monroe Co.; dunes of South Haven, L. H. Bailey; Oscoda Co.; Crawford Co.; Grand Traverse, Winch. Cat.; Keweenaw Co., O. A. Farwell.

597. **C. virescens** Muhl. Dewey. Green Sedge. Detroit, O. A. Farwell; Wayne, C. A. Davis.

598. **C. viridula** Michx. Green Sedge. *C. flava viridula* Bailey. Margins of lakes. Frequent. Th.

*599. **C. vulpinoidea** Michx. Fox Sedge. Low meadows. Common. Th.

600. **C. Willdenovii** Schk. Willdenow's Sedge. Bear Lake, Van Buren Co., E. J. Hill. Rare.

ARACEÆ Neck. Arum Family.

ARISAEMA Mart.

*601. **A. Dracontium** (L.) Schott. Green Dragon. Dragon-root. C. & S.

*602. **A. triphyllum** (L.) Torr. Indian Turnip. Rich woods. Th.

PELTANDRA Raf.

*603. **P. Virginica** (L.) Kunth. Green Arrow-arum. *P. undulata* Raf. S. Mich., Wright Cat.; Huron River, Allmendinger Cat.; Ionia Co.; Bay Co., G. M. Bradford; Flint. C. & S.

CALLA L.

*604. **C. palustris** L. Water Arum. Bogs. Frequent. Th.

SPATHYEMA Raf. SYMPLOCARPUS Salisb.

*605. **S. foetida** (L.) Raf. Skunk Cabbage. *Symplocarpus foetidus* Nutt. Th.

ACORUS L.

*606. **A. Calamus** L. Margin of streams. Infrequent. Th.

LEMNACEÆ Dumort. Duckweed Family.

SPIRODELA Schleid.

*607. **S. polyrhiza** (L.) Schleid. Greater Duckweed. Ponds. Common.

LEMNA L. Duckweed. Duck's-meat.

*608. **L. minor** L. Lesser Duckweed. Th. Ponds Blossoms occasionally in June. Common.

609. **L. perpusilla** Torr. Minute Duckweed. Detroit River, D. H. Campbell; Dr. A. B. Lyons. Rare. C. & S.

*610. **L. trisulca** L. Ivy-leaved Duckweed. Ponds. Frequent. C. & S.

WOLFFIA Horkel.

611. **W. Columbiana** Karst. Columbia Wolffia. Abundant on Maple River, Clinton Co.; Ionia Co.; Detroit, J. M. Bigelow.

612. **W. punctata** Griseb. Brazil Wolffia. W. *Brasiliensis* Engelm., not Wedd. Abundant on Maple River, Clinton Co.; Ionia Co.; Detroit, J. M. Bigelow; Bay Co., G. M. Bradford.

XYRIDACEÆ Lindl. Yellow-eyed Grass Family.

XYRIS L.

613. **X. flexuosa** Muhl. Macomb Co.; S. W., Wright Cat.; Hubbardston, Wheeler; Manistee, F. P. Daniels; Vestaburg, C. A. Davis; S. W., H. S. Pepoon. Rare.

614. **X montana** H. Ries. Yellow-eyed Grass. *X. flexuosa pusilla* A. Gray. Lake Superior, Gray's Manual, 6th edition.

ERIOCAULACEÆ Lindl. Pipewort Family.

ERIOCAULON L.

615. **E. septangulare** With. Margin of Long Lake, Cheboygan Co., B. & K.; S. W., Wright Cat.; Macomb Co.; Cooley; Escanaba, E. J. Hill; Alma, C. A. Davis. Infrequent. L. P.

COMMELINACEÆ Reichenb. Spiderwort Family.

COMMELINA L.

616. **C. Virginica** L. Virginia Day-flower. S. W. Wright Cat.

TRADESCANTIA L. Spiderwort.

617. **T. reflexa** Raf. Grass Lake, Cassopolis, Muskegon, C. F. Wheeler; Algonac Co., W. S. Cooper.

618. **T. Virginica** L. Common Spiderwort. Moist wooks. Ionia Co.; Grand Rapids; Ann Arbor, Allmendinger Cat.; Hillsdale Co. Frequent. C. & S.

PONTEDERIACEÆ Dumort. Pickerel-weed Family.

PONTEDERIA L.

*619. **P. cordata** L. Pickerel-weed. Borders of lakes and slow streams. Lakes in Oakland Co.; Ann Arbor; Ionia Co., and northward. Th.

*620. **P. cordata lancifolia** (Muhl.) Morong. *P. cordata angustifolia* Torr. Lake St. Clair; Pine Lake, Ingham Co.

HETERANTHERA R. & P.

*621. **H. dubia** (Jacq.) MacM. Water Star-grass. *H. graminea* Vahl. Ann Arbor, Allmend. Cat.; Dexter, Dr. Elmore Palmer; Grand Rapids; Bay Co.; Hubbardston; Alma, C. A. Davis. In streams. Common. C. & S.

JUNCACEÆ Vent. Rush Family.

JUNCUS L.

*622. **J. acuminatus** Michx. Short-fruited Rush. Macomb Co.; Fruitport, E. J. Hill; Hubbardston; Keweenaw Co., O. A. Farwell; St. Clair Co., C. K. Dodge; Alma, C. A. Davis.

623. **J. articulatus** L. Pointed Rush. Alger, C. F. Wheeler; Tuscola County, C. A. Davis. Th.

624. **J. Balticus littoralis** Engelm. Sandy shores. S. Haven, Bailey; to Petoskey; Port Huron; Bay Co., G. M. Bradford. Oscodo and northward. A form of this species is found at Hubbardston, Ionia Co., Wheeler; Port Austin, C. A. Davis. Th.

625. **J. brachycarpus** Engelm. Short-fruited Rush. St. Clair Co., C. K. Dodge.

*626. **J. brachycephalus** (Engelm.) Buch. Small-headed Rush. *J. Canadensis brachycephalus* Engelm. Hubbardston; Howell Junction, C. F. Wheeler; Bay Co., G. M. Bradford. S.

*627. **J. bufonius** L. Toad Rush. Roadsides. Common. Th.

*628. **J. Canadensis** J. Gay. Canada Rush. Fruitport, E. J. Hill; Macomb Co.; Hubbardston; northward to Sault Ste. Marie, Burgess; Alma, C .A. Davis. Common. Th.

629. **J. Canadensis brevicaudatus** Engelm. *J. Canadensis coarctatus* Engelm. Detroit; north shore of Lake Superior, Dr. Lyons; Keweenaw Co., O. A. Farwell.

*631. **J. conglomeratus** L. Glomerate Rush. *J. effusus conglomeratus* Engelm. Swamps; not so common as the species.

632. **J. Dudleyi** Wieg. Keweenaw Co., Detroit, O. A. Farwell.

*633. **J. effusus** L. Common or Soft Rush. Marshy grounds. Common. Th.

634. **J. filiformis** L. Thread Rush. Adrian, Tuthill; Saginaw Bay, Winch. Cat.; L. Superior, Jno. Macoun. Rare.

635. **J. Gerardi** Loisel. Black-grass. "Rare about the Great Lakes," Gray's Manual.

636. **J. Greenii** Oakes & Tuck. Green's Rush. Head of Lake Michigan, Gray's Manual, 6th edition; Detroit, Wheeler; Keweenaw Co., O. A. Farwell; St. Clair Co., C. K. Dodge. Rare.

637. **J. interior** Wieg. *J. tenuis secundus* Engelm. Palmer Park, Detroit, C. F. Wheeler; Twin Lake, Muskegon Co., C. F. Wheeler.

638. **J. marginatus** Rostk. Grass-leaved Rush. S. Mich., Winch. Cat.; Macomb Co., Dr. D. Cooley; Grand Rapids, Wheeler; Livingston Co., C. A. Davis. Rare. S.

*639. **J. nodosus** L. Knotted Rush. Gravelly banks. Common. An intermediate form grows with the species on the shore of Litle Traverse Bay, Wheeler.

*640. **J. pelocarpus** E. Meyer. Brownish-fruited Rush. Th. Shore of Woodward Lake, Ionia Co.; Pine Lake, Ingham Co.; Lake Superior; J. Macoun, Can. Cat.; Bear Lake, Manistee Co., E. J. Hill; Macomb Co., Dr. D. Cooley; Vestaburg, C. A. Davis. Infrequent.

*641. **J. Richardsonianus** Schult. Richardson's Rush. *J. alpinus insignis* Fries. "Along the Great Lakes northward and westward," Gray. Common. Along railroad track on Agricultural College Farm. This species has been mistaken by early collectors for J. articulatus, which is confined to the New England States. Engelmann. Th.

642. **J. scirpoides** Lam. Scirpus-like Rush. Sturgis, F. P. Daniels.

644. **J. stygius** L. Moor Rush. "N. shore of Lake Superior, Mr. Wheeler," Gray's Manual, 5th edition; Marquette, E. J. Hill.

*645. **J. tenuis** Willd. Slender Rush. Roadside. Common. Th.

646. **J. Torreyi** Coville. Torrey's Rush. *J. nodosus megacephalus* Torr. Flint; Hubbardston and southward; St. Clair Co., C. K. Dodge; Bay Co., G. M. Bradford.

647. **J. Vaseyi** Engelm. Vasey's Rush. Detroit, Lyons; Lake Superior, John Macoun.

JUNCOIDES Adans. LUZULA DC.

*648. **J campestre** (L.) Kuntze. Common Wood-rush. *Luzula campestris* DC.

*649. **J. pilosum** (L.) Kuntze. Hairy Wood-rush. Bay Co., G. M. Bradford. Common. Th.

650. **J. spicatum** (L.) Kuntze. Spiked Wood-rush. *Luzula spicata* DC. Dr. A. B. Lyons. U. P. Woods. Common. Th.

MELANTHACEÆ R. Br. Bunch-flower Family.

TOFIELDIA Hudson.

*651. **T. glutinosa** (Michx.) Pursh. Glutinous Tofieldia. Sphagnous swamps. Frequent. Th.

652. **T. palustris** Hudson. Scottish Asphodel. Isle Royale, Dr. A. B. Lyons. U. P.

CHAMAELIRIUM Willd.

653. **C. luteum** (L.) A. Gray. Blazing Star. *C. Carolinianum* Willd. Dr. A. B. Lyons. U. P.

ZYGADENUS Michx.

*654. **Z. elegans** Pursh. Th. S. W., Wright Cat.; Dexter, Dr. Elmore Palmer; Ann Arbor, Allmend. Cat.; Grand Ledge; Clarkston, G. H. Hicks; Ionia; Davisburg; Petoskey; Ann Arbor, C. A. Davis; Detroit, O. A. Farwell. Rare.

UVULARIA L.

*655. **U. grandiflora** J. E. Smith. Large-flowered Bellwort. Rich woods. Common. C. & S.

656. **U. perfoliata** L. Perfoliata Bellwort. Rich woods. Marquette Co., Burt MS. Cat.; Flint; Macomb Co. Infrequent.

657. **U. sessilifolia** L. Sessile-leave Bellwort. *Oakesia sessilifolia* S. Wats. Low woods. Monroe Co., Wheeler; Ann Arbor, Allmend. Cat.; Ypsilanti; Flint; Macomb Co.; Crystal Lake, Montcalm Co., E. F. Smith, and northward to Marquette Co., Whitney Cat. Th.

LILIACEÆ Adans. Lily Family.

HEMEROCALLIS L.

*658. **H. fulva** L. Common Day-lily. Escaped from cultivation.

ALLIUM L.

*659. **A. Canadense** Kalm. Wild Garlic. Woods. Common. C. & S.

660. **A. cernuum** Roth. Wild Onion. Ann Arbor, Allmend. Cat. S.

661. **A. Sibiricum** L. *A. Schoenoprasum* Ill. Fl., not L. Chives. Dr. A. B. Lyons; Gray's Manual; N. shore of Lake Superior, Agassiz; Keweenaw Point, O. A .Farwell. U. P.

*662. **A. tricoccum** Ait. Wild Leek. Rich woods. Th.

*663. **A. vineale** L. Field Garlic. In the aboretum of the Agricultural College.

LILIUM L.

664. **L. Canadense** L. Wild Yellow Lily. Meadows and along streams. Northward, frequent.

*665. **L. Philadelphicum** L. Wild Orange-red Lily. Ann Arbor, Allmend. Cat.; Hubbardston; Petoskey; and northward. Not common in the central part of the State. There is some doubt about the presence of this plant in Mich.

*666. **L. superbum** L. Turk's-cap Lily. Low grounds. Ann Arbor (Miss Clark), Winch. Cat.; Flint; S. Haven, Bailey; Alma, C. A. Davis. Frequent. C. & S.

667. **L. umbellatum** Pursh. Western Red Lily. Alpena, Traverse City, C. F. Wheeler; St. Clair Co., C. K. Dodge; Bay Co., G. M. Bradford; Keweenaw Co., O. A. Farwell; Birmingham, S. Alexander.

ERYTHRONIUM L.

*668. **E. albidum** Nutt. White Adder's-tongue Violet. Th. Ann Arbor, Allmend. Cat.; Macomb Co.; Flint; Hubbardston. "At L. Superior Dr. Robbins found a plant like this but yellow flowered, a transition toward E. grandiflorum," Gray's Man.; Keweenaw Co., O. A. Farwell.

670. **E. Americanum** Ker. Yellow Adder's-tongue. Low copses. Common. Th.

QUAMASIA Raf. CAMASSIA Lindl.

670. **Q. hyacinthina** (Raf.) Britton. Wild Hyacinth. *Camassia Fraseri* Torr. Adrian, Mrs. I. H. Wheeler; White Island in the Detroit river opposite Amherstburg, Dr. J. Macoun.

MUSCARI Mill.

MUSCARIA.

670a. **M. botryoides** (L.) Mill. *Grape Hyacinth.* Moist grass lands. Ann Arbor, C. A. Davis.

ALETRIS L.

671. **A. farinosa** L. Colic-root Star-grass. Addison, G. F. Comstock; Howard City; Grand Rapids; Ann Arbor, Allmend. Cat.; Macomb Co.; Flint; Clarkston, G. H. Hicks; Hubbardston; Manistee, F. P. Daniels; Ann Arbor, C. A. Davis. Rare. C. & S.

CONVALLARIACEÆ Link. Lily-of-the-Valley Family.

ASPARAGUS L.

*672. **A. officinalis** L. Garden Asparagus. Sparingly escaped from gardens in older parts of the State.

CLINTONIA Raf.

673. **C. borealis** (Ait.) Raf. Yellow Clintonia. Follows the Lake Michigan shore down as far as S. Haven; on the eastern side of the State reaches to Macomb Co., Dr. D. Cooley; and in the center of the State is found in Ionia Co.; Alma, C. A. Davis. Very common north of latitude 43°.

VAGNERA Adans. SMILACINA Desf.

*674. **V. racemosa** (L.) Morong. Wild Spikenard. *Smilacina racemosa* Desf. Moist grounds. Common. Th.

*675. **V. stellata** (L.) Morong. Star-flowered Solomon's Seal. *Smilacina stellata* Desf. Moist banks. Common. Th.

*676. **V. trifolia** (L.) Morong. Three-leaved Solomon's Seal. *Smilacina trifolia* Desf. Sphagnous swamps. Frequent. Th.

UNIFOLIUM Adans. MAIANTHEMUM Wigg.

*677. **U. Canadense** (Desf.) Greene. False Lily-of-the-valley. *Maianthemum Canadense* Desf. Woods, everywhere.

STREPTOPUS Michx.

678. **S. amplexifolius** (L.) DC. Twisted Stalk. Fort Gratiot, Winch. Cat.; Houghton Lake to Lake Superior, where it is rare, Whitney Cat.

679. **S. roseus** Michx. Sessile-leaved Twisted-stalk. Cedar swamps. Mt. Pleasant, Alma, Chas. A. Davis; Cheboygan Co., B. & K.; Bay Co., G. M. Bradford; Drummond's Is., and Sugar Is., Winch. Cat.; to L. Superior where it is very common, Whitney Cat.

SALOMONIA Heist. POLYGONATUM Adans.

*680. **S. biflora** (Walt.) Britt. Smaller Solomon's Seal. *P. biflorum* Ell. Ann Arbor, Winch. Cat.; Ionia Co., and northward. Open woods. Common. Th.

*681. **S. commutata** (R. & S.) Britt. Smooth Solomon's Seal. Th. *P. giganteum* Dietrich. Great S. S. River banks. Stems often very tall and channeled on one side. Intermediate forms between this and the preceding occur. Ann Arbor, C. A. Davis.

MEDEOLA L.

*682. **M. Virginiana** L. Indian Cucumber-root. L. P. Abundant at Gaylord, where it probably reaches its N. limits, G. L. Stewart; Alma and Wayne, C. A. Davis; S. W., H. S. Pepoon.

TRILLIUM L. Wake Robin. Birthroot.

*683. **T. cernuum** L. Nodding Wake-robin. S. Mich., Winch. Cat.; Macomb Co.; Flint; Grand Rapids, Coleman Cat.; northward to Keweenaw, O. A. Farwell; Alma and Ann Arbor, C. A. Davis. Infrequent.

*684. **T. erectum** L. Ill-scented Wake-robin. Ann Arbor, Winch. Cat.; Constantine; Port Huron, Dodge. The white form with declinate pedicels is most common throughout, and the only form in the northern part of the State. Th.

*685. **T. grandiflorum** (Michx.) Salisb. Large-flowered Wake-robin. Rich woods. Exceedingly variable and apt to sport. Common. Th.

686. **T. nivale** Riddell. Dwarf White Trillium. Low woods. One of our earliest spring flowers. Hubbardston; Ionia; Grand Rapids, Coleman Cat., Niles, I. N. Mitchell. Rare. C. & S.

687. **T. recurvatum** Beck. Prairie Wake-robin. Niles, I. N. Mitchell. S. W.

688. **T. sessile** L. Sessile-flowered Wake-robin. Dr. A. B. Lyons; St. Joseph, Wheeler; Niles, I. N. Mitchell. Rare. S.

689. **T. undulatum** Willd. Painted T. *T. erythrocarpum* Michx. S. Mich., Wright Cat.; not observed in the center of the State; Port Huron, Dodge; Keweenaw Co., O. A. Farwell. Infrequent. Th.

SMILACEÆ Vent. Smilax Family.

SMILAX L.

*690. **S. ecirrhata** (Engelm.) S. Wats. Upright Smilax. St. Clair, C. K. Dodge; Belle Isle, O. A. Farwell; Sturgis, F. P. Daniels; Alma, Ann Arbor, C. A. Davis.

*691. **S. herbacea** L. Carrion-flower. River banks. Flowers much visited by blow-flies. Common. Th.

*692. **S. herbacea pulverulenta** (A. Gray.) Michx. Ann Arbor, Allmend. Cat.; Owosso, G. H. Hicks; Alma, C. A. Davis.

*693. **S. hispida** Muhl. Hispid Greenbrier. The common woody species in Ionia and adjacent counties. Ann Arbor, Winch. Cat.; Hubbardston; Flint; Houghton Lake; Alma, Ann Arbor, C. A. Davis; northward to Lake Superior, Whitney Cat.

694. **S. rotundifolia** L. Common Greenbrier. Horse-brier. Ann Arbor, Allmend. Cat.; Flint; Macomb Co.; Kent Co., Coleman Cat.; Benton Harbor. Infrequent. C. & S.

AMARYLLIDACEÆ Lindl. Amaryllis Family.

HYPOXIS L. Star-grass.

695. **H. hirsuta** (L.) Coville. Star-grass. Tuscola county, Ann Arbor, C. A. Davis. *H. erecta* L. Meadows. Common.

DIOSCOREACEÆ Lindl. Yam Family.

DIOSCOREA L.

*696. **D. villosa** L. Wild Yam-root. Rich woods. Frequent. C. & S.

IRIDACEÆ Lindl. Iris Family.

IRIS L.

697. **I. lacustris** Nutt. Dwarf Lake Iris. Bois Blanc I. and Drummond's I., Winch. Cat.; Mackinac, Whitney Cat.; "Shores of L. Huron and Mich.," A. Gray, Lewis Foote; Mackinaw City, Wheeler; Alpena, C. A. Davis.

*698. **I. versicolor** L. Larger Blue Flag. Low grounds. Common. Th.

SISYRINCHIUM L.

*699. **S. albidum** Raf. White Blue-eyed Grass. Barron Lake, C. F. Wheeler; Belle Isle, O. A. Farwell; Kalamazoo, R. M. Gibbs.

*700. **S. angustifolia** Miller. Northern Blue-eyed Grass. Moist grassy places. Common. Th.

701. **S. apiculatum** Bicknell. Muskegon Co., C. D. McLouth.

702. **S. Farwellii** Bicknell. Near Birmingham, Oakland Co., O. A. Farwell.

*703. **S. graminoides** Bicknell. Belle Isle and elsewhere, O. A. Farwell; *S. anceps* S. Wats. Palo, Petoskey, C. F. Wheeler.

704. **S. hastile** Bicknell. Belle Isle, O. A. Farwell.

705. **S. mucronatum** Michx. Michaux's Blue-eyed Grass. Southeast Michigan, O. A. Farwell.

706. **S. strictum** Bicknell. Montcalm Co., Britton's Manual.

ORCHIDACEÆ Lindl. Orchis Family.

CYPRIPEDIUM L.

*707. C. acaule Ait. Stemless Ladies' Slipper. Dry woods and sphagnous swamps. Frequent. Th.

708. C. arietinum R. Brown. Ram's-head Ladies' Slipper. Isle Royale, Dr. A. B. Lyons; tamarack swamp one-half mile east of Whitmore Lake, W. H. Lewis; Mt. Pleasant, C. A. Davis. Rare.

709. C. candidum Willd. Small White Ladies' Slipper. Tamarack swamps. Ann Arbor, Winch. Cat.; Kalamazoo; Macomb Co.; Flint; Hubbardston; Howell Junction, C. F. Wheeler; Keweenaw Co., O. A. Farwell. Rare.

*710. C. hirsutum Mill. Large Yellow Ladies' Slipper. *C. pubescens* Willd. Woods, in moist or dry ground. Frequent.

*711. C. parviflorum Salib. Smaller Yellow Ladies' Slipper. Very wet swamps. Frequent. Th.

*712. C. reginae Walt. Showy Ladies' Slipper. *C. spectabile* Salisb. Swamps. Ann Arbor, Winch. Cat.; Kalamazoo, Tuthill; Macomb Co.; Flint; Hubbardston; Keweenaw Co.; Alma, C. A. Davis. Frequent. Th.

ORCHIS L.

713. O. rotundifolia Pursh. Small Round-leaved Orchis. Frankfort, E. J. Parker; Marquette, T. H. Danger; Lake Fumeé, E. J. Hill.

GALEORCHIS Rydb. ORCHIS L., in part.

*714. G. spectabilis (L.) Rydb. Showy Orchis. *Orchis spectabilis* L. Rich woods. Scarce. C. & S.

PERULARIA Lindl. HABENARIA Willd., in part.

715. P. flava (L.) O. A. Farwell. Tubercled Orchis. *Habenaria flava* (L.) A. Gray. Macomb Co.; Ann Arbor, Winch. Cat.; S. W. Wright Cat.; Cheboygan Co., B. & K.; Bay Co., G. M. Bradford; Alma, C. A. Davis. Infrequent. Th.

COELOGLOSSUM Hartman. HABENARIA Willd., in part.

*716. C. bracteatum (Willd.) Parl. Long-bracted Orchis. *Habenaria bracteata* (Willd.) R. Br. Ann Arbor and Emmet Co., Winch. Cat.; Macomb Co.; Kalamazoo; Hubbardston; Flint; Alma, and northward. Th.

GYMNANDENIOPSIS Rydb. HABENARIA Willd., in part.

717. G. clavellata (Michx.) Rydb. Small Green Wood Orchis. *Habenaria clavellata* (Michx.) Spreng. Lenawee Co., G. F. Comstock; S. W., Wright Cat.; Ann Arbor, Allmend. Cat.; Hubbardston, Grayling, G. H. Hicks; Alma, C. A. Davis; and northward.

LIMNORCHIS Rydb. HABENARIA Willd., in part.

*718. L. dilatata (Pursh.) Rydb. Tall White Bog Orchis. *Habenaria dilatata* (Pursh.) Hook. S. E. Winch. Cat.; Constantine and northward. Th.

719. L. hyperborea (L.) Rydb. Tall Leafy Green Orchis. *Habenaria hyperborea* (L.) R. Br. Wet woods. Frequent northward. Th.

LYSIAS Salisb. HABENARIA Willd., in part.

*720. L. Hookeriana (A. Gray) Rydb. Hooker's Orchis. *Habenaria Hookeriana* A. Gray. S. E. Winch. Cat.; Hubbardston; Flint; Lake Superior, Whitney Cat.; Alma, C. A. Davis. Rare southward. Th.

*721. L. orbiculata (Pursh.) Rydb. Large Round-leaved Orchis. *Habenaria orbiculata* (Pursh.) Torr. Frequent in the pine region, not rare on U. P., Whitney, Cat.; Hubbardston; Alma; Flint. Th.

LYSIELLA Rydb. HABENARIA Willd., in part.

722. **L. obtusata** (Pursh.) Rydb. Small Northern Bog Orchis. *Habenaria obtusata* (Pursh.) Richards. Cheboygan Co., B. & K.; Pictured Rocks, G. H. Hicks; Isle Royale, Dr. A. B. Lyons; L. Superior, common, Whitney Cat.; Cove I.; L. Huron, Austin. N. & U. P.

BLEPHARIGLOTTIS Raf. HABENARIA Willd., in part.

723. **B. blephariglottis** (Willd.) O. A. Farwell. White-fringed Orchis. *Habenaria blephariglottis* (Willd.) Torr. S. Mich., Winch. Cat.; Stanton. Rare. C. & S.

*724. **B. ciliaris** (L.) Rydb. Yellow-fringed Orchis. *Habenaria ciliaris* (L.) R. Br. Ann Arbor, Winch. Cat.; Macomb Co.; Kalamazoo, Tuthill; Niles, I. N. Mitchell; Algonac, W. S. Cooper; Ann Arbor, C. A. Davis. Rare.

725. **B. ciliaris x blephariglottis.** Bay Co., G. M. Bradford.

*726. **B. lacera** (Michx.) Rydb. Ragged Orchis. *Habenaria lacera* (Michx.) R. Br. Ann Arbor, Winch. Cat.; Woodward Lake; Flint; Macomb Co.; Manistee, F. P. Daniels; north to Keweenaw Co., O. A. Farwell. Infrequent. Th.

*727. **B. leucophaea** (Nutt.) O. A. Farwell. Prairie White-fringed Orchis. *Habenaria leucophaea* (Nutt.) A. Gray. Belle Isle, Foerste; Ann Arbor, Winch. Cat.; Pine Lake, Ingham Co.; Woodward Lake, Ionia Co.; Macomb Co.; Alma, C. A. Davis; Isle Royale, Foote; St. Clair Co., C. K. Dodge. Rare.

*728. **B. psycodes** (L.) Rydb. Smaller Purple-fringed Orchis. *Habenaria psycodes* (L.) A. Gray. Low grounds. Frequent. Th.

POGONIA Juss.

*729. **P. ophioglossoides** (L.) Ker. Snake-mouth. Bogs. Common. Th.

ISOTRIA Raf. POGONIA Juss., in part.

730. **I. verticillata** (Willd.) Raf. *Pogonia verticillata* (Willd.) Nutt. Alma, C. A. Davis; Kalamazoo; Flint; Macomb Co. Rare.

TRIPHORA Nutt. POGONIA Juss., in part.

731. **T. trianthophora** (Sw.) Rydb. Nodding Pogonia. *pogonia pendula* Lindl. Dr. Wright; Calvin, Cass Co., I. N. Mitchell; Alma, C. A. Davis. Rare. S. W.

ARETHUSA Gronov.

*732. **A. bulbosa** L. Arethusa. In sphagnous swamps. S. Mich., Wright Cat.; Ann Arbor, Allmend. Cat.; Kalamazoo, Tuthill; Alma, C. A. Davis; Hubbardston, and northward. Rare. Th.

LIMODORUM L. CALOPOGON R. Br.

*733. **L. tuberosum** L. Grass-pink Colopogon. *Calopogon pulchellus* R. Br. Bogs. Alma, Ann Arbor, C. A. Davis. Common. Th.

GYROSTACHYS Pers. SPIRANTHES L. C. Richard.

*734. **G. cernua** (L.) Kuntze. Nodding Ladies' Tresses. *Spiranthes cernua* Richard. Sphagnous swamps. Frequent. Th.

735. **G. gracilis** (Bigel.) Kuntze. Slender Ladies' Tresses. *Spiranthes gracilis* Bigelow. S. W., Wright's Cat.; Macomb Co.; Grand Rapids, Coleman Cat.; Kalamazoo, Tuthill; Long Lake and Pine Plains, Cheboygan Co., B. & K. L. P.

736. **G. plantaginea** (Raf.) Britt. Wide-leaved Ladies' Tresses. *Spiranthes latifolia* Torr. Drummond's Island, common, and S. E., Winch. Cat.; Hubbardston; Flint; Roscommon. Rare in L. P.

737. **G. stricta** Rydb. Hooded Ladies' Tresses. Not *G. Romanzoffiana* as has been supposed. *Spiranthes Romanzoffiana* Chamisso. Borders of Mud Lake, Northport, E. J. Hill; St. Clair Co., A. F. Foerste; Grayling, G. H. Hicks; Tuscola Co., C. A. Davis; Caribou Islet, Porter; northward to L. Superior. Infrequent southward. Th.

LISTERA R. Br.

738. **L. convallarioides** (Sw.) Torr. Round-lipped Twayblade. Not common. Whitney Cat.; Isle Royale, Dr. A. B. Lyons; abundant at Pictured Rocks, G. H. Hicks; Grand

Traverse Co.; Cheboygan Co., B & K.; Keweenaw Co., O. A. Farwell. N. & U. P.

739. **L. cordata** (L.) R. Br. Twayblade. Whitney Cat.; Isle Royale, Dr. A. B. Lyons; Grayling, G. H. Hicks; Keweenaw Co., O. A. Farwell; cedar swamps, Cheboygan Co., B. & K. N. & U. P.

PERAMIUM Salisb. GOODYERA R. Br.

740. **P. Menziesii** (Lindl.) Morong. Menzies' Rattlesnake Plantain. *Goodyera Menziesii* Lindl. Grayling, G. H. Hicks; Boyne Falls, Northport and Frankfort, E. J. Hill; Petoskey; Isle Royale and Traverse Bay, Dr. A. B. Lyons; Keweenaw Co., O. A. Farwell. Rare in Cheboygan Co., B. & K.

*741. **P. pubescens** (Willd.) MacM. Downy Rattlesnake Plantain. *Goodyera pubescens* R. Br. Woods. Frequent. Th.

742. **P. repens ophioides** (Fernald) Heller. Lesser Rattlesnake Plantain. *Goodyera repens* R. Br. Bangor, Van Buren Co., Bailey; Grand Rapids, Coleman Cat.; Roscommon Co., Dr. D. Cooley; Alma, C. A. Davis; Petoskey and northward. Infrequent. Th.

ACHROANTHES Raf. MICROSTYLIS Nutt.

743. **A. monophylla** (L.) Greene. White Adder's-mouth. *Microstylis monophylla* Lindl. Hubbardston; Flint; Macomb Co.; bogs near Long Lake and Black Lake, Cheboygan Co., B. & K. Rare.

744. **A. unifolia** (Michx.) Raf. Green Adder's-mouth. *Microstylis ophioglossoides* Nutt. Ann Arbor, Allmendinger Cat.; Hubbardston; Bay Co., G. M. Bradford; Lenawee Co., F. G. Comstock. Rare.

LEPTORCHIS Thouars. LIPARIS L. C. Richard.

745. **L. liliifolia** (L.) Kuntze. Large Twayblade. *Liparis liliifolia* Richard. S. W. Wright's Cat.

746. **L. Loesellii** (L.) MacM. Fen Orchis. *Liparis Loeselii* Richard. Ann Arbor. Allmendinger Cat.; abundant in a tamarack swamp near Hubbardston; S. Haven, L. H. Bailey; Flint; Macomb Co.; Keweenaw Co., O. A. Farwell; Sault de Ste. Marie, Porter; Alma, C. A. Davis. Th.

CALYPSO Salisb.

747. **C. borealis** (L.) Oakes. Calypso. Forty-mile Point, Presque Isle Co., Winch. Cat.; shores of Higgins Lake, Dr. D. Cooley; Mount Pleasant, E. F. Smith; Grayling, G. H. Hicks; Frankfort, E. J. Parker; Keweenaw Co., O. A. Farwell; Mackinac, Whitney Cat.; L. Superior, J. Macoun. in Can. Cat.

TIPULARIA Nutt.

748. **T. unifolia** (Muhl.) B. S. P. Crane-fly Orchis. *T. discolor* Nutt. Coleman Cat.; N. Mich., D. Cooley; eastern coast of L. Huron, J. Macoun. Our rarest orchid.

APLECTRUM Nutt.

*749. **A. spicatum** (Wallt.) B. S. P. Adam-and-Eve. Putty-root. *A. hiemale* Nutt. Rich woods. Ann Arbor, Allmend. Cat.; Detroit, Gillman; Macomb Co.; Montcalm Co.; Flint; Hubbardston; Grand Rapids, Coleman's Cat.; Niles, Mitchell; Keweenaw Co., O. A. Farwell; Alma, C. A. Davis. Scarce.

CORALLORHIZA R. Br.

*750. **C. Corallorhiza** (L.) Karst. Early Coral-root. *C. innata* R. Br. S. E., Wright Cat.; Park Lake, Clinton Co.; Harmon; Grayling; Frankfort; L. Superior; S. W., H. S. Pepoon. Not rare. Whitney Cat. Th.

*751. **C. multiflora** Nutt. Large Coral-root. Hubbardston; S. W., Wright Cat.; Cheboygan Co., B. & K.; Detroit, O. A. Farwell; Alma, C. A. Davis; northward to Lake Superior. Th.

752. **C. odontorhiza** (Willd.) Nutt. Small-flowered Coral-root. Hubbardston; Flint; Frankfort; Oscoda; northward to Lake Superior, Whitney Cat. Th.

753. **C. striata** Lindl. Striped Coral-root. Frankfort; Comins, Oscoda Co.; abundant at Mackinac; St. Clair Co., C. K. Dodge; Keweenaw Co., O. A. Farwell. Becoming scarce, N. & U. P.

SAURURACEÆ Lindl. Lizard's-tail Family.

SAURURUS L.

*752. **S. cernuus** L. Lizard's-tail. Swamps, river-sides. Common. C. & S.

JUGLANDACEÆ Lindl. Walnut Family.

JUGLANS L.

*753. **J. cinerea** L. Butternut. Low rich woods.
*754. **J. nigra** L. Black Walnut. Becoming scarce from Bay City south. C. & S.

HICORIA Raf. CARYA Nutt.

*755. **H. alba** (L.) Britton. Mocker-nut. *Carya tomentosa* Nutt. Dry Woods. Flint; Grand Rapids, Coleman's Cat.; Three Rivers; Cassopolis, C. F. Wheeler; Ann Arbor, C. A. Davis. Rare. C. & S.
756. **H. borealis** Ashe. Northern Hickory. Belle Isle, Detroit, O. A. Farwell; Rochester, W. A. Brotherton, and probably elsewhere.
757. **H. glabra** (Mill.) Britton. Pig-nut Hickory. *Carya porcina* Nutt. Woods. Alma, Ann Arbor, C. A. Davis. C. & S.
758. **H. laciniosa** (Michx. f.) Sarg. King-nut. *Carya sulcata* Nutt. Monroe Co., White Pigeon, W. J. B.; Teconsha, G. W. Davis; Ann Arbor, C. A. Davis. River bottoms. Rare, except in the extreme south.
759. **H. microcarpa** (Nutt.) Britton. Small-fruited Hickory. *Carya microcarpa* Nutt. Ann Arbor; Cassopolis; Three Rivers. S.
*760. **H. minima** (Marsh.) Britton. Bitter-nut. *Carya amara* Nutt. Common. C. & S.
*761. **H. ovata** (Mill.) Britton. Shag-bark. C. & S. *Carya alba* Nutt. Common. C. & S.
762. **H. villosa** (Sargent) Ashe. Belle Isle, O. A. Farwell.

MYRICACEÆ Dumort. Sweet-Gale Family.

MYRICA L.

763. **M. cerifera** L. Bayberry. Wax-myrtle. S. Mich., Winch. Cat.; Wayne Co., C. A. Davis.
764. **M. Gale** L. Sweet Gale. Swamp near Crooked Lake, Emmet Co.; Harrisville; Manistee, E. J. Hill; Isle Royale, Dr. A. B. Lyons; Roscommon, C. A. Davis. N. & U. P.

COMPTONIA Banks.

764. **C. peregrina** (L.) Coulter. Sweet-fern. *Myrica asplenifolia* L. South, Wr. Cat.; Detroit, Dr. A. B. Lyons; Alma, C. A. Davis. Very common in the center of the State and northward throughout the pine country, of which it is a chracteristic species.

SALICACEÆ Lindl. Willow Family.

POPULUS L.

*765. **P. alba** L. White Poplar. Abele. Occasionally escaped from cultivation.
*766. **P. balsamifera** L. Balsam Poplar. River banks. A small tree in Michigan. Northward. Th.
766a. **P. candicans** Ait. Balm of Gilead. *P. balsamifera candicans* A. Gray. Cultivated, occasional. Indigenous northward.
*768. **P. deltoides** Marsh. Cottonwood. *P. monilifera* Ait. Sometimes a large tree, three feet in diameter. Infrequent. Collected by the Forestry Commission, June, 1888, in Alcona Co. L. P.
*769. **P. dilatata** Ait. Lombardy Poplar. Occasionally escaped from cultivation.
*770. **P. grandidentata** Michx. Large-toothed Aspen. Woods. Common northward, but "rare in U. P." Whitney Cat.; Alma, Ann Arbor, C. A. Davis.

771. P. heterophylla L. Swamp or Downy Poplar. Cass Co., J. H. Roy.
*772. P. tremuloides Michx. American Aspen. Woods and lake shores; most abundant in U. P., Whitney. Common. Th.

SALIX L.

773. S. adenophylla Hook. Tomentose Willow. Beach sand, Lake Michigan, and northward. Petoskey; St. Jo., Dr. Wright in Torr. Herb., Bebb. Hooker's original specimens came from Labrador.
*774. S. alba vitellina (L.) Koch. Golden Willow. Naturalized from Europe.
*755. S. amygdaloides Anders. Peach-leaved Willow. Flint, Dr. D. Clark; Hubbardston; Harrisville; Detroit, O. A. Farwell. Th.
776. S. balsamifera (Hook.) Barratt. Balsam Willow. Flint, Dr. D. Clark; Keweenaw Co., O. A. Farwell; near Port Huron, C. K. Dodge.
777. S. balsamifera lanceolata Bebb. Keweenaw Co., O. A. Farwell.
778. S. balsamifera vegeta Bebb. Keweenaw Co., O. A. Farwell.
*779. S. Bebbiana Sarg. Bebbs' Willow. *S. rostrata* Richardson. Moist or dry ground. Common. Th.
779a. S. Babbiana x petiolaris No. 37 Bebb. Herb. Salicum. Flint, Dr. D. Clark.
*780. S. candida Fluegge. Hoary Willow. Usually in tamarack swamps. Rare in S. part of the State. Common northward. Alma, Ann Arbor, C. A. Davis. Th.
781. S. candida x cordata Bebb. Flint, Dr. D. Clark; Hubbardston, C. F. Wheeler.
*782. S. cordata Muhl. Heart-leaved Willow. Along streams. Narrow-leaved forms occur in the central and southern parts of the State; at Petoskey, broadly ovate-heart-shaped leaved forms occur. Common. Th.
783. S. cordata angustata (Pursh.) Anders. Ypsilanti, Detroit, O. A. Farwell.
*784. S. cordata x sericea Bebb. L. H. Bailey; Flint, Dr. D. Clark.
*785. S. discolor Muhl. Glaucous Willow. River banks. Common.
786. S. eriocephala Michx. Keweenaw Co., O. A. Farwell. *S. discolor eriocephala* Anders.
789. S. fragilis L. Brittle Willow. Planted in cities and villages. Birmingham, S. Alexander.
*790. S. fragilis x alba Wimmer. Coleman's Catalogue; Palmer's Cat.
*791. S. glaucophylla Bebb. Broad-leaved Willow. Hubbardston; Petoskey, frequent along the shores of Little Traverse Bay, and shores of the Great Lakes. Rare in the interior. Th.
792. S. glaucophylla angustifolia Bebb. Sturgeon Point, Alcona Co.
793. S. glaucophylla brevifolia Bebb. Shores of Little Traverse Bay, C. F. Wheeler.
*794. S. humilis Marshall. Prairie Willow. Common northward to Marquette Co., Burt MS. Cat. A very broad-leaved form with the young leaves round obovate to nearly round, collected by O. A. Farwell, Keweenaw Co.; Ann Arbor, C. A. Davis.
*795. S. humilis x discolor Bebb. Keweenaw Co., O. A. Farwell.
795a. S. interior Rowlee. Wet places. Common. Th.
796. S. interior Wheeleri Rowlee. Bay Co., G. M. Bradford; Belle Isle, O. A. Farwell.
*797. S. lucida Muhl. Shining Willow. Along streams. Variable. Common. Th.
*798. S. myrtilloides L. Myrtle Willow. Sphagnous swamps. S. E., Winch. Cat.; Ionia to L. Superior. Frequent. Th.
799. S. myrtilloides pedicellaris Anders. Keweenaw Co., O. A. Farwell.
*800. S. nigra Marshall. Black Willow. Along streams; a small tree. Th.
801. S. nigra falcata (Pursh.) Torr. Bay City and vicinity, G. M. Bradford.
*802. S. petiolaris J. L. Smith. Petioled Willow. With the last. From Sault de Ste. Marie southward. Ionia Co. Frequent. Th.
803. S. petiolaris gracilis Anders. Slender Willow. S. western part of the State.
804. S. petiolaris x candida Bebb. Herb Salicum, No. 30. Originally from Hascall's swamp, near Flint, Mich., where it was discovered by Daniel Clarke, M. D., in 1872—the locality being soon after obliterated. Should it be deemed advisable hereafter to treat supposed hybrids as quasi-species, after the manner of Anderson, Kerner and others, I very much wish that this beautiful willow should be called S. Clarkei, to commemorate the name of a botanist who has done more than any other to give an impetus to the study of hybrid willows in this country, M. S. Bebb, 1880. Swamp near Hubbardston, Ionia Co., C. F. Wheeler.
*805. S. proinoides Pursh. *S. discolor prinoides* (Pursh.) Anders. Common.
*806. S. sericea Marshall. Silky-Willow. Drummond's Is., Winch. Cat.; Ionia Co., and southward to S. Haven, L. H. Bailey; shore of Black Lake, Cheboygan Co. Very common in central part of the State; Alma, Ann Arbor, C. A. Davis.

807. **S. sericea x candida** Bebb., *forma denudata* Bebb. Herb. Salicum; Flint, Dr. Clark.

808. **S. sericea x candida** Bebb. Herb. Salicum, No. 32. Flint, Dr. D. Clark; Hubbardston, C. F. Wheeler.

809. **S. tristis** Ait. Dwarf Gray Willow. Alcona Co.; Barron Lake, Cass Co.; Monroe Co. Infrequent.

*810. **S. viminalis** L. Basket Osier. Woodward Lake, Ionia Co.; S. Haven, L. H. Bailey.

BETULACEÆ Agardh. Birch Family.

CARPINUS L.

*811. **C. Caroliniana** Walter. American Hornbeam. Blue or Water Beech. Along streams. Th.

OSTRYA Scop.

*812. **O. Virginica** (Mill.) Willd. American Hop-Hornbeam. Lever-wood. Rich woods. Common. Th.

CORYLUS L.

*813. **C. Americana** Walt. Wild Hazel-nut. Thickets. Common. Th.

814. **C. rostrata** Ait. Beaked Hazel-nut. Hubbardston; and common northward. St. Clair Co., C. K. Dodge; Alma, C. A. Davis.

BETULA L.

815. **B. glandulosa** Michx. Dwarf Birch. Burt's MS. Cat.; Keweenaw Co., O. A. Farwell. U. P.

*816. **B. lenta** L. Cherry Birch. Sweet or Black Birch. Ann Arbor, Allmend. Cat.; S. Haven, L. H. Bailey; Lenawee Co., W. J. B.; Hubbardston; Flint and northward to L. Superior. Rare in the south, but attains a "monstrous size" on Drummond's Island, Winch. Cat. Th.

817. **B. lutea** Michx. f. Yellow or Gray Birch. Ann Arbor, Winch. Cat.; S. Haven, L. H. Bailey, to L. Superior. Common along the line of the F. & P. M. R. R. and northward to the Traverse country; a large tree south of the Grand-Saginaw valley; Alma, Ann Arbor, C. A. Davis.

*818. **B. papyrifera** Marshall. Paper or Canoe Birch. White Birch. Extends southward to Lansing and perhaps further. Frequent at Crystal Lake, Montcalm Co., as a small tree.

*819. **B. pumila** L. Low Birch. Swamps. Variable. Frequent. Th.

ALNUS Gaertn.

820. **A. Alnobetula** (Ehrh.) K. Koch. Green Alder. *A. viridis* DC. "Dry rocky land," Whitney's Cat.; Isle Royale, Dr. A. B. Lyons; high hills, Escanaba, E. J. Hill; Keweenaw Co., O. A. Farwell. Common. U. P.

*821. **A. incana** (L.) Willd. Speckled or Hoary Alder. Borders of streams. The prevailing alder in center of the State and in U. P. Common.

822. **A. rugosa** (DuRoi) K. Koch. Smooth Alder. *A. serrulata* Willd. Smooth Alder. Macomb Co.; Traverse City and S. Mich., Winch. Cat.; Burt's MS. Cat. Rare or local.

FAGACEÆ Drude. Beech Family.

FAGUS L.

*823. **F. Americana** Sweet. American Beech. *F. ferruginea* Ait. Common in L. P., but rare in U. P. Occurs at Mackinac and Pictured Rocks; St. Mary's River, Macoun. Th.

CASTANEA Adans.

824. **C. dentata** (Marsh.) Borkh. American Chestnut. *C. sativa Americana* S. Wats. Occurs abundantly along an outcrop of Helderberg limestone in E. Monroe Co. and Wayne Co., C. F. Wheeler; Ann Arbor, Dr. Steere; Detroit River to Lake St. Clair, Macoun, in Can. Cat.; St. Clair Co., C. K. Dodge. **S. E.**

QUERCUS L.

*825. **Q. acuminata** (Michx.) Houda. Chestnut or Yellow Oak. *Q. Muhlenbergii* Engelm. Rich woods. A medium sized tree. Alma, Ann Arbor, C. A. Davis. C. & S.

*826. **Q. alba** L. White Oak. Rich woods. Rare in U. P., Menominee Co., Burt. MS. Cat. Common. Th.

827. **Q. Alexanderi** Britton. Alexander's Oak. Birmingham, Oakland Co. S. Alexander; Addison, Lenawee Co., O. C. McLouth. Moist land adapted to swamp white oak, bitternut and American Elm.

828. **Q. borealis** Michx. Gray Oak. *Q. rubra borealis* (Michx.) O. A. Farwell, similar to Red Oak. Keweenaw Co., O. A. Farwell. Referred by Elgelmann to *Q. rubra* L.

*829. **Q. coccinea** Wang. Scarlet Oak. Frequent. L. P.

830. **Q. coccinea x palustris** Hill. Detroit, O. A. Farwell. Frequent.

831. **Q. ellipsoidalis** E. J. Hill. Hill's Oak. A tall tree on moist sandy land. West of Ann Arbor, C. A. Davis.

832. **Q. imbricaria** Michx. Laurel or Shingle Oak. Barrens, Galesburg, H. Dale Adams; Ann Arbor, C. A. Davis; Sturgis, F. P. Daniels; S. Mich., Dr. Wright. Rare.

833. **Q. Leana** Nutt. *Q. imbricaria x velutina* Jackson, S. H. Camp; Ann Arbor, C. A. Davis.

*834. **Q. macrocarpa** Michx. Burr Oak. Over-cup or Mossy-cup Oak. Rich soil. Common. A form of this oak occurs in Cheboygan Co., B. & K. Lower falls of Menominee River, the farthest north of any station in the State, C. F. Wheeler. Th.

835. **Q. palustris** DuRoi. Swamp, Spanish, or Pin Oak. Only seen in the S. portion of the State; Port Huron, C. K. Dodge; Belle Isle; Monroe Co.; Jackson, Marshall, W. J. B.; Algonac, W. S. Cooper; Ypsilanti, C. A. Davis.

*836. **Q. platanoides** (Lam.) Sudw. Swamp White Oak. *Q. bicolor* Willd. Low ground. A large tree. Common. C. & S.

837. **Q. prinoides** Willd. Dwarf Chestnut Oak. A low shrub or small tree. Macomb Co.; Barron Lake, Cass Co.; Brighton, Dr. J. B. Steere; Hubbardston; Muir; Gratiot Co., Washtenaw Co., C. A. Davis. Infrequent. C. & S.

*838. **Q. rubra** L. Red Oak. In the C. & S., a large tree.

839. **Q. Schneckii** Britton. Schneck's Red Oak. *Q. Texana* Sargent. Texas Oak. Wet woods, east of Ypsilanti, C. A. Davis.

*840. **Q. velutina** Lam. Black Oak. L. P. *Q. coccinea tinctoria* A. Gray. This species and Q. coccinea occur together in the C., and are seldom large—usually 40-50 ft., and 12-15 inches in diameter.

ULMACEÆ Mirbel. Elm Family.

ULMUS L.

*841. **U. Americana** L. White or American Elm. Low grounds. Common. Th.

*842. **U. fulva** Michx. Slippery or Red Elm. Rich soil. Frequent. Th.

*843. **U. racemosa** Thomas. Cork or Rock Elm. River banks. Frequent. Th.

CELTIS L.

*844. **C. occidentalis** L. Hackberry. Sugarberry. River banks. Frequent. C. & S.

MORACEÆ Lindl. Mulberry Family.

MORUS L.

845. **M. alba** L. White Mulberry. Ypsilanti, O. A. Farwell; Birmingham, S. Alexander.

*846. **M. rubra** L. Red Mulberry. A small tree on river bottoms. Ann Arbor, C. A. Davis. C. & S.

HUMULUS L.

847. **H. Lupulus** L. Common Hop. Banks of streams. Frequent northward. Th.

CANNABIS L.

*848. **C. sativa** L. Hemp. Waste places. Frequent.

URTICACEÆ Reichenb. Nettle Family.

URTICA L.

849. **U. dioica** L. Great Nettle. Waste places. Bay Co., G. M. Bradford; Manistee, F. P. Daniels. Occasional.

*850. **U. gracilis** Ait. Slender Nettle. Moist ground. Common. Th.

URTICASTRUM Fabr. LAPORTEA Gaud.

*851. **U. divaricatum** (L.) Kuntze. Wood Nettle. *Laportea Canadensis* Gaud. Thick woods along streams. Common. C. & S.

ADICEA Raf. PILEA Lindl.

852. **A. pumila** (L.) Raf. Richweed. *Pilea pumila* A. Gray. Low woods. Common. C. & S.

BOEHMERIA Jacq.

*523. **B. cylindrica** (L.) Willd. False Nettle. Moist ground. Common. C. & S.

LORANTHACEÆ D. Don. Mistletoe Family.

RAZOUMOFSKYA Hoffm.

854. **R. pusilla** (Peck) Kuntze. Small Mistletoe. *Arceuthobium pusillum* Peck. Chatham, C. F. Wheeler; Turin, B. Barlow; south of Cadillac, C. A. Davis. Parasitic on spruces distorting the branches.

SANTALACEÆ R. Br. Sandalwood Family.

COMANDRA Nutt.

855. **C. livida** A. DC. Northern Comandra. "Sandy shores, L. Superior," A. Gray; Traverse City, Dr. A. B. Lyons; Isle Royale, Whitney Cat.; Keweenaw Co., O. A. Farwell. U. P.

*856. **C. umbellata** (L.) Nutt. Bastard Toad-flax. Dry ground. Indifferently parasitic on roots. Common. Th.

ARISTOLOCHIACEÆ Blume. Birthwort Family.

ASARUM L.

*857. **A. acuminatum** (Ashe.) Bicknell. Long-tipped Wild Ginger. Agricultural College, C. F. Wheeler; Keweenaw Co., O. A. Farwell; Alma, W. S. Cooper.

*858. **A. Canadense** L. Wild Ginger. Moist woods. Common.

*859. **A. reflexum** Bicknell. Short-lobed Wild Ginger. Muskegon, C. D. McLouth; Saginaw, W. S. Cooper; Belle Isle, O. A. Farwell.

860. **A. reflexum ambiguum** Bicknell. Keweenaw Co., O. A. Farwell.

ARISTOLOCHIA L.

861. **A. Serpentaria** L. Virginia Snakeroot. Manistee, F. P. Daniels; Detroit, O. A. Farwell. Rare.

POLYGONACEÆ Lindl. Buckwheat Family.

RUMEX L.

862. R. Acetosa L. Sorrel Dock. Very abundant at Point-aux-Pins, above Sault de Ste. Marie, Macoun; N. shore of Lake Superior, Pitcher, Trelease's Revision of Rumex; Bay Co., G. M. Bradford.

*863. R. Acetosella L. Field or Sheep Sorrel. Sterile fields. Common. Th.

*864. R. altissimus Wood. Pale Dock. Sault de Ste. Marie, Winch. Cat.; Ionia; near Ann Arbor, C. A. Davis. Apparently introduced at the College. Rare.

*865. R. Britannica L. Great Water-dock. Wet places. Ann Arbor, C. A. Davis; Ionia Co.; Flint; Macomb Co.; and northward. Frequent.

*866. R. crispus L. Curled Dock. Narrow Dock. Everywhere in fields. Th.

866a. R. obtusifolius L. Broad-leaved Dock. Th.

*867. R. obusifolius x crispus Trelease. North Manitou Isle., Mrs. Wislizenus. Trelease, Revision of Rumex.

*868. R. Patientia L. Patience Dock. Adventitious at Portland, C. F. Wheeler.

869. R. salicifolius Weinm. White Dock. Shore of Little Traverse Bay, and northward. Scarce. N. & U. P.

870. R. sanguineus L. Red-veined Dock. Introduced from Europe.

*871. R. verticillatus L. Swamp Dock. River banks. Frequent. L. P.

FAGOPYRUM Gaertn.

*872. F. Fagopyrum (L.) Karst. Buckwheat. *F. esculentum* Moench. Persistent in fields.

POLYGONUM L.

*873. P. amphibium L. Water Persicaria. Borders of ponds. Frequent. Th.

874. P. arifolium L. Halberd-leaved Tear-thumb. Low grounds. South Haven, L. H. Bailey; Gros. Cap, L. Mich. Winch. Cat.; Bay Co., G. M. Bradford; S. Mich., Wright Cat. Infrequent. L. P.

*875. P. aviculare L. Knot-grass. The commonest of weeds. Th.

876. P. Careyi Olney. Carey's Persicaria. Fort Gratiot. Dr. Z. Pitcher.

877. P. cilinode Michx. Fringed Black Bindweed. Copses. S. Haven, L. H. Bailey; Huron and Roscommon counties, C. A. Davis. Common northward. Th.

*878. P. Convolvulus L. Black Bindweed. Waste grounds. Common. Th.

*879. P. dumetorun L. Detroit, O. A. Farwell.

*880. P. emersum (Michx.) Britton. Swamp Persicaria. *P. Muhlenbergii* S. Wats. East shore of Lake Huron, J. Macoun; Ionia Co.; Flint; Grand Rapids; Indian River; Black Lake; Cheboygan Co.; Alma, Ann Arbor; C. A. Davis; S. W., H. S. Pepoon.

*881. P. erectum L. Erect Knot grass. Waste places. Common.

*882. P. Hartwrightii A. Gray. Hart Wright's Persicaria. Kalamazoo, Tuthill; Indian River; Black Lake, Cheboygan Co., B. and K.; Lenawee Co., W. J. B.; Keweenaw Co., O. A. Farwell; Ann Arbor, C. A. Davis.

*883. P. Hydropiper L. Common Smartweed or Water-pepper. Moist grounds. Common. Th.

*884. P. hydropiperoides Michx. Mild Water-pepper. Wet places. Common. C. & S.

885. P. hydropiperoides Macouni Small. Detroit, O. A. Farwell.

*886. P. incarnatum Ell. Slender Pink Persicaria. Frequent.

*887. P. lapathifolium L. Dock-leaved Persicaria. River banks. Ionia Co.; Grand Rapids, Coleman Cat.; Keweenaw Co., O. A. Farwell. Frequent.

888. P. lapathifolium incanum (Shmidt.) Koch. Keweenaw Co., O. A. Farwell; L. Superior, O. B. Wheeler. Th.

889. P. lapathifolium nodosum (Pers.) Small. Escanaba, C. F. Wheeler; Detroit, O. A. Farwell.

*890. P. littorale Link. Shore Knotweed. Frequent. Th.

*891. P. orientale L. Prince's Feather. Sparingly escaped from gardens.

*892. P. Pennsylvanicum L. Pennsylvania Persicaria. Low grounds. Ionia Co.; Clinton Co.; Flint; S. Mich., Wright Cat. Frequent. C. & S.

*893. P. Persicaria L. Lady's Thumb. Waste places. Common. Th.

*894. P. punctatum Ell. Water Smartweed. *P. acre* H. B. K. Wet places. Ann Arbor; Ionia Co.; Mackinac, Winch. Cat.; Flint. Frequent. L. P.

895. P. ramosissium Michx. Bushy Knotweed. Les Cheneaux Islands, Coryell; Alpena, C. F. Wheeler.

896. P. Rayi Babing. Ray's Knotweed. Belle Isle, O. A. Farwell. Frequent.

*897. P. sagittatum L. Arrow-leaved Tear-thumb. Low grounds. Frequent. Th.

*898. P. scandens L. Climbing False Buckwheat. *P. dumetorum scandens* A. Gray. Moist thickets. Frequent. Th.

*899. P. tenue Michx. Slender Knot-grass. Sterile soil. S. Mich., Wright Cat.; common in Ionia Co.; Macomb Co., and northward.

*900. P. Virginianum L. Virginia Knotweed. Thickets. Common. C. & S.

901. P. viviparum L. Alpine Bistort. Shore of L. Superior, A. Gray; Isle Royale, Whitney Cat.; Keweenaw Co., O. A. Farwell. Common.

POLYGONELLA Michx.

902. P. articulata (L.) Meisn. Coast Jointweed. Traverse City, Winch. Cat.; L. Superior, Whitney Cat.; Oscoda; Black Lake, Cheboygan Co., B. & K.; Indian River, C. F. Wheeler; Harrison, W. J. B.; Crawford Co., O. Palmer; shores of Cable Lake, Berrien Co., H. S. Pepoon. N. & U. P.

CHENOPODIACEÆ Dumort. Goosefoot Family.

CHENOPODIUM L.

*903. C. album L. Lamb's quarters. Pigweed. Waste and cultivated ground. Common. Th.

*904. C. album viride (L.) Moq. Keweenaw Co., Detroit, Ypsilanti, O. A. Farwell.

*905. C. ambrosioides L. Mexican Tea. Waste places. Macomb Co.; Detroit; Ann Arbor, Miss Clark; S. W., Wright. Cat. Scarce. S.

906. C. anthelminticum L. Wormseed. Detroit, O. A. Farwell; Port Huron, C. K. Dodge.

907. C. Bonus-Henricus L. Good King Henry. Flint, Dr. Clark. Infrequent.

*908. C. Botrys L. Jerusalem Oak. Feather Geranium. Escaped from gardens. Abundant at Indian River, C. F. Wheeler; Detroit, O. A. Farwell; Port Crescent, C. A. Davis.

*909. C. glaucum L. Oak-leaved Goosefoot. Detroit and Keweenaw Co., O. A. Farwell; Alma, C. A. Davis.

*910. C. hybridum L. Maple-leaved Goosefoot. Waste grounds. Common. Th.

911. C. leptophyllum (Moq.) Nutt. Narrow-leaved Goosefoot. Bay City, Charlevoix, C. F. Wheeler; Keweenaw Co., O. A. Farwell.

912. C. murale L. Nettle-leaved Goosefoot. Grand Rapids, Coleman Cat.; Ypsilanti and Detroit, O. A. Farwell.

913. C. rubrum L. Red Goosefoot. Grand Rapids, H. C. Skeels; Bay City, G. M. Bradford.

*914. C. urbicum L. Upright Goosefoot. Waste grounds. Ionia Co.; Flint; Grand Rapids, Coleman Cat.; Ann Arbor, C. A. Davis. Infrequent. S.

BLITUM L.

*915. B. capitatum L. Strawberry Blite. *Chenopodium capitatum* Aschers. Rich shady ground. Common. Th.

CYCLOLOMA Moquin.

*916. C. atriplicifolium (Spreng.) Coulter. Winged Pigweed. *C. platyphyllum* Moq. Keweenaw Co., O. A. Farwell; Port Huron, C. K. Dodge; Manistee, F. P. Daniels.

ATRIPLEX L.

*917. A. hastata L. Halberd-leaved Orache. *A. patulum hastatum* A. Gray. Common at Detroit, Dr. A. B. Lyons; Bay Co., G. M. Bradford.

918. A. patula L. Spreading Orache. *A. littoralis* (L.) Sturgis, F. P. Daniels; along the Great Lakes.

CORISPERMUM L.

919. **C. hyssopifolium** L. Bug-seed. Frankfort, C. A. Davis; Detroit, O. A. Farwell; S. Haven, L. H. Bailey; and northward to L. Superior, along the shores of the Great Lakes.

SALSOLA L.

920. **S. Tragus** L. Russian Thistle. *S. Kali Tragus* Moq. Well scattered over the State.

AMARANTHACEÆ J. St. Hil. Amaranth Family.

AMARANTHUS L.

*921. **A. blitoides** S. Wats. Prostrate Amaranth. Lately introduced from the west, spreading rapidly along railroad tracks.

*922. **A. graecizans** L. Tumble-weed. *A. albus* L. Fields and gardens. Frequent. Th.

923. **A. hybridus** L. Slender Pigweed. *A. hypochondriacus* L. *A. chlorostachys* Willd. Scarcely escaped from gardens. Ann Arbor, Miss Clark; Detroit, O. A. Farwell.

*924. **A. retroflexus** L. Rough Pigweed. A common weed in gardens. Th.

925. **A. spinosus** L. Spiny Amaranth. Detroit, O. A. Farwell.

ACNIDA L.

*926. **A. tamariscina concatenata** (Moq.) Uline & Bray. Frequent in Grand River Valley; Detroit, O. A. Farwell.

*927. **A. tamariscina tuberculata** (Moq.) Uline & Bray. Low grounds. S. Mich., Winch. Cat.; Ionia Co.; Macomb Co. Frequent. C. & S.

PHYTOLACCACEÆ Lindl. Pokeweed Family.

PHYTOLACCA L.

*928. **P. decandra** L. Garget. Poke. Scoke. Pigeon-berry. Fields. Frequent. C. & S.

NYCTAGINACEÆ Lindl. Four-o'clock Family.

ALLIONIA Loefl. OXYPHABUS L'Her.

929. **A. hirsuta** Pursh. Hairy Umbrella-wort. *Oxybaphus albidus* Choisy. Grand Rapids, C. W. Follass.

*930. **A. nyctaginea** Michx. Heart-leaved Umbrella-wort. *Oxybaphus nyctagineus* Sweet. Richmond, W. A. Brotherton.

AIZOACEÆ A. Br. Carpet-weed Family.

MOLLUGO L.

*931. **M. verticillata** L. Carpet-weed. Roadsides and sandy fields. Common. C. & S.

PORTULACACEÆ Reichenb. Purslane Family.

CLAYTONIA L.

932. **C. Caroliniana** Michx. Carolina Spring-beauty. Only found in the northern part of the State. Frankfort, E. J. Parker; Mackinac, July, 1888, G. H. Hicks; Keweenaw Co., O. A. Farwell; Alpena, C. A. Davis.

*933. **C. Virginica** L. Spring-beauty. An early and very pretty spring flower. Common. Th.

PORTULACA L.

934. **P. grandiflora** Hook. Garden Portulaca. Ypsilanti, Detroit, O. A. Farwell.

*935. **P. oleracea** L. Purslane. Pussley. Very tenacious of life—a vile weed in gardens. Very common.

CARYOPHYLLACEÆ Reich. Pink Family.

AGROSTEMMA L.

*936. **A. Githago** L. Corn Cockle. *Lychnis Githago* Scop. In wheat fields, but easily eradicated by sowing clean seed-wheat. Common.

SILENE L.

*937. **S. antirrhina** L. Sleepy Catchfly. Common.

938. **S. Armeria** L. Sweet William Catchfly. Baldwin, W. J. B.; Ypsilanti, O. A. Farwell.

*939. **S. noctiflora** L. Night-flowering Catchfly. Frequent in cultivated grounds. Th.

940. **S. stellata** (L.) Aiton. Starry Campion. Dr. Wright; Constantine, C. F. Wheeler. S.

941. **S. Virginica** L. Fire Pink. Catchfly. Winchell Cat.; Keweenaw Co., O. A. Farwell. Open woods, Lake Huron, Todd; islands in Detroit River. Maclagan, Canadian Catalogue. Th.

942. **S. vulgaris** (Moench.) Garcke. Bladder Campion. *S. Cucubalus* Wibel. Port Huron, C. K. Dodge; Manistee, F. P. Daniels.

LYNCHNIS L.

944. **L. alba** Mill. Evening Lynchnis. White Campion. *L. vespertina* Sibth. L. P. Rather common.

*945. **L. Coronaria** (L.) Desr. Mullein Pink. Keweenaw Co.; Grand Traverse, A. B. Lyons; Alma, C. A. Davis.

GYPSOPHILA L.

946. **G. muralis** L. Manistee, F. P. Daniels.

SAPONARIA L.

*947. **S. officinalis** L. Bouncing Bet. Soapwort. Waste places and roadsides. Old Mission, E. J. Hill; Keweenaw Co., O. A. Farwell. Th.

VACCARIA Medic. SAPONARIA L., in part.

*948. **V. Vaccaria** (L.) Britton. Cow-Herb. *Saponaria Vaccaria* L. Muir; S. Mich., Wright Cat.; Keweenaw Co., O. A. Farwell; St. Clair Co., C. K. Dodge. Sparingly introduced. Th.

DIANTHUS L.

949. **D. Armeria** L. Deptford Pink. Midland, E. P. Rice; Rochester, W. A. Brotherton; Detroit, O. A. Farwell.

950. **D. barbatus** L. Sweet William. Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels; St. Clair, C. K. Dodge.

951. **D. deltoides** L. Port Huron, C. K. Dodge.

ALSINE L. STELLARIA L.

952. **A. borealis** (Bigel.) Britton. Northern Stitchwort. *Stellaria borealis* Bigelow. Point au Barques, L. Huron, and Isle aux Train, Gillman; also, Gray in Manual, and Can. Cat.; Isle Royale, A. E. Foote. Infrequent. U. P.

953. **A. borealis alpestris** (Fries.) Britton. *Stellaria borealis alpestris* A. Gray. Alpena, Escanaba, Marquette, C. F. Wheeler.

954. **A. crassifolia** (Ehrh.) Britton. Fleshy Stitchwort. *Stellaria crassifolia* Ehrh. Dr. Lyons; Carson City, C. F. Wheeler. Rare.

*955. **A. graminea** (L.) Britton. Lesser Starwort. *Stellaria gramminea* L. Low ground. Common. Th.

955a. **A. longifolia** (Muhl.) Britton. Long-leaved Stitchwort. Low grounds. Common. Th.

956. **A. longipes** (Goldie) Coville. *Stellaria longipes* Goldie. Long-stalked Stitchwort. Gros Cap, L. Mich., abundant in pure sand, Winch. Cat.; Lake Superior, Dr. A. B. Lyons. Rare. N. & U. P.

957. A. media L. Common Chickweed. *Stellaria media* Cyr. Gardens and fields. A very abundant and hardy little weed. Th.
958. A. uliginosa (Murr.) Britton. Bog Starwort. *Stellaria uliginosa* Murr. Dr. Lyons. Infrequent. U. P.

CERASTIUM L.

959. C. arvense L. Field Chickweed. Lake Superior, Can. Cat.; Mackinac, G. H. Hicks. Infrequent. Th.
960. C. arvense oblongifolium (Torr.) Holl. & Britt. S. Mich., Winch. Cat.; wet woods close to Amherstburgh, Ont., Macoun; Belle Isle, O. A. Farwell.
961. C. longipedunculatum Muhl. Nodding Chickweed. *C. nutans* Raf. Macomb Co.; Flint; Lyons; S. W., H. S. Pepoon. Found on low grounds from Louisiana to Hudson's Bay. Rare.
962. C. semidecandrum L. Keweenaw Co., O. A. Farwell.
963. C. viscosum L. Larger Mouse-ear. Th. Not common.
*964. C. vulgatum L. Mouse-ear Chickweed. Common. Th.

SAGINA L.

965. S. nodosa (L.) Fenzl. Knotted Pearlwort. L. Superior and northward A. Gray; Isle Royale, Whitney's Cat.; Keweenaw Co., O. A. Farwell; Isle Royale, F. E. Wood. U. P.
966. S. procumbens L. Procumbent Pearlwort. Champion, Mich., E. J. Hill.

ARENARIA L. ALSINE Wahl.

*967. A. serpyllifolia L. Thyme-leaved Sandwort. *Alsine serpyllifolia* L. Sandy fields. Common. Th.
967a. A. serpyllifolia tenuior Roch. Keweenaw Co., O. A. Farwell.
968. A. stricta Michx. Rock Sandwort. *Alsine Michauxii* Hook, f. S. Michigan, Wright Cat.; Macomb Co.; Montcalm Co.; L. Sup., Can. Cat.; Constantine; Three Rivers; Alcona Co.; Crystal Lake, Benzie Co. Th.

MOEHRINGIA L. ARENARIA L., in part.

*969. M. lateriflora (L.) Fenzl. *Arenaria lateriflora* L. St. Clair Co., C. K. Dodge.
970. M. macrophylla (Hook.) Torr. Lake Superior, Britton & Brown. *Arenaria macrophylla* Hook.

SPERGULA L.

971. S. arvensis L. Corn Spurrey. Dr. Wright; Dr. Clark; Alma, C. A. Davis. Introduced from Europe.

TISSA Adans. BUDA Adans. SPERGULARIA Pers., in part.

972. T. rubra (L.) Britton. Purple Sandwort. *Spergularia rubra* Presl. Litchfield, W. T. Wallace; Rochester, W. A. Brotherton.

ANYCHIA Michx.

973. A. Canadensis (L.) B. S. P. Norvell, C. F. Wheeler; Jonesville, W. T. Wallace.
974. A. dichotoma Michx. S. Mich., Dr. Wright; Ann Arbor; Watkins Sta., Dr. A. B. Lyons.

SCLERANTHUS L.

*975. S annuus L. Knawel. Naturalized on the Agricultural College grounds; Ann Arbor, C. A. Davis.

NYMPHAEACEÆ DC. Water Lily Family.

BRASENIA Schreber.

*976. B. purpurea (Michx.) Casp. Water-shield. *B. peltata* Pursh. Greenville; Ionia; Ann Arbor; Fife Lake; Oscoda; Manistee, F. P. Daniels; St. Clair, W. S. Cooper; Alma, C. A. Davis; S. W., H. S. Pepoon. Infrequent. L. P.

NYMPHAEA L. NUPHAR Sibth. & Smith.

*977. **N. advena** Soland. Large Yellow Pond Lily. *Nuphar advena* R. Br. In company with water-lilies, but often a dirty plant seeming to delight in filth. Common. Th.

978. **N. advena minor** Morong. Long Lake, Cheboygan Co., B. & K.; Baldwin, W. J. B.

979. **N. Kalmiana** (Michx.) Sims. Small Yellow Pond-Lily. *Nuphar Kalmianum* R. Br. "Sag. Bay & S. W.," Winch. Cat.; S. tier of counties, Wright's Cat.; N. shore of Lake Superior, Agassiz. Rare.

CASTALIA Salisb.

980. **C. odorata** (Dryand.) Woodv. & Wood. Sweet-scented Water-Lily. *Nymphaea odorata* Dryand. Petoskey, E. J. Hill; Vestaburg, C. A. Davis. A form with pink flowers is found in Otsego Lake, Otsego County. Infrequent.

*981. **C. tuberosa** (Paine) Greene. Tuberous White Water-Lily. *Nymphaea tuberosa* Paine. In all our ponds and slow streams. Flowers large and delicately beautiful, fragrant. Th.

NELUMBO Adans.

982. **N. lutea** (Willd.) Pers. Yellow Nelumbo. Water Chinquapin. Lotus. River Rouge, south of Detroit; Mill pond, Vicksburg, Tuthill; River Raisin at Monroe, where it is abundant. Perhaps introduced by the Indians. Local.

CERATOPHYLLACEÆ A. Gray. Hornwort Family.

CERATOPHYLLUM L.

*983. **C. demersum** L. Hornwort. Ponds. Fruit in August. Common. Th.

MAGNOLIACEÆ J. St. Hil. Magnolia Family.

LIRIODENDRON L.

*984. **L. Tulipifera** L. Tulip-tree. Whitewood. A large sized tree, frequent at Ionia, Saranac, Lansing and southward, but not seen north of Grand River Valley. Formerly common but becoming infrequent.

ANONACEÆ DC. Custard-apple Family.

ASIMINA Adans.

*985. **A. triloba** (L.) Dunal. Common Papaw. A low tree, fruit edible. Frequent in the valleys of the Grand and Maple Rivers, whence it probably reaches its northern limit. Common southward. C. & S.

RANUNCULACEÆ Juss. Crowfoot Family.

HYDRASTIS Ellis.

*986. **H. Canadensis** L. Golden Seal. Yellow Puccoon. Rich, moist woods. Rather local. C. & S.

CALTHA L.

*987. **C. palustris** L. Marsh Marigold. In swamps. Frequently called "Cowslip." Very common. Th.

TROLLIUS L.

988. **T. laxus** Salisb. American Globe Flower. Linden, A. W. Chase.

COPTIS Salisb.

*989. **C. trifolia** (L.) Salisb. Three-leaved Goldthread. Bogs and coniferous woods. Common. Th.

ISOPYRUM L.

*990. **I. biternatum** (Raf.) Torr. & Gray. False Rue Anemone. Very common on "beech and maple" land, but not on oak. C. & S.

ACTAEA L.

*991. **A. alba** (L.) Mill. White Baneberry. Cohosh. Moist woods and hillsides. Frequent. Th.

992. **A. eburnea** Rydb. Ivory Baneberry. Keweenaw Co., O. A. Farwell.

*993. **A. rubra** (Ait.) Willd. Red Baneberry. Cohosh. *A. spicata rubra* Ait. Frequent. Th.

CIMICIFUGA L.

994. **C. racemosa** (L.) Nutt. Black Snakeroot. Black Cohosh. Bugbane. S. E., Winch. Cat.; Jackson, University Herb; U. P., Burt. Rare in Michigan.

AQUILEGIA L.

*995. **A. Canadensis** L. Wild Columbine. Frequent. Th.

996. **A. vulgaris** L. European Columbine. Keweenaw Co., O. A. Farwell. Escaped from gardens.

ANEMONE L.

*997. **A. Canadensis** L. Canada Anemone. Wind-flower. *A. Pennsylvanica* L. On low ground along streams; on higher land northward. Common. Th.

*998. **A. cylindrica** A. Gray. Long-fruited Anemone. Poor soil. Infrequent. L. P.

999. **A. Hudsoniana** Richards. Cut-leaved Anemone. Mackinaw, Whitney Cat. Mouth of Saginaw R., Winchell Cat.; Lake Superior, Gray; Frankfort; shore of Grand Traverse Bay, and Torch Lake near the landing, E. J. Hill; eight miles below Rock Harbor, Isle Royal, Porter; Bay Co., Tuscola Co., C. A. Davis. Has been confused with *A. multifida* Poir; a South American species.

1000. **A. parviflora** Michx. Northern Anemone. Dr. Lyons; Asa Gray; Pic River, Lake Superior, Macoun. Rare. U. P.

*1001. **A. quinquefolia** L. Wind-flower. Wood Anemone. *A. nemorosa* Michx. Bay Co., G. M. Bradford. Common. Th.

1002. **A. riparia** Fernald. Rochester and Detroit, O. A. Farwell.

*1003. **A. Virginiana** L. Frequent, preferring sand or gravel. Th.

HEPATICA Scop.

*1004. **H. acuta** (Pursh.) Britton. Sharp-lobed Liver-leaf. This species is very common on beech and maple land, while *Hepatica Hepatica* prefers oak soil. Th.

*1005. **H. Hepatica** (L.) Karst. Round-lobed Liver-leaf. *H. triloba* Chaix. Less frequent than the preceding, at least in the center. Th.

SYNDESMON Hoffmg. ANEMONELLA Spach.

*1006. **S. thalictroides** (L.) Hoffmg. Rue-Anemone. *Anemonella thalictroides* Spach. Quite local through C. & S.; Keweenaw Co., O. A. Farwell; Sturgis, F. P. Daniels; Ann Arbor, C. A. Davis.

PULSATILLA Adans.

1007. **P. hirsutissima** (Pursh.) Britton. Nuttall's Pasque Flower. *Anemone patens Nuttalliana* A. Gray. Norway, S. M. Tobey. Infrequent.

CLEMATIS L.

*1008. **C. Virginiana** L. Common Virgin's Bower. Frequent on low land. Th.

ATRAGENE L.

1009. **A. Americana** Sims. Purple Virgin's Bower. *Clematis verticillaris* DC. Norway, S. M. Tobey. Rare.

RANUNCULUS L.

*1010. **R. abortivus** L. Small-flowered Crowfoot. Common. Th.

*1011. **R. acris** L. Tall Crowfoot or Buttercup. "Classed by Hooker f. as indigenous." S. Watson. So. Haven; Macomb Co.; Flint; Huron shore, Winch. Cat.; and Lake Superior. Gradually spreading over the State. Infrequent. Th.

1012. **R. bulbosus** L. Bulbous Crowfoot or Buttercup. Sault Ste. Marie, Porter; Bay Co., G. M. Bradford; Keweenaw Co., O. A. Farwell. A bad weed in meadows.

*1013. **R. delphinifolius** Torr. Yellow Water-Crowfoot. *R. multifidus* Pursh. Ponds and slow streams. Perennial by rooting from the nodes of floating stems after flowering. The young plants rooting in mud are pubescent so far as observed in many parts of the State. Common. Th.

*1014. **R. fascicularis** Muhl. Early Crowfoot. Flowers sometimes double or with reversion of essential organs to leaves. Ann Arbor, C. A. Davis; Berrien Co., H. S. Pepoon. Hills and sandy plains. Infrequent from the center of the State southward.

1015. **R. Lapponicus** L. Lapland Buttercup. Thunder Bay, Lake Superior, Britton & Brown.

1016. **R. Macounii** Britton. Macoun's Buttercup. *R. hispidus* Hook. Lake Superior, Britton and Brown.

1017. **R. micranthus** Nutt. Rock Crowfoot. *R. obortivus micranthus* A. Gary. North shore of Lake Superior, Agassiz; Keweenaw Co., O. A. Farwell.

1018. **R. obtusiusculus** Raf. Water Plantain Spearwort. *R. ambigens* S. Wats. Dr. Lyons. St. Clair Co., A. F. Foerste and W. S. Cooper.

1019. **R. ovalis** Raf. Prairie Crowfoot. *R. rhomboideus* Goldie. On light sand; Muir and Palo in Ionia County; Lake Superior, Can. Cat. Prairies, Mich. A. Gray. Rare.

1020. **R. Pennsylvanicus** L. f. Bristly Crowfoot. Frequent on low land. Th.

1021. **R. Purshii** Richards. Pursh's Buttercup. Topinabee, S. H. Camp; Chandler's Falls, Atlanta; Alpena, C. F. Wheeler.

*1022. **R. recurvatus** Poir. Hooked Crowfoot. Woods in rich soil. Common. Th.

*1023. **R. repens** L. Creeping Buttercup. Port Huron, C. K. Dodge; Rochester, W. A. Brotherton and O. A. Farwell; Montreal River, Keweenaw Co., O. A. Farwell; in the lawn Agricultural College, C. F. Wheeler.

1024. **R. reptans** L. Creeping Spearwort. *R. Flammula reptans* E. Meyer. Infrequent except northward, and not seen in the center of the State. Sandy shore of Black Lake, Cheboygan Co., B. & K.; Belle Isle, O. A. Farwell; Algonac, W. S. Cooper. N. U. P.

1025. **R. reptans intermedius** (Hook.) Torr. & Gray. *R. Flammula intermedius* Hook. Keweenaw Co., O. A. Farwell.

*1026. **R. sceleratus** L. Cursed Crowfoot. Ditch Crowfoot. Ditches and low ground. Exceedingly variable, stems sometimes two inches in diameter. Frequent. Th.

*1027. **R. septentrionalis** Poir. Swamp Buttercup. Frequent and variable. Th.

BATRACHIUM S. F. Gray.

*1028. **B. divaricatum** (Schrank) Wimm. Stiff Water-Crowfoot. *Ranunculus circinatus* Sibth. Bear River, Petoskey, E. J. Hill; Ann Arbor, Allmendinger Cat.; Huron R., Lyons. The common form in L. P. Frequent. Th.

1029. **B. trichophyllum** (Chaix.) Bossch. White Water-Crowfoot. *Ranunculus aquatilis trichophyllus* A. Gray. Common. Not observed in E. and S. Black Lake, Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell, St. Clair River and Lake, W. S. Cooper.

OXYGRAPHIS Bunge.

1030. **O. Cymbalaria** (Pursh.) Prantl. Seaside Crowfoot. *Ranunculus Cymbalaria* Pursh. South Haven, L. H. Bailey. Rare.

THALICTRUM L.

*1031. **T. dioicum** L. Early Meadow-Rue. Common along river banks. Th.

1032. **T. polygamum** Muhl. Tall Meadow-Rue. Keweenaw Co., O. A. Farwell.

*1033. **T. purpurascens** L. Purplish Meadow-Rue. Wet meadows. Common. Th.

NIGELLA L.

1033a. **N. Damascena** L. Fennel-flower. Escaped from gardens. Infrequent.

BERBERIDACEÆ T. G. Barberry Family.

BERBERIS L.

*1034. **B. vulgaris** L. Common Barberry. Sometimes escaped from cultivation; Flint, Dr. Clark; Tuscola Co., Ann Arbor, C. A. Davis; Bay Co., G. M. Bradford.

CAULOPHYLLUM Michx.

*1035. **C. thalictroides** (L.) Michx. Pappoose-root. Blue Cohosh. Common in L. P.

JEFFERSONIA Barton.

*1036. **J. diphylla** (L.) Persoon. Rheumatism-root. Twin-leaf. Infrequent. C. & S.

PODOPHYLLUM L.

*1037. **P. pelatum** L. May-Apple. Wild Mandrake. Very common. C. & S.

MENISPERMACEÆ DC. Moonseed Family.

MENISPERMUM L.

*1038. **M. Canadense** L. Canada Moonseed. Woods and moist thickets. Frequent.

LAURACEÆ Lindl. Laurel Family.

SASSAFRAS Nees & Eberm.

*1039. **S. Sassafras** (L.) Karst. Sassafras. *S. officinale* Nees & Eberm. Woods, sandy soil, Manistee, F. P. Daniels; Hamlin Lake, Mason Co., C. E. St. Johns; Bay Co., G. M. Bradford. Frequent. L. P.

BENZOIN Fabric.

*1040. **B. Benzoin** (L.) Coulter. Spice-bush. *Lindera Benzoin* Blume. Damp woods. Frequent. C. & S.

PAPAVERACEÆ B. Juss. Poppy Family.

Including also **Fumariaceæ.**

PAPAVER L.

1041. **P. somniferum** L. Common Poppy. Opium Poppy. Occasional. Escaped from cultivation.

SANGUINARIA L.

*1042. **S. Canadensis** L. Bloodroot. Common. Th.

STYLOPHORUM Nutt.

1043. **S. diphyllum** (Michx.) Nutt. Celandine Poppy. Rich woods, Oceana County and southward. Local. C. & S.

CHELIDONIUM L.

*1044. **C. majus** L. Celandine. Lebanon, Clinton Co., Van Vleck; Sturgis, F. P. Daniels.

BICUCULLA Adans. DICENTRA Bernh.

*1045. **B. Canadensis** (Goldie) Millsp. Squirrel Corn. *Dicentra Canadensis* Walp. From Frankfort southward. Frequent. L. P.

*1046. **B. Cucullaria** (L.) Millsp. Dutchman's Breeches. *Dicentra cucullaria* Torr. Frankfort, southward. Frequent. L. P.

ADLUMIA Raf.

1047. **A. fungosa** (Ait.) Greene. Climbing Fumitory. *A. cirrhosa* Raf. Hemlock woods in vicinity of Houghton Lake; Grand Rapids, Miss Clark; So. Haven; Marquette, E. J. Hill. Not common. Th.

CAPNOIDES Adans. CORYDALIS Vent.

1048. **C. aureum** (Willd.) Kuntze. Golden Corydalis. *Corydalis aurea* Willd. Sturgis, F. P. Daniels. Frequent northward in the U. P. Common.

1049. **C. sempervirens** (L.) Borck. Pale Corydalis. *Corydalis glauca* Pursh. Grand Haven; Clare Co.; L. Sup. Infrequent except northward. Oscoda; Alger's camp, Alcona Co. Th.

FUMARIA L.

1050. **F. officinalis** L. Fumitory. Escaped from cultivation at Ypsilanti. O .A. Farwell.

CRUCIFERÆ B. Juss. Mustard Family.

LEPIDIUM L.

*1051. **L. apetalum** Willd. Apetalous Pepper-grass. *L. intermedium* A. Gray. Alma, Ann Arbor, C. A. Davis; Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels.

1052. **L. campestre** (L.) R. Br. Field or Cow Cress. Bay Co., G. M. Bradford; Macomb Co., and Detroit. Infrequent. S. E.

1053. **L. sativum** L. Garden Pepper-grass. Port Huron, C. K. Dodge. Escaped from cultivation.

*1054. **L. Virginicum** L. Wild Pepper-grass. Alma, Ann Arbor. Th.

IBERIS L.

1055. **L. amara** L. Escaped from gardens. Keweenaw Co., O. A. Farwell.

CONRINGIA Link.

1056. **C. orientalis** (L.) Dumort. A bad weed in grain fields, introduced into northern Michigan from the northwest. O. A. Farwell.

THLASPI L.

*1057. **T. arvense** L. Field Pennycress. Ann Arbor. Allmendinger Cat.; "shores of Lake Huron," A. Gray; Monroe, C. A. Davis; Bay Co., G. M. Bradford.

SISYMBRIUM L.

1058. **S. altissimum** L. Tall Sisymbrium. A bad weed from Europe introduced into the Canadian Northwest. Benton Harbor in 1896, C. F. W.; later in many localities. Tuscola Co., C. A. Davis; near Detroit, O. A. Farwell.

1059. **S. humile** Meyer. Northern Rock-cress. Isle Royale, Gillman; Keweenaw Co., O. A. Farwell; Mackinac, G. H. Hicks; Macomb Co.; Grand Haven. Infrequent.

*1060. **S. officinale** (L.) Scop. Hedge Mustard. Road-sides and wet places. Frequent. Th.

CAKILE Gaertn.

1061. **C. edentula** (Bigel.) Hook. Sea-rocket. *C. Americana* Nutt. Shores of the great lakes. Common. Th.

SINAPIS L.

*1062. **S. alba** L. White Mustard. *Brassica alba* Boiss. Infrequent. Th.

BRASSICA L.

*1063. **B. arvensis** (L.) B. S. P. Charlock. *Brassica Sinapistrum* Boiss. A bad weed, becoming too frequent. Th.

1064. **B. campestris** L. Turnip. Keweenaw Co., O. A. Farwell.

*1065. **B. juncea** (L.) Cosson. Indian Mustard. Lapeer, Mrs. M. Owen; Keweenaw Co., O. A. Farwell; Port Huron, W. S. Cooper.

*1066. **B. nigra** (L.) Koch. Black Mustard. Common. Th.

DIPLOTAXIS DC.

*1067. **D muralis** (L.) DC. Sand Rocket. Grand Rapids, H. C. Skeels.

RAPHANUS L.

1067a. **R. Raphanistrum** L. White Charlock. Ballast grounds. Detroit, O. A. Farwell.

1067b. **R. satinus** L. Garden Radish. Persists for some time in old gardens and waste places.

BARBAREA R. Br.

1069. **B. Barbarea** (L.) MacM. Yellow Rocket or Cress. *B. vulgaris* R. Br. South Haven; Ann Arbor, Allmendinger Cat.; Macomb Co.; Alma, C. A. Davis; Hubbardston; Bay Co., G. M. Bradford; and N. into the U. P. where it is indigenous and frequent. Th.

1070. **B. praecox** (J. E. Smith) R. Br. Belle Isle Cress. Sandy ground. St. Clair Co., C. K. Dodge.

1071. **B. stricta** Andrz. Erect-fruited Winter Cress. *B. vulgaris stricta* A. Gray. Keweenaw Co., O. A. Farwell; St. Clair Co. near Capac, C. K. Dodge; Detroit, W. S. Cooper.

RORIPA Scop. NASTURTIUM R. Br.

1072. **R. Americana** (A. Gray) Britton. Lake Water-cress. *Nasturtium lacustre* A. Gray. From Fish Creek and Maple River southward. Infrequent. C. & S.

*1073. **R. Armoracia** (L.) Hitchcock. Horseradish. *Nasturtium Armoracia* Fries. Escaped from gardens into waste places. Frequent.

*1074. **R. hispida** (Desv.) Britton. Hispid Yellow Cress. *Nasturtium palustre hispidum* A. Gray. Alma, Ann Arbor, C. A. Davis. Th.

*1075. **R. Nasturtium** (L.) Rusby. True Water-cress. *Nasturtium officinale* R. Br. Ionia; South Haven; Ann Arbor, Alma, C. A. Davis. Frequent in brooks.

1076. **R. obtusa** (Nutt.) Britton. Blunt-leaved Yellow Cress. *Nasturtium obtusum* Nutt. Keweenaw Co., O. A. Farwell.

*1077. **R. palustris** (L.) Bess. Yellow Water-cress. *N. palustre* DC. Frequent. Th.

1078. **R. sylvestris** (L.) Bess. Creeping Yellow Water-cress. *Nasturtium sylvestre* R. Br. Detroit; O. A. Farwell.

CARDAMINE L.

*1079. **C. bulbosa** (Schreb.) B. S. P. Bulbous Cress. *C. rhomboidea* DC. Common. Th.

*1080. **C. hirsuta** L. Small Bitter Cress. New Buffalo, C. F. Wheeler; near Black Lake, Cheboygan Co., B. & K.; Belle Isle, O. A. Farwell; Alma, C. A. Davis. Th.

1081. **C. parviflora** L. Small-flowered Bitter-cress. New Buffalo, C. F. W.; Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell; St. Clair Co., J. W. Stacey.

1082. **C. Pennsylvanica** Muhl. Pennsylvania Bitter-cress. Keweenaw Co., O. A. Farwell; Bay Co., G. M. Bradford.

*1083. **C. pratensis** L. Cuckoo-Flower. Bogs. Rare S., frequent in C., and common N. Th.

*1084. **C. purpurea** (Torr.) Britton. Purple Cress. *C. rhomboidea purpurea* DC. An early spring flower. Th.

MATTHIOLA L.

1085. **M. bicornis** (B. & S.) DC. Growing in sand along shore of Lake Huron. U. S. Dept. of Agriculture reported it as the only place in the country where it had escaped.

DENTARIA L.

*1086. **D. diphylla** Michx. Two-leaved Toothwort. Common. Th.
*1087. **D. laciniata** Muhl. Cut-leaved Toothwort. Keweenaw Co. and southward. Common. Th.
1088. **D. maxima** Nutt. Large Toothwort. Bluffs along Black River near Abbottsford, C. K. Dodge; the only station known in the State.

BURSA Weber. CAPSELLA Medic.

*1089. **B. Bursa-pastoris** (L.) Britton. Shepherd's Purse. *Capsella Bursa-pastoris* Medic. The commonest of weeds. Th.

CAMELINA Crantz.

*1090. **C. microcarpa** Andrz. Small-fruited False Flax. A weed which is becoming naturalized from Europe.
1091. **C. sativa** (L.) Crantz. Gold-of-Pleasure. False Flax. Road-sides and waste places. Flint; Ann Arbor; Macomb Co.; Sturgis; Bay Co.; Keweenaw Co., F. Not common. Th.

DRABA L.

1092. **D. Caroliniana** Walt. Carolina Whitlow-grass. Ionia Co. and southward. Rare. C. & S.
1093. **D. incana arabisans** (Michx.) S. Wats. Shores of Great Lakes. Mackinac, G. H. Hicks. Infrequent.
*1094. **D. nemorosa** L. Wood Whitlow-grass. "Fort Gratiot and northwestward." A. Gray. Collected by Dr. Pitcher. Infrequent.
*1095. **D. verna** L. Whitlow-grass. S. W., H. S. Pepoon; Detroit, Dr. Lyons. Rare. S.

SOPHIA Adans.

1096. **S. pinnata** (Walt.) Britt. Tansy Mustard. *Sisymbrium canescens* Nutt. Shores of Great Lakes, Pitcher, Houghton, Winchell; Barry Co., L. H. Bailey. Infrequent. Th.
1096a. **S. pinnata brachycarpa** (Richards) O. A. Farwell. Belle Isle, O. A. Farwell.
1097. **S. Sophia** (L.) Britton. Flixweed. Keweenaw Co., O. A. Farwell.

ARABIS L.

1098. **A. brachycarpa** (T. & G.) Britton. Purple Rock-cress. *A. confinis brachycarpa* Watson and Coulter. Keweenaw Co., O. A. Farwell; Marquette, Alpena, Hubbardston, C. F. Wheeler. Not common.
*1099. **A. Canadensis** L. Sickle-pod. Fields and rocky woods. Not common.
*1100. **A. dentata** T. & G. Toothed Rock-cress. Low lands along Grand River and southward. Not common. C. & S.
1100a. **A. Drummondii** A. Gray. *A. confinis* S. Wats., in part. Keweenaw Co., O. A. Farwell.
*1101. **A. glabra** (L.) Bernh. Tower Mustard. *A. perfoliata* Lam. Alma, Ann Arbor, C. A. Davis. Fields. Infrequent. Th.
*1102. **A. hirsuta** (L.) Scop. West Bay City, G. M. Bradford.
1103. **A. Holboellii** Hornem. Holboell's Rock-cress. Thunder Bay, Alpena Co., the most easterly known station for this western species, C. F. Wheeler.
*1104. **A. laevigata** (Muhl.) Poir. Smooth Rock-cress. River banks. Infrequent. Th.
1105. **A. lyrata** L. Lyre-leaved Rock-cress. Sandy fields, Hubbardston; South Haven; Flint; Petoskey; U. P., Whitney Cat. Infrequent. Th.
1106. **A. lyrata occidentaliss** S. Wats. Bower's Harbor, Grand Traverse Bay, Alpena, C. F. Wheeler.

ERYSIMUM L.

*1107. **E. cheiranthoides** L. Worm-seed Mustard. Gratiot Co.; Grayling, G. H. Hicks; Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell; Port Huron, C. K. Dodge; Alma, Ann Arbor, C. A. Davis.
*1108. **E. inconspicuum** (S. Wats.) MacM. Small Erysimum. *E. parviflorum* Nutt. Probably introduced from the west. Clifton, Keweenaw Co., O. A. Farwell; St. Clair Co., C. K. Dodge.

ALYSSUM L.

*1109. **A. alyssoides** (L.) Gouan. Yellow Alyssum. *A. calycinum* L. Hubbardston, Ionia, Flint, C. F. Wheeler; Alma, Ann Arbor, C. A. Davis; Bay Co., G. M. Bradford. A recently introduced plant, becoming common in many localities. Door-yards and waste places.

HESPERIS L.

1110. **H. matronalis** L. Dame's Violet. Escaped from cultivation at Cassopolis, C. F. Wheeler; shore of Lake Huron, T. & G.; Ann Arbor, C. A. Davis; Keweenaw Co., Ypsilanti, O. A. Farwell.

CAPPARIDACEÆ Lindl. Caper Family.

CLEOME L.

1111. **C. serrulata** Pursh. Pink Cleome. *C. integrifolia* T. & G. Vestaburg, C. A. Davis. At one time introduced in several places near the Agricultural College, but now apparently extinct.

POLANISIA Raf.

1112. **P. graveolens** Raf. Clammy-weed. Shores of Great Lakes; Bay Co., G. M. Bradford; South Haven; Detroit; Put-in-Bay. Not rare.

RESEDACEÆ S. F. Gray. Mignonette Family.

RESEDA L.

1113. **R. lutea** L. Yellow cut-leaved Mignonette. Jackson, S. H. Camp.

SARRACENIACEÆ La Pyl. Pitcher-Plant Family.

SARRACENIA L.

*1114. **S. purpurea** L. Side-saddle Flower. Huntsman's Cup. Pitcher-plant. Common in sphagnous swamps. Th.

1115. **S. purpurea heterophylla** (Eaton) Torr. Keweenaw Co., O. A. Farwell.

DROSERACEÆ S. F. Gray. Sundew Family.

DROSERA L.

1116. **D. intermedia** Hayne. Spatulate-leaved Sundew. ***D. intermedia*** var. *Americana* DC. S. Mich., Wright. Cat.; Ann Arbor, Allmendinger Cat.; Grand Rapids, R. H. Wolcott; Keweenaw Co., O. A. Farwell; Vestaburg, C. A. Davis. Th.

1117. **D. linearis** Goldie. Slender Sundew. Livingston Co.; L. Superior, Dr. Lyons; Keweenaw Co., O. A. Farwell; La Pointe, D. Houghton; Copper Harbor, F. E. Wood. Rare.

1118. **D. longifolia** L. Oblong-leaved Sundew. Mud Lake, Oakland Co., W. A. Brotherton.

*1119. **D. rotundifolia** L. Round-leaved Sundew. In sphagnous swamps. An insectivorous plant. Th.

PODOSTEMACEÆ Lindl. River-Weed Family.

PODOSTEMON Michx.

1120. **P. Ceratophyllum** Michx. River-weed. Thread-foot. Detroit, Dr. A. B. Lyons.

CRASSULACEÆ DC. Orpine Family.

SEDUM L.

*1121. **S. acre** L. Mossy Stone-crop. Escaped from cultivation; Alma, Whitmore Lake, C. A. Davis; Bay Co., G. M. Bradford; Keweenaw Co., O. A. Farwell.
*1122. **S. Telephium** L. Live-forever. Garden Orpine. Escaped from cultivation. Th.

PENTHORACEÆ Rydb.

PENTHORUM L.

*1123. **P. sedoides** L. Ditch Stonecrop. Roadsides and ditches; shore of Black Lake, Cheboygan Co., B. & K. Frequent. L. P.

SAXIFRAGACEÆ Dumort. Saxifrage Family.

SAXIFRAGA L.

1124. **S. autumnalis** L. Yellow Mountain Saxifrage. *S. aizoides* L. Northern Michigan, A. Gray; Britton & Brown. U. P.
1125. **S. Aizoon** Jacq. Livelong Saxifrage. Upper Mich., Gray; Isle Royale, Lyons, and Whitney. U. P.
*1126. **S. Pennsylvanica** L. Swamp Saxifrage. Bogs. Common. Th.
1127. **S. tricuspidata** Retz. Three-toothed Saxifrage. Shore of L. Superior and northward, A. Gray; Isle Royale, Dr. Lyons and Whitney Cat. U. P.
1128. **S. Virginiensis** Michx. Early Saxifrage. L. Superior, Can. Cat.; Keweenaw Co., O. A. Farwell. U. P.

TIARELLA L.

*1129. **T. cordifolia** L. False Mitrewort. Birmingham, Belle Isle, O. A. Farwell; Ft. Gratiot; Macomb Co.; Flint; Stanton, and northward. Common in N. & U. P.; rare in C. & S. W. Th.

HEUCHERA L.

1130. **H. Americana** L. Common Alum-root. Grand Rapids, Coleman; Ann Arbor, Winch. Cat.; Monroe Co., C. F. Wheeler; Ypsilanti, O. A. Farwell. C. & S.
*1131. **H. hispida** Pursh. Common in Grand-Saginaw Valley; Ypsilanti, O. A Farwell C. & S.

MITELLA L.

*1132. **M. diphylla** L. Mitre-wort. Hillsides in rich woods. Very common. Th.
*1133. **M. nuda** L. Shaded swamps in Sphagnum. Frequent. Th.

CHRYSOSPLENIUM L.

*1134. **C. Americanum** Schwein. South Haven; Hubbardston; Flint; Alma, C. A. Davis. Northward. Infrequent.

PARNASSIACEÆ Dumort. Grass-of-Parnassus Family.

PARNASSIA L.

*1135. **P. Caroliniana** Michx. Alma, C. A. Davis. Swamps and wet banks. Rare northward; common in C. & S.
1136. **P. palustris** L. Drummond's Is., Winch. Cat.; "shore of L. Superior and northward," A. Gray, from Dr. Pitcher; Harbor Point, C. A. Davis; Keweenaw Co., O. A. Farwell.
1137. **P. parviflora** DC. L. Sup., Can. Cat.; Grand Island, Henry Gillman; Harbor Point, C. A. Davis; also, northwest shore of Lake Michigan, in Wisconsin, Gillman. U. P.

GROSSULARIACEÆ Dumort. Gooseberry Family.

RIBES L.

1138. **R. aureum** Pursh. Permanently established on shore of Lake Huron, C. K. Dodge.

*1139. **R. Cynosbati** L. Gooseberry. Common. Th.

*1140. **R. floridum** L'Her. Wild Black Currant. Cheboygan Co., B. & K. Southward. Common. Th.

*1141. **R. gracile** Michx. Slender Gooseberry. St. Joseph's Island, and Sitting Rabbit, Winchell's Cat.; Chandler's, C. F. Wheeler.

1142. **R. lactustre** (Pers.) Poir. Swamp Gooseberry. As far south as Houghton Lake; Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell.

*1143. **R. oxyacanthoides** L. Swamp Gooseberry. Low grounds along Fish Creek and Maple River; Flint; Macomb Co., northward and shore of Long Lake, B. & K.; Keweenaw Co., O. A. Farwell; Alma, Ann Arbor, C. A. Davis; Bay Co., G. M. Bradford.

1144. **R. prostratum** L'Her. Fetid Currant. Flint; Isabella Co.; Missaukee Co.; St. Joseph's Is., Winch. Cat.; Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell. Frequent. C. N. & P.

1145. **R. rotundifolium** Michx. Thunder Bay Island, C. F. Wheeler.

*1146. **R. rubrum** L. Red Currant. Deep swamps and cold woods, under tamaracks. Ann Arbor; Ionia; Stanton; Cheboygan Co., B. & K. Infrequent. Th.

HAMAMELIDACEÆ Lindl. Witch Hazel Family.

HAMAMELIS L.

*1147. **H. Virginiana** L. Common. Th.

PLATANACEÆ Lindl. Plane-Tree Family.

PLATANUS L.

*1148. **P. occidentalis** L. Along our rivers. "The largest tree of the Atlantic forests." Sargent. C. & S.

ROSACEÆ B. Juss. Rose Family.

OPULASTER Medic. PHYSOCARPA Raf.

1149. **O. opulifolius** (L.) Kuntze. Ninebark. *Physocarpus opulifolius* Maxim. Along streams. Th.

SPIRAEA L.

*1150. **S. salicifolia** L. Common Meadow-Sweet. In marshes. Common. Th.

1151. **S. sorbifolia** L. Port Huron, C. K. Dodge.

*1152. **S. tomentosa** L. Hardhack. Steeple-bush. Grand Rapids, G. D. Sones; north of the College about three and one-half miles, W. J. B.; Sturgis, F. P. Daniels; Ann Arbor, C. A. Davis; Ionia Co., and southward. Infrequent. C. & S.

PORTERANTHUS Britton. GILLENIA Moench.

1152a. **P stipulatus** (Muhl.) Britton. Near Detroit, O. A. Farwell.

1153. **P. trifoliatus** (L.) Britton. Indian Physic. *Gillenia trifoliata* Moench. Bowman's Root. Winchell Cat. Rare. S.

RUBUS L.

*1154. **R. Americanus** (Pers.) O. A. F. Dwarf Raspberry. *R. triflorus* Richard. Frequent. Th.

1154a. **R. argutus** Link. Tall Blackberry. Keweenaw Co., O. A. Farwell.

*1155. **B. Baileyanus** Britton. Belle Isle, O. A. Farwell.

*1156. **R. Canadensis** L. Dewberry. Low Blackberry. *R. Millspaughii* Britt. Lake, Cheboygan Co., C. F. Wheeler; Alma, Ann Arbor, C. A. Davis; Keweenaw Co., O. A. Farwell. Black. Frequent. Th.

1157. **R. Ensleni** Tratt. Enslen's Dewberry. Detroit and vicinity, O. A. Farwell.

1157a. **R. frondosus** Bigel. Leafy-bracted Blackberry. *R. villosus frondosus* Bigel. Keweenaw Co., and Detroit, O. A. Farwell.

*1158. **R. hispidus** L. Running Swamp-B. Very abundant through the central in the pine country. Th.

*1160. **R. neglectus** Peck. Hubbardston; Birmingham; Belle Isle; Alma, C. A. Davis.

1161. **R. nigricans** Rydb. Peck's Dewberry. Detroit and vicinity, O. A. Farwell.

*1161a. **R. nigrobaccus** Bailey. High Bush Blackberry. *R. villosus* A. Gray. Very common in Central Mich., rare in U. P.

*1162. **R. occidentalis** L. Black R. Thimbleberry. Hybrids between this and *R. strigosus* frequently occur. Common. Th.

1163. **R. odoratus** L. Purple Flowering Raspberry. From Ft. Gratiot (Winchell) northward. Not observed in the interior of the C. & S. Common in N. and U. P. Th.

1164. **R. parviflorus** Nutt. Salmon-berry. *R. Nutkanus* Mocino. Salmon-berry. Frequent at Petoskey and common farther north; Elk Rapids, W. S. Cooper. N. & U. P.

1164a. **R. procumbens** Muhl. Low Running Blackberry. *R. Canadensis* T. & G., not L. Keweenaw Co., Detroit, O. A. Farwell.

*1165. **R. setosus** Bigel. Bristly Blackberry. Vestaburg, C. F. Wheeler.

*1166. **R. strigosus** Michx. Wild Red Raspberry. Variable. Common. Th.

DALIBARDA L.

1169. **D. repens** L. Ann Arbor, Mary Clark; Macomb Co., Cooley. May occur farther north, and perhaps throughout. Lake Huron, Bell, Canadian Catalogue.

DRYMOCALLIS Fourr. POTENTILLA, in part.

*1170. **D. arguta** (Pursh.) Rydb. Tall Cinquefoil. *Potentilla arguta* Pursh. Flowers either white or yellow, generally all in one locality of one color. Alma; Ann Arbor; Macomb Co.; Ionia; and N. to Isle Royal. Frequent on light sand; in places common. Th.

DASIPHORA Raf. POTENTILLA, in part.

*1171. **D. fruticosa** (L.) Rydb. Shrubby Cinquefoil. *Potentilla fruticosa* L. Edges of swamps. Ranges N. to Arctic America. Common. Th.

SIBBALDIOPSIS Rydb. POTENTILLA, in part.

1172. **S. tridentata** (Soland.) Rydb. Three-toothed Cinquefoil. *Potentilla tridentata* Soland. Barrens of Missaukee Co.; Grayling; shores of the Upper Great Lakes, A. Gray; Isle Royal, Dr. Lyons; Roscommon, C. A. Davis. N. & U. P. Frequent.

FRAGARIA L.

1173. **F. Americana** (Porter) Britton. American Wood Strawberry. *F. vesca Americana* T. G. Porter. Keweenaw Co., O. A. Farwell; Baldwin, W. J. B.; Vestaburg, C. A. Davis; Alpena, C. F. Wheeler.

1174. **F. Canadensis** Michx. Northern Wild Strawberry. N.

*1175. **F. Virginiana** Duchesne. Virginia Strawberry. *F. Virginiana Illinoensis* Prince. Common. Th.

COMARUM L. POTENTILLA, in part.

*1176. **C. palustre** (L.) Marsh Cinquefoil. *Potentilla plustris* Scop. Swamps throughout, but nowhere abundant.

ARGENTINA Lam. POTENTILLA, in part.

1177. **A. Anserina** (L.) Rydb. *Potentilla Anserina* L. Silver-weed. Frequent. along the Great Lakes, but rare in the interior. Banks of Higgins Lake, G. H. Cannon; Baldwin, W. J. B.; Otsego Co., Guy L. Stewart; Huron Co., C. A. Davis.

POTENTILLA L.

*1178. **P. argentea** L. Silvery Cinquefoil. Ann Arbor, Clark and Allmendinger; Alma, Davis; Inland, Benzie Co.; Behaves like a weed at Hanover, Mich., running out clover, G. E. Simmons; St. Clair Co., C. K. Dodge.

*1179. **P. Canadensis** L. Common Cinquefoil. Five-finger. Common. Th.

1180. **P. intermedia** L. Downy Cinquefoil. Well established in Livingston and Washtenaw counties, C. F. Wheeler.

1181. **P. Monspeliensis** L. Rough Cinquefoil. *P. Norvegica* L. Common. Th.

1182. **P. paradoxa** Nutt. Bushy Cinquefoil. Shores of Great Lakes.

1183. **P. Pennsylvanica** L. L. Superior, Gray's Man. Prof. Ellis, in Canadian Catalogue.

1184. **P. Robbinsiana** Oakes. Robbin's Cinquefoil. *P. frigida* A. Gray. (Not Villar.) Dr. Lyons. Rare. U. P.

*1185. **P. sulphurea** Lam. Much like *P. recta* L. Ypsilanti, O. A. Farwell; Howell Junction, C. F. Wheeler.

WALDSTEINIA Willd.

*1186. **W. fragarioides** (Michx.) Tratt. Barren Strawberry. Livingston Co., Winch. Cat.; Ionia; Flint to L. Sup., Whitney Cat. Rather local, but abundant when found at all. Th.

GEUM L.

*1187. **G. Canadense** Jacq. White Avens. *G. album* Gmelin. Common. L. P.

1188. **G. macrophyllum** Willd. Rare in L. P. Cheboygan Co., B. & K.; N. shore of Lake Superior, Agassiz.

*1189. **G. rivale** L. Water or Purple Avens. Swamps and wet places. Common. Th.

1190. **G. strictum** Aiton. New Haven, Gratiot Co.; Petoskey; Keweenaw Co., O. A. Farwell; St. Clair Co., A. F. Foreste.

1191. **G. vernum** (Raf.) T. & G. Spring Avens. St. Clair, Miss Gurd; Belle Isle, O. A. Farwell. Rare.

*1192. **G. Virginianum** L. Rough Avens. Common. C. & S.

SIEVERSIA R. Br. GEUM, in part.

1193. **S. ciliata** (Pursh.) Rydb. *Geum triflorum* Pursh. Otisco Tp., Ionia Co., A. B. Morse; Montcalm Co.; Ada, Greenville, C. A. Davis.

ULMARIA Hill.

1194. **U rubra** Hill. Queen-of-the-Prairie. *Spiraea lobata* Gronov. "Meadows and prairies, Penn. Co., Mich.," A. Gray. Occurs only in the S. W: R. R. track, near Augusta, L. H. Bailey; Calhoun Co.; University Herb. St. Joseph, V. Willoughby. S.

AGRIMONIA L.

*1195. **A. hirsuta** (Muhl.) Bicknell. Tall Hairy Agrimony. *A. Eupatoria hivsuta* Muhl. Dry soil. Common. Th.

*1196. **A. mollis** (T. & G.) Britton. Soft Agrimony. *A. Eupatoria mollis* T. & G. St. Clair Co., J. W. Stacey; Belle Isle, O. A. Farwell. Frequent.

1197. **A. parviflora** Soland. Many-flowered Agrimony. Detroit, Miss Clark; Macomb Co.; St. Clair Co., A. F. Foerste; S. W., H. S. Pepoon. Infrequent. S. E.

1198. **A. pumila** Muhl. Belle Isle, O. A. Farwell.

1199. **A. striata** Michx. Belle Isle, O. A. Farwell. *A. glabra* (Muhl.) Bicknell.

SANGUISORBA L. POTERIUM L.

1200. **S. Canadensis** L. American Great Burnet. *Poterium Canadense* A. Gray. South Haven, L. H. Bailey; Ann Arbor, Allmendinger. Rare.

ROSA L. Rose.

1201. **R. Arkansana** Porter. Harbor Springs, C. F. Wheeler; Keweenaw Co., O. A. Farwell. Rare.

1202. **R. blanda** Aiton. Ionia; Hubbardston; Flint; Ft. Gratiot, Winch. Cat.; Alma, Ann Arbor, C. A. Davis; Lake shore, New Buffalo, C. F. Wheeler, northward to

Keweenaw Co., O. A. Farwell. Common northward, but infrequent south of lat. 43° except along the shore of Lake Michigan. Th.

1203. **R. canina** L. Mackinac Island, Ypsilanti, O. A. Farwell.

*1205. **R. Carolina** L. Swamps. Common. Th.

*1206. **R. Carolina x humilis** C. F. Wheeler. A supposed hybrid. Border of a swamp near the Agricultural College.

1207. **R. centifolia** L. Along railway track St. Clair Co., C. K. Dodge.

1208. **R. Engelmanni** S. Wats. Prickly Rose. Cheboygan Co.; Mackinaw City: Petoskey, C. F. Wheeler; Keweenaw Co., O. A. Farwell.

*1209. **R. humilis** Marsh. Abundant and pretty. In the C. it is our common wild rose. Dry soil. Alma, Ann Arbor, C. A. Davis. A low form on hills about Ionia, has narrow leaves, with peduncles and ripe fruit glandular-bristly.

*1210. **R. rubiginosa** L. Sweetbrier. Eglantine. Roadsides. Frequent. C. & S.

1211. **R. Sayi** Schwein. Indian River, Black Lake, Cheboygan Co.; Mackinaw City; Petoskey, C. F. Wheeler; Oscoda; Keweenaw Co., O. A. Farwell. N. & U. P.

1212. **R. setigera** Michx. Climbing or Prairie Rose. South Haven, L. H. Bailey; Jackson Co., Winchell; Flint, Dr. Clark; Macomb Co., Cooley; Grosse Isle, Miss Clark; Belle Isle Park, Foerste; near Adrian, Mrs. I. H. Wheeler; along Bean Creek, Addison to Morenci, W. J. B. Indigenous, but rare or local. C. & S.

1213. **R. Woodsii** Lindl., Keweenaw Co., O. A. Farwell.

POMACEÆ L. Apple Family.

SORBUS L.

1214. **S. Aucuparia** L. European Mountain Ash. In a swamp near Bay City, G. M. Bradford.

1215. **S. Americana** Marsh. American Mountain Ash. *Pyrus Americana* DC. Ludington, and north along the Michigan shore to Charlevoix; Sault Ste. Marie; and into Canada where it is common; also westward, through U. P. Burt and Witney; near St. Clair River, C. K. Dodge. N. & U. P.

1216. **S. sambucifolia** (C. & S.) Roem. Western Mountain Ash. *Pyrus sambucifolia* Cham. & Schlect. Mackinac Island, H. Mann; Keweenaw Co., O. A. Farwell; near Houghton Lake, C. A. Davis.

PYRUS L.

1217. **P. communis** L. Common Pear. Ypsilanti, Detroit as an escape, O. A. Farwell.

MALUS Hill. PYRUS, in part.

*1218. **M. coronaria** (L.) Mill. American Crab Apple. *Pyrus cornaria* L. Thickets and along streams. Common. C. & S.

*1219. **M. Malus** (L.) Britton. Apple. Escaped from cultivation. Th.

ARONIA Medic. PYRUS, in part.

1220. **A. arbutifolia** (L.) Medic. Red Choke-berry. *Pyrus arbutifolia* L. f. Hubbardston, New Buffalo, Wheeler; St. Clair Co., C. K. Dodge, and northward.

*1221. **A. nigra** (Willd.) Britton. Black Choke-berry. *Pyrus arbutifolia melanocarpa* Michx. Burt. MS.; Isle Royale, Whitney's Cat. Frequent in swamps. Th.

AMELANCHIER Medic.

1222. **A. alnifolia** Nutt. Presque Isle, Winchell; Keweenaw Co., O. A. Farwell. Rare. N. & U. P.

*1223. **A. Canadensis** (L.) Medic. Shad-bush. Service-berry. Common. Th.

*1224. **A. Botryapium** (L. f.) DC. Shad-bush. *A. Canadensis oblongifolia* T. & G. Ionia; Flint; S. Mich., Winchell Cat.; Alma, Ann Arbor, C. A. Davis. Infrequent. Th.

1225. **A. oligocarpa** (Michx.) Roem. Shores of Lake Superior, Gray's Man.; Keweenaw Co., O. A. Farwell. U. P.

1226. **A. rotundifolia** (Michx.) Roem. Round-leaved June-berry. Grand River Valley, C. F. W.; Elk Rapids, W. S. Cooper. Not common.

1227. **A. spicata** (Lam.) Dec. Low June Berry. Shore of Thunder Bay, near Alpena, Grayling, C. F. Wheeler; Keweenaw Co., O. A. Farwell.

CRATAEGUS L.

1228. C. acutiloba Sargent. Detroit, O. A. Farwell.
1229. C. albicans Ashe. Detroit, O. A. Farwell.
1230. C. altrix Ashe. Detroit, O. A. Farwell.
1231. C. ater Ashe. St. Clair Co., W. W. Ashe.
1232. C. attenuata Ashe. St. Clair Co., W. W. Ashe
1233. C. borealis Ashe. St. Clair Co., W. W. Ashe; Keweenaw Co., O. A. Farwell.
1234. C. brevispina (Dougl.) Farwell. Keweenaw Co., O. A. Farwell. *C. punctata brevispina* Dougl.
1235. C. caesa Ashe. St. Clair Co., W. W. Ashe.
*1236. C. coccinea L. Common. Th.
*1237. C. Crus-galli L. Common. Th.
1238. C. decans Ashe. Detroit, O. A. Farwell.
1239. C. Dodgei Ashe. St. Clair Co., W. W. Ashe.
1240. C. fallax Ashe. St. Clair Co., W. W. Ashe.
1241. C. filipes Ashe. St. Clair Co., W. W. Ashe.
1242. C. gemmosa Sargent. Grand Rapids, Emma J. Cole; St. Clair Co., C. K. Dodge; Detroit, O. A. Farwell.
1243. C. glareola Ashe. St. Clair Co., W. W. Ashe.
1244. C. immanis Ashe. St. Clair Co., W. W. Ashe.
1245. C. latisepala Ashe. St. Clair Co., W. W. Ashe.
1246. C. lanta Ashe. St. Clair Co., W. W. Ashe.
1247. C. lumaria Ashe. St. Clair Co., W. W. Ashe.
*1248. C. macrantha Lodd. Long-spined Thorn. *C. coccinea macrantha* Dudley. Common. Th.
1249. C. Michiganensis Ashe. St. Clair Co., W. W. Ashe.
*1250. C. mollis (T. & G.) Scheele. Red-fruited Thorn. *C. coccinea mollis* T. & G. Frequent in C.; Monroe Co., C. F. Wheeler; Ontonagon, Mary H. Clark; Alma, Ann Arbor, C. A. Davis.
1251. C. nuperia Ashe. Detroit, O. A. Farwell.
1252. C. obtecta Ashe. St. Clair Co., W. W. Ashe.
1253. C. onusta Ashe. St. Clair Co., W. W. Ashe.
1254. C. Oxyacantha L. English Hawthorn. Detroit, O. A. Farwell.
1255. C. pascens Ashe. St. Clair Co., W. W. Ashe.
1256. C. pastora Sargent. Detroit, O. A. Farwell.
1257. C. prona Ashe. St. Clair Co., W. W. Ashe.
1258. C. prunifolia (Marsh.) Pers. Detroit, O. A. Farwell.
1259. C. pubifolia Ashe. St. Clair Co., W. W. Ashe.
1260. C. pubipes Ashe. St. Clair Co., W. W. Ashe.
*1261. C. punctata Jacq. Very variable. Common in south.
1262. C. rotundifolia (Ehrh.) Borck. Glandular Thorn. Grand Rapids, Island Lake, Grass Lake, C. F. Wheeler.
1263. C. redolans Ashe. Detroit, O. A. Farwell.
1264. C. sera Sargent. Detroit, O. A. Farwell.
1265. C. structilis Ashe. Detroit, O. A. Farwell.
1266. C. tenax Ashe. St. Clair Co., W. W. Ashe.
*1268. C. tomentosa L. Pear Thorn. Common. Th.
1269. C. virella Ashe. St. Clair Co., W. W. Ashe.

DRUPACEÆ DC. Plum Family.

PRUNUS L.

*1270. P. Americana Marsh. Wild Yellow or Red Plum. Black River, Cheboygan Co., B. & K.; St. Clair Co., C. K. Dodge; Detroit, O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Th.
*1271. P. Cerasus L. Sour Cherry. Escaped from cultivation by the aid of birds. Keweenaw Co., O. A. Farwell, and in many other places.
1272. P. cuneata Raf. Appalachian Cherry. Bay City and vicinity, G. M. Bradford; Keweenaw Co., O. A. Farwell.
1273. P. domestica L. Well established. St. Clair Co., C. K. Dodge; Ypsilanti, O. A. Farwell.
1274. P. Mahaleb L. Mahaleb Cherry. Well established near St. Clair, C. K. Dodge; Detroit, O. A. Farwell.

*1275. **P. nigra** Ait. Canada Plum. Port Huron northward; Detroit, O .A. Farwell.

*1276. **P. Pennsylvanica** L. f. Wild Red Cherry. Very abundant on sandy land in the N. half of the State, but less common southward, where P. serotina takes it place.

1277. **P. pumila** L. Dwarf Cherry. Sand Cherry. L. Sup.; Emmet Co.; Houghton Lake; Mecosta Co.; South Haven; Saginaw Bay, Winchell, etc.; St. Clair Co., C. K. Dodge. Frequent in the N. half of the L. P., but not yet found in the interior. S. of Saranac, Ionia Co., where it occurs in a dry glade.

*1278. **P. serotina** Ehrh. Wild Black Cherry. Frequent in C. and S. Rare in N. and U. P.

*1279. **P. Virginiana** L. Choke Cherry. A shrub or small tree. Common. Th.

AMYGDALUS L.

1279a. **A. Persica** L. Peach. Well established at Ypsilanti, O. A. Farwell.

CAESALPINACEÆ Kl. & Garcke. Senna Family.

CERCIS L.

*1280. **C. Canadensis** L. Red-bud. Judas-tree. Indigenous throughout the southern part of the State, and as far N. in the west as Grand River Valley. Plaster Creek, Grand Rapids, Garfield; Ionia, Le Valley; banks of Thornapple River. Eaton Co.; Ann Arbor; Adrian; South Haven; near Medina a tree was found 26 inches in diameter! W. J. B.

CASSIA L.

*1281. **C. Marilandica** L. Wild Senna. Grand Haven; Ann Arbor, river banks; Bay Co., G. M. Bradford. Infrequent. C. & S.

1282. **C. nictitans** L. Sturgis, F. P. Daniels; S. W., H. S. Pepoon.

GLEDITSIA L.

1283. **G. triacanthos** L. Three-thorned Acacia. Honey-Locust. Grows along the River Raisin, and is certainly indigenous. Often two feet in diameter, W. J. B. Along the St. Joseph, also, and in other localities in the extreme S. Dundee, Niles, C. F. Wheeler; Detroit, O. A. Farwell; Milan, C. A. Davis.

GYMNOCLADUS Lam.

1284. **G. dioica** (L.) Koch. Kentucky Coffee-tree. C. & S. *G. Canadensis* Lam. A slender tree along the river banks as far N. as Maple River, in Clinton Co.; also Fish Creek, Montcalm Co.; banks of Grand River; Macomb Co., W. J. B.; Belle Isle, O. A. Farwell and W. J. B.; Sturgis, F. P. Daniels; Ann Arbor, C. A. Davis. Infrequent.

PAPILIONACEÆ L. Pea Family.

BAPTISIA Vent.

1285. **B. leucantha** T. & G. Large White Wild Indigo. Calhoun Co., Winch. Cat.; Ann Arbor, Allmendinger Cat.; shore of Lake Erie, Goldie, Canadian Cat.; near Detroit, O. A. Farwell. Rare.

*1286. **B. tinctoria** (L.) R. Br. Wild Indigo. Woodward Lake, Ionia Co.; Flint; Macomb Co.; Ann Arbor, Winch. Cat.; Alma, C. A. Davis. Abundant in places but not well distributed. C. & S.

CROTALARIA L.

1287. **C. sagittalis** L. Rattle-box. Detroit, O. A. Farwell.

LUPINUS L.

*1288. **L. perennis** L. Wild Lupine. C. & S. Abundant in light sand.

1289. **L. perennis occidentalis** S. Wats. Benton Harbor, C. F. Wheeler.

1290. **L. polycarpus** Greene. Ballast grounds. Detroit, O. A. Farwell.

MEDICAGO L.

1291. **M. denticulata** Willd. Toothed Medick. Detroit, O. A. Farwell.
*1292. **M. lupulina** L. Black Medick. Nonesuch. Waste places. Mackinac; Jackson; Ann Arbor; Alma, C. A. Davis. L. P.
*1293. **M. sativa** L. Lucerne. Alfalfa. Escaped from cultivation in a few places.

MELILOTUS Juss.

*1294. **M. alba** Desv. White Melilot. Very common in the older portions of the State where it has often been sown by the roadside to furnish "pasture" for bees. L. P.
*1295. **M. officinalis** (L.) Lam. Yellow Melilot. It needs heavy soil. Rare. C. & S.

TRIFOLIUM L.

*1296. **T. arvense** L. Rabbit-foot or Stone Clover. Becoming naturalized in light soil.
1297. **T. aureum** Poll. Yellow or Hop Clover. *T. agrarium* L., in part. Port Huron, C. K. Dodge; Keweenaw Co., O. A. Farwell. Not common. Th.
1298. **T. depauperatum** Desv. Introduced at Detroit, O. A. Farwell.
1299. **T. dubium** Sibth. Grand Rapids, Miss E. J. Cole; Detroit and Ypsilanti, O. A. Farwell.
1300. **T. furcatum** Lindl. Introduced at Detroit, O A Farwell
*1301. **T. hybridum** L. Alsike Clover. Escaped from cultivation.
1302. **T. medium** L. Occasional in roads and fields. Port Huron, C. K. Dodge.
*1303. **T. pratense** L. Red Clover. Meadows and fields. Th.
*1304. **T. procumbens** L. Low Hop-Clover. Infrequent. C. & S.
*1305. **T. repens** L. White Clover. Meadows, pastures and roadsides. Very common. Th.
1306. **T. stoloniferum** Muhl. Running Buffalo Clover. Kalamazoo, Tuthill. Introduced (?)

AMORPHA L.

1307. **A. canescens** Pursh. Lead-plant. W. Mich., Winch. Cat.; Ann Arbor, Prof. M. W. Harrington; Kalamazoo; Tuthill; Klinger Lake; Barron Lake, C. F. Wheeler; Sturgis, F P. Daniels; Berrien Co., H. S. Pepoon. Rare.

KUHNISTERA Lam. PETALOSTEMON Michx.

1308. **K. purpurea** (Vent.) MacM. Violet Prairie-clover. *Petalostemon violaceus* Michx. Berrien Co., H. S. Pepoon.

CRACCA L. TEPHROSIA Pers.

1309. **C. Virginiana** L. Goat's Rue. Catgut. *Tephrosia Virginiana* Pers. Clinton Co.; Montcalm Co.; Newaygo Co.; Macomb Co.; Livingston Co., Miss Clark. On light sand. Infrequent. C. & S.

ROBINIA L.

*1310. **R. Pseudacacia** L. Common Locust or False Acacia. Escaped from cultivation.
1311. **R. viscosa** Vent. Clammy Locust. Escaped from cultivation at Ionia, C. F. Wheeler.

ASTRAGALUS L.

*1312. **A. Carolinianus** L. Carolina Milk Vetch. *A. Canadensis* L. Put-in-Bay; Ann Arbor; Macomb Co.; Orchard Lake; Montcalm Co.; river bank, North Lansing; Shiawassee Co., G. H. Hicks; Whitmore Lake, C. A. Davis; and northward. Well distributed, but infrequent. Th.

PHACA L. Some authors.

*1313. **P. neglecta** T. & G. Cooper's Milk Vetch. *Astragalus Cooperi* A. Gray. Dr. A. B. Lyons; Cooley MS.; Park Lake, Clinton Co., L. H. Bailey; Alma. Rare. C. & S.

STROPHOSTYLES Ell.

1314. **S. helvola** (L.) Britton. Trailing Wild Bean. *S. angulosa* Ell. S. E. along shore, and on the islands of Lake Erie, and shore of Lake Michigan at South Haven, L. H. Bailey; Belle Isle, O. A. Farwell. Probably not much farther north, and not found in the interior.

MEIBOMIA Heist. DESMODIUM Desv.

*1315. **M. bracteosa** (Michx.) Kuntze. Large-bracted Tick-trefoil. *Desmodium cuspidatum* Hook. Hubbardston; Flint; Ann Arbor; Macomb Co.; S. Mich., Wright; Alma. Usually on oak land. Frequent. C. & S.

*1316. **M. Canadensis** (L.) Kuntze. Showy Tick-trefoil. *Desmodium Canadense* DC. Hubbardston, and probably father N.; Flint; Ann Arbor, Allmendinger Cat. and Winch. Cat. Flowers early. Frequent. Th.

1317. **M. canescens** (L.) Kuntze. Hoary Tick-trefoil. *Desmodium canescens* DC. Ann Arbor, Allmend. Cat.; S. West, Wright, Cat. Rare. S.

*1318. **M. Dillenii** (Darl.) Kuntze. Dillen's Tick-trefoil. *Desmodium Dillenii* Darlington. Oak woods. Hubbardston; Flint; Macomb Co.; S. Mich., Wright Cat.; Mont Lake, Miss Clark. Frequent. C. & S.

*1319. **M. grandiflora** (Walt.) Kuntze. Pointed-leaved Tick-trefoil. *Desmodium acuminatum* DC. Woodlands. Very common. C. & S.

1320. **M. Illinoensis** (A. Gray) Kuntze. Illinois Tick-trefoil. *Desmodium Illinoense* A. Gray. University campus, Ann Arbor; White Pigeon, W. J. B.; S. W., H. S. Pepoon.

*1321. **M. Marilandica** (L.) Kuntze. Small-leaved Tick-trefoil. *Desmodium Marilandicum* F. Boott. Dry hills; Ionia, and southward; near Park Lake, Clinton Co., W. J. B.; Orion, O. A. Farwell. Infrequent. C. & S.

*1322. **M. Michauxii** Vail. Prostrate Tick-trefoil. *Desmodium rotundifolium* DC. Hubbardston; Ionia; Flint; South Haven; Macomb Co.; Ann Arbor, Allmend. Cat.; Orion, O. A. Farwell. Infrequent. C. & S.

*1323. **M. nudiflora** (L.) Kuntze. Naked-flowered Tick-trefoil. *Desmodium nudiflorum* DC. Crystal Lake, Montcalm Co.; Flint; Macomb Co.; S. Mich., Wright. Cat. Common. C. & S.

1324. **M. obtusa** (Muhl.) Vail. Hairy Small-leaved Tick-trefoil. *Desmodium ciliare* DC. Dr. Wright. Rare. S.

*1325. **M. paniculata** (L.) Kuntze. Panicled Tick-trefoil. *Desmodium paniculatum* DC. Oak woods. Hubbardston; Flint; Macomb Co.; Alma; South Haven; S. Mich., Wright's Cat. Frequent. C. & S.

*1326. **M pauciflora** (Nutt.) Kuntze. Few-flowered Tick-trefoil. *Desmodium pauciflorum* DC. Mont Lake, Miss Clark in Winch. Cat. Rare. S.

1327. **M. rigida** Ell. Kuntze. Rigid Tick-trefoil. *Desmodium rigidum* DC. Hubbardston; Flint; Macomb Co.; Ann Arbor and S. W., Winch. Cat. Infrequent. C. & S.

1328. **M. sessilifolia** (Torr.) Kuntze. Sessile-leaved Tick-trefoil. *Desmodium sessilifolium* T. & G. Michigan, A. Gray; S. Mich., Winch. Cat.; University Herb., Niles, Ames; Greenville; Rochester, O. A. Farwell. Rare. S.

LESPEDEZA Michx.

1329. **L. augustifolia** (Pursh.) Ell. Narrow-leaved Bush-clover. Sandy fields. Infrequent. C. & S.

*1330. **L. capitata** Michx. Round-headed Bush-clover. Abundant in old fields; prefers light sand. Common. C. & S.

*1331. **L. frutescens** (L.) Britton. Wand-like Bush-clover. *L. Stuvei intermedia* S. Watts. *L. reticulata* S. Wats. Gray's Manual; Britton and Brown; Orion and Detroit, O. A. Farwell; S. W., H. S. Pepoon. S.

*1332. **L. hirta** (L.) Ell. Hairy Bush-clover. *L. polystachya* Michx. Common. C. & S.

1333. **L. Nuttallii** Darl. Nuttall's Bush-clover. Britton and Brown.

1334. **L. procumbens** Michx. Trailing Bush-clover. S. W., H. S. Pepoon.

1335. **L. Stuvei** Nutt. Stuve's Bush-clover. Winchell Cat.; Gray in Manual. Rare. S.

*1336. **L. violacea** (L.) Pers. Bush-clover. Ann Arbor, Miss Clark; S. W., Wright; Rochester, O. A. Farwell; Vestaburg, C. A. Davis. Infrequent.

1337. **L. Virginica** (L.) Britton. Slender Bush-clover Munith, G. H. Hicks; Norvell, C. F. Wheeler; Orion, O. A. Farwell.

VICIA L.

*1338. **V. Americana** Muhl. Ann Arbor; Ionia; Montcalm Co.; N. to Lake Superior; Whitmore Lake, C. A. Davis. Spreads rapidly in C., along railroads and highways and acts like an immigrant. Th.

1339. **V. angustifolia** Roth. Smaller Vetch. *V. sativa angustifolia* Ser. Keweenaw Co., O. A. Farwell.

*1340. **V Caroliniana** Walter. Dry soil. Common. C. & S.

1341. **V. Cracca** L. Cow Vetch. S. Mich., Wright Cat.; Ann Arbor, Prof. M. W. Harrington; Alma, C. A. Davis. Rare or local.

1342. **V. sativa** L. Common Vetch. Tare. Detroit, Dr. Lyons; Springwells, Henry Gillman; Port Huron, C. K. Dodge; Belle Isle, O. A. Farwell; Alma. Infrequent.

LATHYRUS L.

1343. **L. maritimus** (L.) Bigel. Beach Pea. All around the Great Lakes, but seldom seen in the interior. Shore of Higgins Lake, G. H. Cannon. Th.

1344. **L. myrtifolius** Muhl. Myrtle-leaved Marsh Pea. *L. palustris myrtifolius* A. Gray. Infrequent. Th.

*1345. **L. ochroleucus** Hooker. Hillsides and dry uplands. Keweenaw Co., O. A. Farwell; Alma, and southward. Infrequent. Th.

*1346. **L. palustris** L. Ann Arbor; Ionia; Bay City, Antrim Co.; Alma; Petoskey to L. Sup. Th.

1347. **L. venosus** Muhl. Ionia Co.; Clinton Co.; Flint; Macomb Co.; Ann Arbor; Vestaburg. Northward to Keweenaw Co., O. A. Farwell. Infrequent. Th.

FALCATA Gmel. AMPHICARPAEA Ell.

*1348. **F. comosa** (L.) Kuntze. Hog Pea-nut. *Amphicarpaea monoica* Ell. Fields and woods. Common. Th.

*1349. **F. Pitcheri** (T. & G.) Kuntze. Pitcher's Hog Pea-nut. *Amphicarpaea Pitcheri* T. & G. Moist woodlands. Hubbardston, C. F. Wheeler; Lenawee Co., W. J. B.; Manistee, F. P. Daniels; Alma, Whitmore Lake, C. A. Davis.

APIOS Moench.

*1350. **Apios Apios** (L.) MacM. Ground-nut. *A. tuberosa* Moench. Common. C. & S.

GERANIACEÆ J. St. Hil. Geranium Family.

GERANIUM L.

1351. **G. Bicknellii** Britton. Bicknell's Crane's-bill. Thunder Bay Island, Alpena Co., C. F. Wheeler; Keweenaw Co., O. A. Farwell.

*1352. **G. Carolinianum** L. Carolina Crane's-bill. Macomb Co.; Clinton Co.; Alma. Rare in S. & C., abundant in vicinity of Farwell, Clare Co., thence N. to L. Superior. Th.

1353. **G. columbinum** L. Long-stalked Crane's-bill. Grand Detour, T. C. Porter.

*1354. **G. maculatum** L. Wild Crane's-bill. Canada to Florida. T. & G. Common. C. & S.

1355. **G. molle** L. Harbor Springs, Hon. G. L. Maurice.

*1356. **G. pusillum** Burm. f. Small-flowered Crane's-bill. Flint, Dr. Clark; Benzie Co., W. J. B.; Constantine, C. F. Wheeler; St. Clair Co., C. K. Dodge; Ann Arbor, C. A. Davis. Adventive.

1357. **G. Robertianum** L. Herb Robert. Put-in-Bay, Lake Erie; Montcalm Co.; Saginaw Bay; Mackinac, and Drummond's Is., Winch. Cat.; L. Sup. Whitney's Cat.; Keweenaw Co., O. A. Farwell; Island near Algonac, C. K. Dodge. Common around the Great Lakes, but seldom seen in the interior.

1358. **G rotundifolium** L. Round-leaved Crane's-bill. Britton & Brown.

ERODIUM L'Her.

1359. **E. cicutarium** (L.) L'Her. Stork's-bill. Alfilarilla. Oceana Co.; Ionia Co.; Kalamazoo; Port Huron, C. K. Dodge. Not yet common.

OXALIDACEÆ Lindl. Wood-Sorrel Family.

OXALIS L.

1360. **O. Acetosella** L. Common Wood-sorrel. Macomb Co.; L. Sup. and northward, A. Gray. .Rare in S. Peninsula. Th.
1361. **O. corniculata** L. Rochester, O. A. Farwell.
1362. **O. cymosa** Small. Tall Yellow Wood-sorrel. Marquette Co., Burt's MS., Rochester, O. A. Farwell. Southward. Common.
*1363. **O. stricta** L. Upright Yellow Wood-sorrel. Marquette southward. Common.
1364. **O. violacea** L. Violet Wood-sorrel. Winchell Cat.; S. W., H. S. Pepoon. Rare. S. E.

LINACEÆ Dumort. Flax Family.

LINUM L.

1365. **L. humule** Mill. Belle Isle, rare, O. A. Farwell.
1366. **L. medium** (Planch.) Britton. Port Huron, C. K. Dodge; Belle Isle, O. A. Farwell.
1367. **L. striatum** Walt. Ridged Yellow Flax. S. W., H. S. Pepoon.
1368. **L. sulcatum** Riddell. Grooved Yellow Flax. Dr. Wright; University Herb. Rare. S.
1369. **L. usitatissimum** L. Common Flax. Dr. Clark, and Dr. Wright; Keweenaw Co., O. A. Farwell; Alma, C. A. Davis. Escaped from cultivation. Infrequent.
*1370. **L. Virginianum** L. Wild Yellow Flax. Ann Arbor, Dr. Lyons; S. Mich., Wright Cat.; Lenawee Co., G. F. Comstock. Infrequent. S.

RUTACEÆ Juss. Rue Family.

XANTHOXYLUM L.

*1371. **X. Americanum** Mill. Northern Prickly Ash. Toothache-tree. Everywhere along streams and on low ground. L. P.

PTELEA L.

*1372. **P. trifoliata** L. Hop-tree Shrubby Trefoil. Very abundant on the low, rocky islands in the western end of Lake Erie, and frequent along river banks and the shore of the Great Lakes. Occurs in the interior as far north as Montcalm Co. Does not occur at Flint, and is not mentioned in Dr. Cooley's MS. Cat., but is given in Dr. Wright. Alma, Ann Arbor, C. A. Davis. C. & S.

SIMARUBACEÆ DC. Ailanthus Family.

AILANTHUS Desf.

1373. **A. glandulosa** Desf. Tree-of-Heaven. Port Huron, C. K. Dodge; Detroit, O. A. Farwell.

POLYGALACEÆ Reichenb. Milkwort Family.

POLYGALA L.

1374. **P. ambigua** Nutt. Loose-spiked Milkwort. *P. verticillata ambigua* Wood. Ypsilanti, Mrs. Lucy Osband; Port Huron, C. K. Dodge.
1375. **P. cruciata** L. S. Mich., Wright Cat. Rare. S.
1376. **P. incarnata** L. Pink Milkwort. Port Huron, Walpole Island, C. K. Dodge. Rare.
*1377. **P. paucifolia** Willd. Flowering Wintergreen. Fringed Polygala. Common on pine land. Th.

*1378. **P. polygama** Walt. Sandy soil. Ionia, Clinton and Montcalm Counties; Grass Lake, Vestaburg, C. A. Davis. Southward. Abundant near Quinnesec, Menominee Co., E. J. Hill. Th.

*1379. **P. Senega** L. Seneca Snakeroot. Alma, C. A. Davis; Bay Co., G. M. Bradford; near Detroit, O. A. Farwell. Th.

1380. **P. Senega latifolia** Torr. & Gray. Hubbardston; Flint; Ann Arbor, C. A. Davis. Rare. Th.

*1381. **P. verticillata** L. Whorled Milkwort. Flint; Macomb Co.; Wayne Co. Common in some places, rare in others.

*1382. **P. viridescens** L. Purple Milkwort. *P. sanguinea* L. Belding; Ionia; Flint; Macomb Co.; Grass Lake, Whitmore Lake, C. A. Davis. Local, but usually abundant when found at all. C. & S.

EUPHORBIACEÆ J. St. Hil. Spurge Family.

ACALYPHA L.

1383. **A. gracilens** A. Gray. Slender Three-seeded Mercury. *A. Virginica gracilens* Muell. Detroit, Dr. D. Clark.

*1384. **A. Virginica** L. Three-seeded Mercury. Open woods. Variable. Common. Th.

EUPHORBIA L.

1385. **E. commutata** Engelm. Tinted Spurge. Ann Arbor, C. A. Davis; Flint. Infrequent.

*1386. **E. corollata** L. Flowering Spurge. Sandy soil. Frequent. C. & S.

*1387. **E. Cyparissias** L. Cypress Spurge. Escaped from cultivation. Frequent.

*1388. **E. Esula** L. Leafy Spurge. Escaped from cultivation.

1389. **E. glyptosperma** Engelm. Ridge-seeded Spurge. Escanaba, Marquette, C. F. Wheeler.

1390. **E. Helioscopia** L. Sun Spurge. Common at Detroit, Dr. A. B. Lyons; St. Clair Co., C. K. Dodge; Mackinac Island, O. A. Farwell.

1391. **E. hirsuta** (Torr.) Weigand. Hairy Spurge. *E. hypericifolia hirsuta* Torr. Clair Co., C. K. Dodge; Mackinac Island, O. A. Farwell.

1392. **E. humistrata** Engelm. Hairy Spreading Spurge. Flint, Dr. D. Clark; St. Clair Co., C. K. Dodge; Bay Co., G. M. Bradford.

*1393. **E. maculata** L. Spotted Spurge. Roadsides and fields, everywhere.

1394. **E. Nicaeensis** All. Nicaean Spurge. Port Huron, C. K. Dodge

*1395. **E. nutans** Lag. Upright Spotted Spurge. *E. Preslii* Guss. Cultivated soil, and waste places. Detroit, Dr. A. B. Lyons; Ionia; Grand Rapids. Infrequent. C. & S.

*1396. **E. obtusata** Pursh. Belle Isle, rare, O. A. Farwell.

*1397. **E. Peplus** L. Petty Spurge. Grand Rapids, H. C. Skeels; Olivet, H. L. Clark.

1398. **E. platyphylla** L. Broad-leaved Spurge. Macomb Co.; "along the Great Lakes," A. Gray; Lake Huron, Dr. Todd.

1399. **E. polygonifolia** L. Knotweed Spurge. Sandy shores of the Great Lakes. South Haven, L. H. Bailey; Fort Gratiot, Winch. Cat.; Roscommon and Ann Arbor, C. A. Davis. Frequent. Th.

1400. **E. serpyllifolia** Pers. Thyme-leaved Spurge. Keweenaw Co., O. A. Farwell.

CALLITRICHACEÆ Lindl. Water-Starwort Family.

CALLITRICHE L.

1401. **C. bifida** (L.) Morong. Northern Water Starwort. *C. autumnalis* L. Flint, Dr. Clark; L. Superior, A. Gray; Alma, C. A. Davis; Sault de Ste. Marie, T. Morong.

1402. **C. palustris** L. Water Starwort. Water Fennel. *C. verna* L. Ponds, Macomb Co., Cooley; Keweenaw Co., O. A. Farwell; Norway, E. J. Hill; Alma, C. A. Davis.

EMPETRACEÆ Dumort. Crowberry Family.

EMPETRUM L.

1403. **E. nigrum** L. Black Crowberry. Whitney Cat. Pictured Rocks, G. H. Hicks. U. P.

LIMMANTHACEÆ Lindl. False Mermaid Family.

FLOERKEA Willd.

*1404. **F. proserpinacoides** Willd. False Mermaid. Ionia; Hubbardston; Flint; Ann Arbor, C. A. Davis. Not rare but usually overlooked.

ANACARDIACEÆ Lindl. Sumac Family.

RHUS L.

1405. **R. aromatica** Ait. Fragrant Sumach. *R. Canadensis* Marsh. Less frequent than other members of the genus. Found chiefly on bluffs and sandy hills through the middle counties of the L. P. "N. to the Saskatchawan," Torr. & Gr.; Saginaw Co., Washtenaw Co., C. A. Davis. Th.

*1406. **R. copallina** L. Dwarf Sumach. Abundant in the pine country on light soil. C. & S.

*1407. **R. glabra** L. Smooth Sumach. Alma, Ann Arbor, C. A. Davis. Very common in C. Th.

1407a. **R. glabra borealis** Britton. Keweenaw Co., O. A. Farwell.

*1408. **R. hirta** (L.) Sudw. Staghorn Sumach. *R. typhina* L. Alma, Ann Arbor, C. A. Davis.

1409. **R. pubescens** (Mill) O. A. F. Northern Poison Oak. *R. Toxicodendron quercifolia* Michx. Belle Isle, O. A. Farwell.

*1410. **R. radicans** L. Poison, climbing or Three-leaved Ivy. Poison Oak. L. P.

1411. **R. Rydbergii** Small. Keweenaw Co., O. A. Farwell.

*1412. **R Toxicodendron** L. Tall-climbing, often to tops of forest trees; stems occasionally several inches in diameter. Th.

*1413. **R. Vernix** L. Poison Sumach. Poison Elder. *R. venenata* DC. Common in swamps. Green Lake, G. Traverse Co. and southward. L. P.

ILICACEÆ Lowe. Holly Family.

ILEX L.

*1414. **I. verticillata** (L.) A. Gray. Black Alder. Winterberry. Low grounds. Th.

ILICIOIDES Dumont. NEMOPANTHES Raf.

*1415. **I. mucronata** (L.) Britton. Mountain Holly. *Nemopanthes fascicularis* Raf. Borders of swamps. S. Mich., Wright Cat.; Ionia Co.; Alma, Ann Arbor, C. A. Davis; Montcalm Co.; Flint; Macomb Co., and northward. Frequent. Th.

CELASTRACEÆ Lindl. Staff-tree Family.

EUONYMUS L.

*1416. **E. atropurpureus** Jacq. Wahoo. Burning Bush. Low river banks. Well distributed, but nowhere very common. C. & S.

*1417. **E. obovatus** Nutt. Strawberry Bush. *E. Americanus obobatus* T. & G. Trailing. Common. C. & S.

CELASTRUS L.

*1418. **C. scandens** L. Wax-work. Climbing Bittersweet. Frequent. Th.

STAPHYLEACEÆ DC. Bladder-nut Family.

STAPHYLEA L.

*1419. **S. trifolia** L. American Bladder-nut. Marquette Co., and southward, Burt. Frequent along river banks in L. P. Th.

ACERACEÆ J. St. Hil. Maple Family.

ACER L.

*1420. **A. Negundo** L. Box Elder. Ash-leaved Maple. Th. *Negundo aceroides* Moench. River banks. A small tree. West of Lake Superior, and north to Saskatchewan Valley. Macoun in Candian Cat. Frequent in Grand River Valley; escaped from cultivation in St. Clair Co., C. K. Dodge; Belle Isle, Alma, Ann Arbor, C. A. Davis. Th.

*1420a. **A. nigrum** Michx. Black Sugar Maple. *A. saccharinum nigrum* T. & G. Common. C. & S.

1421. **A. Pennsylvanicum** L. Striped Maple. Abundant in U. P. Whitney; common at Petoskey, and occasional as far S. on the Huron shore as Alcona Co., Winchell Cat.; in the interior as far as Houghton Lake. N. & U. P.

1422. **A. platanoides** L. Norway Maple. Banks of Huron River, Ypsilanti, O. A. Farwell. Introduced from Europe.

*1424. **A. rubrum** L. Red or Swamp Maple. On low ground and along streams. Very common. Th.

*1425. **A. saccharinum** L. Silver Maple. *A. dasycarpum* Ehrh. Low ground along rivers. Common. C. & S.

*1426. **A. Saccharum** Marsh. Rock or Sugar Maple. *A. saccharinum* Wang. Frequent in the U. P., and abundant in the lower, forming extensive groves, either alone or in connection with beech. Th.

*1427. **A. spictatum** Lam. Mountain Maple. Common in U. P.; Alcona Co., Winch. Cat.; Crystal Lake, Montcalm Co.; Alma, C. N. & U. P.

HIPPOCASTANACEÆ T. & G. Buckeye Family.

AESCULUS L.

1428. **AE. glabra** Willd. Ohio Buckeye. Fetid B. Lenawee Co., W. J. B.; Washtenaw Co., O. A. Farwell, and probably in other localities in the southern tier of counties. S

1429. **AE. Hippocastenum** L. Horse-chestnut. Escaped from cultivation. St. Clair Co., C. K. Dodge; Washtenaw and Wayne counties, O. A. Farwell.

BALSAMINACEÆ Lindl. Jewel-weed Family.

IMPATIENS L.

*1430. **I. aurea** Muhl. Pale Touch-me-not. *I. pallida* Nutt. Macomb Co.; Ionia Co.; Sugar Is., Winch. Cat.; Ann Arbor, C. A. Davis. Much rarer than the next. Th.

*1431. **I. biflora** Walt. Spotted Touch-me-not. *I. fulva* Nutt. Keweenaw Co., O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Southward.

RHAMNACEÆ Dumort. Buckthorn Family.

RHAMNUS L.

*1432. **R. alnifolia** L'Her. Alma, Ann Arbor, C. A. Davis. Common. Th.

*1433. **R. cathartica** L. Buckthorn. Escaped from cultivation. Lansing, and very likely in other portions of the state.

CEANOTHUS L.

*1434. **C. Americanus** L. New Jersey Tea. Red-root. Sandy woods, rarely on beech and maple land. Alma, Ann Arbor, C. A. Davis. Frequent.

1435. **C. ovatus** Desf. Smaller Red-root. East shore L. Huron, E. Ont. and L. Sup., Can. Cat.; Cheboygan Co., B. & K.; Alcona Co.; Grayling; Oscoda; Manistee, F. P. Daniels.

VITACEÆ Lindl. Grape Family.

VITIS L.

1436. **V. aestivalis** Michx. Summer Grape. Sturgis, F. P. Daniels.

*1437. **V. bicolor** Le Conte. Winter Grape. Lyons; Stanton; Crystal Lake; Montcalm Co.; Flint; Ann Arbor, Winch. Cat.; Bay Co., G. M. Bradford; S. Mich., Wright Cat. Infrequent. C. & S.

1438. **V. cordifolia** Michx. Frost Grape. Sturgis, F. P. Daniels.

*1439. **V. Labrusca** L. Northern Fox-grape. Muskegon, W. J. B.

1440. **V. rotundifolia** Michx. Rochester, O. A. Farwell.

*1441. **V. vulpina** L. Riverside Grape. *riparia* Michx. Sturgis, F. P. Daniels. Common along our rivers; infrequent in the pine region, where V. bicolor is very abundant. Th.

PARTHENOCISSUS Planch.

*1442. **P. quinquefolia** (L.) Planch. Virginia Creeper. American Ivy. *Ampelopsis quinquefolia* Michx. Common. Th.

1443. **P. quinquefolia laciniata** (Planch.) *R. vitacea* A. S. Hitchcock. Frequent.

1444. **P. quinquefolia hirsuta** (Donn.) Torr. & Gray. Bald-head Park, Saugatuck, C. F. Wheeler; Belle Isle, O. A. Farwell.

TILIACEÆ Juss. Linden Family.

TILIA L.

*1445. **T. Americana** L. Basswood. Linden. Abundant in C. & S., common in Emmet Co., and frequent in U. P., "especially in Ontonagon Valley," Whitney Cat. Th.

MALVACEÆ Neck. Mallow Family.

ALTHAEA L.

1446. **A. officinalis** L. Marsh-Mallow. Bay Co., G. M. Bradford; Detroit, O. A. Farwell.

MALVA L.

*1447. **M. Alcea** L. European Mallow. Adventive at the Agricultural College; Manistee, F. P. Daniels.

*1447a. **M. crispa** L. Curled Mallow. *M. verticillata crispa* L. Flint, Dr. Clark; Detroit, O. A. Farwell.

*1448. **M. moschata** L. Musk Mallow. Keweenaw Co., O. A. Farwell; Alma, C. A. Davis. Southward. Not rare.

*1449. **M. rotundifolia** L. Common Mallow. Keweenaw Co., O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Southward. Common. Th.

*1450. **M. sylvestris** L. High Mallow. Occasional. L. P.

SIDA L.

1452. **S. hermaphrodita** (L.) Rusby. Virginia Mallow. *S. Napaea* Cav. Kalamazoo, R. R. track, Tuthill. Rare. S.

1453. **S. spinosa** L. Prickly Sida. Britton and Brown.

ABUTILON Gaertn.

*1454. **A. Abutilon** (L.) Rusby. Velvet-leaf. *A. Avicennae* Gaert. Roadsides and river banks. Very common in places, and, along with Datura Stramonium, forming regular thickets. Alma, Ann Arbor, C. A. Davis. C. & S.

HIBISCUS L.

1455. **H. Moscheutos** L. Swamp Rose-mallow. Monroe; Flint; Put-in-Bay, Lake Erie; St. Joseph, V. Willoughby; Detroit, D. H. Campbell; Saugatuck, Tuthill. Rare. C. & S.

*1456. **H. Trionum** L. Bladder Ketmia. South, Dr. Wright; Ann Arbor, Winch. Cat.; Flint; Ypsilanti, O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Rare.

HYPERICACEÆ Lindl. St. John's-wort Family.

HYPERICUM L.

***1457. H. Ascyron** L. Great St. John's Wort. South Haven; Ann Arbor; Ft. Gratiot; Macomb Co.; Ionia; Stanton; Hubbardston; Alma; Crystal Lake; Flint; Ontonagon River, Whitney Cat.; Grand Rapids, Sones; Keweenaw Co., O. A. Farwell. Occurs at wide intervals and in small patches. Th.

1458. **H. boreale** (Britton) Bicknell. *H. Canadense boreale* Britton. Keweenaw Co., O. A. Farwell.

*1459. **H. Canadense** L. Canadian St. John's-wort. Ann Arbor, Miss Clark; L. Sup., A. Gray; Keweenaw Co., O. A. Farwell; Carp River, Porter; Alma. Infrequent.

1460. **H. ellipticum** Hook. Ann Arbor, Miss Clark; Keweenaw Co., O. A. Farwell; Carp River, T. C. Porter. Infrequent.

1461. **H. Kalmianum** L. Kalm's St. John's-wort. Along the lakes; more common northward. Manistee, F. P. Daniels; Fort Gratiot, Winch. Cat.; Petoskey; Port Austin, C. A. Davis; S. W., H. S. Pepoon. Th.

*1462. **H. maculatum** Walt. Spotted St. John's-wort. Common. Th.

*1463. **H. majus** (A. Gray) Britton. Larger Canadian St. John's-wort. *H. Canadense majus* A. Gray. St. Clair Co., Foerste; Flint; Fruitport and Old Mission, E. J. Hill; L. Sup., A. Gray, Vestaburg. Th.

*1464. **H. mutilum** L. Dwarf St. John's-wort. Common. Th.

*1465. **H. perforatum** L. Common St. John's-wort. Ionia; Flint; Keweenaw Co., O. A. Farwell. Infrequent.

*1466. **H. prolificum** L. Shrubby St. J. A low, compact shrub. Frequent from C. Southward. Th.

SAROTHRA L.

1467. **S. gentianoides** L. Pine-weed. *Hypericum nudicaule* Walt. Ypsilanti. O. A. Farwell.

TRIADENUM Raf. ELODEA Pursh, not Michx.

*1468. **T. Virginicum** (L.) Raf. Marsh St. John's-wort. *Elodea campanulata* Pursh. Common. Th.

CISTACEÆ Lindl. Rock-rose Family.

HELIANTHEMUM Pers.

*1469. **H. Canadense** (L.) Michx. Frost-weed. Frequent. Th.

1470. **H. majus** (L.) B. S. P. Hoary Frost-weed. Vestaburg, Davis and Wheeler; Muskegon, W. J. B.; Fife Lake, Alpena, C. F. Wheeler; Detroit, O. A. Farwell.

HUDSONIA L.

1471. **H. tomentosa** Nutt. False Heather. Shores of Great Lakes, A. Gray; S. Mich., Winch. Cat.; L. Sup., Can. Cat.; Oscoda. Frequent. Th.

LECHEA L.

1472. **L. intermedia** Leggett. Large-podded Pin-weed. Port Huron, C. K. Dodge; Bay Co., G. M. Bradford.

1473. **L. minor** L. Thyme-leaved Pin-weed. Pointe aux Pins, Lake Superior, Dawson; Au Sable, W. J. B.; St. Clair Co., C. K. Dodge.

1474. **L. stricta** Leggett. Prairie Pin-weed. Lower falls of the Menominee River, C. F. Wheeler.

*1475. **L. villosa** Ell. Hairy Pin-weed. *L. major* Michx. Common on poor soil. C. & S.

VIOLACEÆ DC. Violet Family.

VIOLA L.

*1477. **V. arenaria** D C. Sand Violet. Sandy land. Crawford Co., W. J. B.

*1478. **V. blanda** Willd. Sweet White Violet. Low ground. Common and variable. Th.

*1479. **V. Canadensis** L. Canada Violet. Abundant on beech and maple land, but seldom, or never found under oaks. Th.

*1479a. **V. cuculata** Ait. Marsh Blue Violet. *V. affinis* Le Conte. *V. obliqua* Hill. Very common. Th.

1479a. **V. crassula** Greene. Near Jackson. E. L. Greene.

1480. **V. cuspidata** Greene. Utica in Macomb Co., W. S. Cooper.

1481. **V. emarginata** (Nutt.) Le Conte. Triangle-leaved Violet. Munith, G. H. Hicks; Muskegon, W. J. B.; Detroit, O. A. Farwell.

1482. **V. fimbriatula** J. E. Smith. *V. ovata* Nutt. S. W., H. S. Pepoon. Very rare. Ann Arbor, Grand Rapids, C. F. Wheeler; Bay City, G. M. Bradford; Detroit, O. A. Farwell.

*1483. **V. Labradorica** Schrank. American Dog Violet. *V. canina Muhlenbergii* Traut. Common. Th.

1484. **V. lanceolata** L. Lance-leaved Violet. W. Harrisville, Glade Pt.; Kalkaska Co.; Roscommon Co.; L. Sup.; Belle Isle, O. A. Farwell; Vestaburg, C. A. Davis.

1484a. **V. Le Conteana** Don. *V. blanda amoena* (Le Conte) B. S. P. Towar's swamp. near Lansing; Cheboygan Co., B. & K.; shores of Barron Lake, Cass Co.; Keweenaw Co., O. A. Farwell.

1484a. **V. nodosa** Green. Marengo, Mich. E. L. Greene.

*1485. **V. palmata** L. Early Blue Violet. Dry to moist woods. Ann Arbor, C. A. Davis.

1485a. **V. papilionacea** Ph. Hooded Blue Violet. Very common.

*1486. **V. pedata** L. Bird-foot Violet. Common on light sand; our finest species N. to British Am.; lat. 55°. Torr. & Gr., Fl. N. Am. Th.

1486a. **V. peramoena** Greene. Marengo. E. L. Greene.

*1487. **V. pubescens** Aiton. Downy Yellow Violet. Beech woods. Common. Th.

*1488. **V. pubescens eriocarpa** (Schw.) Nutt. Detroit, O. A. Farwell.

*1489. **V. renifolia** A. Gray. Kidney-leaved Violet. *V. blanda renifolia* A. Gray. Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell, Gratiot Co., C. A. Davis.

*1490. **V. rostrata** Pursh. Long-spurred Violet. Well distributed and frequent. Alma, Ann Arbor, C. A. Davis. C. & S.

1491. **V. rotundifolia** Michx. Round-leaved Violet. Sugar Island, Winch. Cat.; Michigan, A. Gray. U. P.

1492. **V. sagittata** Aiton. Arrow-leaved Violet. Rare in some sections, common in others. Howard City and Baldwin, W. J. B.; Keweenaw Co., O. A. Farwell. Th.

*1493. **V. scabriuscula** (T. & G.) Schwein. Smoothish Yellow Violet. *V. pubescens scrabiuscula* T. & G. Frequent.

1494. **V. Selkirkii** Pursh. Great-spurred Violet. Gillman; A. Gray; Keweenaw Co., O. A. Farwell. U. P.

1495. **V. septentrionalis** Greene. S. W., H. S. Pepoon.

1496. **V. sororia** Willd. Belle Isle, O. A. Farwell.

*1497. **V. striata** Aiton. Pale Violet. Keweenaw Co., O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Southward. Common. Th.

1498. **V. Rafinesqueii** Greene. Field Pansy *V. tenella* Muhl. Croswell, W. W. Wier; Detroit, O. A. Farwell.

1499. **V. tricolor** L. Pansy. Heart's-ease. Keweenaw Co., O. A. Farwell.

1500. **V. villosa** Walt. Keweenaw Co., O. A. Farwell.

CALCEOLARIA Loefl. SOLEA Spreng. IONIDIUM Vent.

*1501. C. verticillata (Ort.) Kuntze. Nodding Violet. Green Violet. *Solea concolor* Ging. Hubbardston; Ann Arbor, Allm. Cat. Scarce and local. Near Grand Rapids, Miss E. J. Cole. C. & S.

CACTACEÆ Lindl. Cactus Family.

OPUNTIA Mill.

1502. O. humifusa Raf. Western Prickly Pear. *O. Rafinesquii* Engelm. Common in Newaygo Co. along the Muskegon River, Cedar Creek Tp., Muskegon Co., C. F. Wheeler; and a stunted variety on sand barrens near Greenville, J. Satterlee; also, northward into British Am., Engelmann.

THYMELEACEÆ Reichenb. Mezereon Family.

DIRCA L.

*1503. D. palustris L. Leather-wood. Moose-wood. Woods. The tough bark used for thongs by Indians. Ann Arbor, Winch. Cat.; South Haven, L. H. Bailey; Alma, C. A. Davis, and northward. Frequent. Th.

ELAEAGNACEÆ Lindl. Oleaster Family.

LEPARGYRAEA Raf. SHEPHERDIA Nutt.

*1504. L. Canadensis (L.) Greene. Canadian Buffalo-berry. *Shepherdia Canadensis* Nutt. Gravelly banks. "On the western islands in Lake Erie," Dr. D. Cooley; Ann Arbor, Winch. Cat.; South Haven, L. H. Bailey; Flint, and northward. Abundant at Petoskey. A common short shrub, L. Superior, Whitney Cat.; Benton Harbor, C. F. Wheeler. Th.

LYTHRACEÆ Lindl. Loosestrife Family.

DECODON Gmelin.

*1505. D. verticillatus (L.) Ell. Swamp Loosestrife. Crystal Lake, Montcalm Co.; Flint; Birmingham, Oakland Co.; South Haven; Orion; Grass Lake; Fife Lake, farthest station north known. Infrequent. L. P.

LYTHRUM L.

1506. L. alatum Pursh. Near Detroit, Cooley MS., O. A. Farwell; S. Mich., Wright Cat.; Kalamazoo, Tuthill; Bay, Tuscola and Huron Counties, C. A. Davis. Malden, Detroit River,—Maclagan Canadian Cat. O. A. Farwell

1507. L. Salicaria L. Purple Loosestrife. Detroit to Bay City, Port Huron, C. K. Dodge.

MELASTOMACEÆ R. Br. Meadow-Beauty Family.

RHEXIA L.

1508. R. Virginica L. Deer-grass. Muskegon, C. D. McLouth; S. W., H. S. Pepoon.

ONAGRACEÆ Dumort. Evening-primrose Family.

ISNARDIA L. LUDWIGIA L., in part.

1509. I. palustris L. Marsh Purslane. *Ludwigia palustris* Ell. Water Purslane. Common. L. P.

LUDWIGIA L.

1510. **L. alternifolia** L. Seed-box. Dr. Wright; near Detroit, Dr. Cooley; near Algonac, C. K. Dodge; S. W., H. S. Pepoon. S. W.

1511. **L. polycarpa** Short & Peter. Dr. Pitcher and Dr. Clark. St. Clair Co., A. F. Foerste, C. K. Dodge. C. & S.

CHAMAENERION Adans. EPILOBIUM L., in part.

*1512. **C. angustifolium** (L) Scop. Great Willow-herb. *Epilobium spicatum.* Lam. *E. angustifolium* L. Great Willow Herb Fire-weed. Springs up abundantly where forests have been burned over, hence one common name.

1512a. **C. angustifolium canescens** (Wood.) Trelease. Keweenaw Co., O. A. Farwell.

EPILOBIUM L.

1513. **E. adenocaulon** Haussk. Northern Willow-herb. Trelease, Revision of Epilobium, p. 95. Keweenaw Co., O. A. Farwell. Frequent.

*1514. **E. coloratum** Muhl. Purple-leaved Willow-herb. Common. Th.

1515. **E. Hornemanii** Reichenb. Hornman's Willow-herb. Upper Wisconsin and Michigan. A. Gray. U. P.

*1516. **E. lineare** Muhl. Linear-leaved Willow-herb. Hubbardston; Flint; Alma; Stanton, and north to L. Sup. Infrequent in C. and rare or not at all in S.

1517. **E. palustre** L. Marsh Willow-herb. Frequent in swamps.

1518. **E. paniculatum** Nutt. Panicled Willow-herb. Britton and Brown.

*1519. **E. strictum** Muhl. Soft Willow-herb. Hubbardston; Flint; Ann Arbor; Macomb Co., Alma; northward to Keweenaw Co., O. A. Farwell. Frequent.

ONAGRA Adans. ŒNOTHERA L., in part.

*1520. **O. biennis** (L.) Scop. Common Evening Primrose. *Oenothera biennis* L. Common. Th.

1521. **O. Oakesiana** (A. Gray.) Britton. Oakes Evening Primrose. *Oenothera biennis Oakesiana* A. Gray. Shores of the Great Lakes.

OENOTHERA L.

1522. **O. laciniata** Hill. *O. sinuata* L. Detroit, O. A. Farwell. Rare.

1523. **O. rhombipetala** Nutt. S. W., H. S. Pepoon.

KNEIFFIA Spach. ŒNOTHERA L., in part.

1524. **K. fruticosa** (L.) Raimann. Common Sundrops. *Oenothera fruticosa* L. Palo, Ionia Co.; Flint; Macomb Co.; Ann Arbor, Allmend. Cat.; Keweenaw Co., O. A. Farwell. Th.

1525. **K. pumila** (L.) Spach. Small Sundrop. *Oenothera pumila* L. Ontonagon Falls, Whitney Cat.; So. Mich., Wr. Cat.; Oscoda, shore Lake Huron. Infrequent. Th.

HARTMANNIA Spach. ŒNOTHERA L., in part.

1526. **H. speciosa** (Nutt.) Small. Showy Primrose. Bay City, G. M. Bradford.

GAURA L.

1527. **G. biennis** L. Biennial Gaura. Dr. Wright. Malden, Ont., Maclagan, Canadian Catalogue. Detroit, O. A. Farwell.

1528. **G. coccinea** Pursh. Scarlet Gaura. White Pigeon, W. J. B.

CIRCAEA L.

*1529. **C. alpina** L. Smaller Enchanter's Nightshade. Woods. Common. Th.

*1530. **C. Lutetiana** L. Enchanter's Nightshade. Woods. Very common. Th.

HALORAGIDACEÆ Kl. & Garcke. Water-Milfoil Family.

HIPPURIS L.

1531. **H. vulgaris** L. Mare's tail. L. Sup., Can. Cat.; S. Mich., Wright Cat.; Keweenaw Co., O. A. Farwell; Black River, Cheboygan Co., B. & K.; Escanaba, E. J. Hill. Rare. Th.

1532. **H. vulgaris fluviatilis** Hart. Keweenaw peninsula, Robbins; Indian River, Cheboygan Co., C. F. Wheeler.

PROSERPINACA L.

*1533. **P. palustris** L. Mermaid-Weed. In swamps along with Alopecurus aristulatus and Ludwigia palustris. Keweenaw Co., O. A. Farwell; Cheboygan Co., B. & K.; Ann Arbor, Huron, Tuscola, Alpena Counties, C. A. Davis.

MYRIOPHYLLUM L.

1534. **M. alterniflorum** DC. St. Clair River, W. S. Cooper; Bay Co., G. M. Bradford.

1535. **M. Farwellii** Morong. Farwell's Water-Milfoil. Keweenaw Co., O. A. Farwell.

1536. **M. heterophyllum** Michx. Loose-flowered Water-Milfoil. Fruitport, E. J. Hill; Huron R. at Ypsilanti, Lyons; Macomb Co., Cooley; Manistee, F. P. Daniels. Rare.

*1537. **M. spicatum** L. Spiked Water-Milfoil. Abundant, L. Sup., Can. Cat.; Black Lake, Cheboygan Co., B. & K.; Pine Lake, Ingham Co. Common. Th.

1538. **M. tenellum** Bigelow. Slender Water-Milfoil. Gray's Manual.

*1539. **M. verticillatum** L. Whorled Water-Milfoil. Hubbardston; Flint; Macomb Co.; S. Mich., Wright Cat. Common. L. P. Ann Arbor, Alma, C. A. Davis.

ARALIACEÆ Vent. Ginseng Family.

ARALIA L.

*1540. **A. hispida** Vent. Bristly Sarsaparilla. Wild Elder. In a peat bog near the Agricultural College; also on dry clay soil, lot 21, Collegeville, Ingham Co.; Washtenaw Co., Alma, C. A. Davis. Apparently not common south of the pine region.

*1541. **A. nudicaulis** L. Wild Sarsaparilla. Alma, Ann Arbor, C. A. Davis. Common. Th.

*1542. **A. racemosa** L. Spikenard. Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell; Roscommon, Ann Arbor, C. A. Davis. Frequent. Th.

1543. **A. spinosa** L. Spreading from Detroit, O. A. Farwell.

PANAX L.

*1544. **P. quinquefolium** L. Ginseng. *Aralia quinquefolia* Decsne. & Planch. Sault Ste. Marie; Ludington; Stanton, formerly in great abundance; Hubbardston; Flint; Macomb Co.; southwest, Wright Cat. Th.

1545. **P. trifolium** L. Dwarf Ginseng or Ground-nut. *Aralia trifolia* Decsne. & Planch. Frequent northward to Keweenaw Co., O. A. Farwell. Th.

UNBELLIFERÆ B. Juss. Corrot Family.

DAUCUS L.

1546. **D. Carota** L. Wild Carrot. Keweenaw Co., O. A. Farwell; Ann Arbor, C. A. Davis; St. Clair Co., C. K. Dodge; near Grand Rapids, G. D. Sones.

ANGELICA L. ARCHANGELICA Hoffm.

*1547. **A. atropurpurea** L. Great-purple-stemmed Angelica. Alma, occasional in S., common in C. & N.; also L. Sup. Can. Cat. Th.

1548. **A. villosa** (Walt.) B. S. P. *A. hirsuta* Muhl. Emmet Co., Winch. Cat.; Pontiac; Detroit, Ann Arbor, C. A. Davis. Dry banks and open woods. Infrequent. L. P.

CONIOSELINUM Hoffm.

*1549. **C. Chinense** (L.) B. S. P. Hemlock Parsley. Ann Arbor, All. Cat.; Macomb Co.; Flint; Hubbardston and north, C. F. Wheeler. Infrequent. Th.

OXYPOLIS Raf. TIEDEMANNIA DC.

*1550. **O. rigidus** (L.) Britton. Cowbane. *Tiedemannia rigida* Coult. & Rose. Hubbardston; Ann Arbor, All. Cat. Not much N. of lat. 43. Infrequent. C. & S.

HERACLEUM L.

*1551. **H. lanatum** Michx. Low ground. Common. Th.

PASTINACA L.

*1552. **P. sativa** L. Wild Parsnip. Fields. Infrequent. Th.

IMPERATORIA L.

1553. **I. Ostruthium** L. Masterwort. Muskegon, C. D. McLouth.

POLYTAENIA DC.

1554. **P. Nuttallii** DC. Dr. Wright, also A. Gray. Rare. S.

FOENICULUM Adans.

1555. **F. Foeniculum** (L.) Karst. Fennel. Ballast grounds, Detroit, O. A. Farwell.

THASPIUM Nutt.

1556. **T. barbinode** (Michx.) Nutt. Hairy-jointed Meadow Parsnip. Jackson, S. H. Camp.

1557. **T. trifoliatum** (L.) Britton. Purple Meadow Parsnip. Confined to the two southern tiers of counties.

1558. **T. trifoliatum aureum** (Nutt.) Britton. Belle Isle, O. A. Farwell.

ERYNGIUM L.

1559. **E. aquaticum** L. Button Snakeroot. *E. yuccaefolium* Michx. White Pigeon, 1838, Dr. Wright, in University Herb; Sturgis, F. P. Daniels; S. W., H. S. Pepoon. Rare. S. W.

SANICULA L.

*1560. **S. Canadensis** L. Short-styled Snakeroot. *S. Marylandica Canadensis* Torr. Hubbardston; Flint; Ann Arbor; All. Cat.; Alma. Beech and maple woods. Infrequent. C. & S.

*1561. **S. gregaria** Bicknell. Clustered Snake-root. Alma, C. A. Davis; Agricultural College, C. F. Wheeler; Belle Isle, O. A. Farwell.

*1562. **S. Marylandica** L. Black Snake-root. Oak woods. Common. Th.

1563. **S. trifoliata** Bicknell. Large-fruited Snake-root. Alma, C. A. Davis and C. F. Wheeler.

TAENIDIA Drude. PIMPINELLA L., in part.

*1564. **T. integerrima** (L.) Drude. Yellow Pimpernel. *Pimpinella integerrima* A. Gray. Dry soil. Very abundant in Grand-Saginaw Valley. Th.

CHAEROPHYLLUM L.

*1565. **C. procumbens** (L.) Crantz. Spreading Chervil. Low woods. Frequent in Grand River Valley; Valley of the Raisin River near Dundee, C. F. Wheeler; Macomb Co. C. & S.

WASHINGTONIA Raf. OSMORRHIZA Raf.

*1566. **W. Claytoni** (Michx.) Britton. Woolly Sweet-cicely. *Osmorrhiza brevistylis* DC. The prevailing species, Winchell; probably true of counties along the Huron shore. Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Common. Th.

*1567. **W. longistylis** (Torr.) Britton. Smoother Sweet-Cicely. *Osmorrhiza longistylis* DC. Alma, C. A. Davis; Bay Co., G. M. Bradford. Common. Th.

CONIUM L.

*1568. **C. maculatum** L. Poison Hemlock. St. Clair Co., C. K. Dodge. Infrequent in C. & S. Common at Mackinac, Winch. Cat.

SIUM L.

*1569. **S. cicutaefolium** Gmel. Water-Parsnip. Clinton Co.; S. Mich., Winch. Cat.; Alma, Ann Arbor, C. A. Davis. Frequent. Th.

ZIZIA Koch.

*1570. **Z. aurea** (L.) Koch. Golden Meadow-Parsnip. Low ground. Common. Th.

1571. **Z. cordata** (Walt.) DC. Heart-leaved Alexanders. Flint; Put-in-Bay and S. W., Dr. Wright; Barron Lake, Cass Co., C. F. Wheeler. Rare. C. & S.

CARUM L.

1572. **C Copticum** (L.) Benth. Ballast grounds. Detroit, O. A. Farwell.

*1573. **Carui** L. Caraway. Naturalized in many places.

CICUTA L.

*1574. **C. bulbifera** L. Bulb-bearing Water Hemlock. Swamps. Common. Th.

*1575. **C. maculata** L. Spotted Cowbane. Beaver Poison. Musquash Root. Th.

DERINGA Adans. CRYPTOTAENIA DC.

*1576. **D. Canadensis** (L.) Kuntze. Hornwort. *Cryptotaenia Canadensis* DC. Low woods. Common. C. & S.

BERULA Hoffm.

1577. **B. erecta** (Huds.) Coville. Cut-leaved Water Parsnip. *B. angustifolia* Mert. & Koch. Winchell, and A. Gray; Kalamazoo, Tuthill; Manistee, F. P. Daniels. Infrequent. S.

HYDROCOTYLE L.

*1578. **H. Americana** L. American Marsh-Pennywort. Common in C.; Alma, C. A. Davis. L. P.

*1579. **H. umbellata** L. Umbellate Marsh-Pennywort. Woodward Lake, Ionia Co.; S. Mich., Wright Cat.; Alma, C. A, Davis. Either grows on sandy shores or is wholly aquatic with floating leaves.

ERIGENIA Nutt.

*1580. **E. bulbosa** (Michx.) Nutt. Harbinger of Spring. Common. C. & S.

CORNACEÆ Link. Dogwood Family.

CORNUS L.

*1581. **C. alternifolia** L. f. Alternate-leaved Cornel. Banks. Frequent. The dead stems bright yellow. Keweenaw Co., O. A. Farwell. Rare in Cheboygan Co., B. & K.; Alma, Ann Arbor, C. A. Davis. Th.

*1582. **C. Amomum** Mill. Silky Cornel. Kinnikinnik. *C. sericea* L. Near Detroit, O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Frequent. C. & S.

1583. **C. asperifolia** Michx. Rough-leaved Dogwood. Belle Isle, O. A. Farwell; Homer, C. F. Wheeler.

*1584. **C. Baileyi** Coulter & Evans. Bailey's Cornel. Petoskey, New Buffalo, C. F. Wheeler; abundant along E. shore of Lake Michigan on sand dunes, occasional in Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell.

*1585. **C. candidissima** Mill. Panicled Cornel. Common. L. P.

*1586. **C. Canadensis** L. Dwarf Cornel. Bunch-berry. Rare south. Constantine, C. F. Wheeler; Ann Arbor, G. D. Sones. Northward very common. Th.

1587. **C. circinata** L'Her. Round-leaved Dogwood. Klinger Lake; Hubbardston to Cheboygan Co., B. & K.; Alma, Ann Arbor, C. A. Davis and northward to Keweenaw Co., O. A. Farwell. Infrequent. Th.

*1588. **C. florida** L. Flowering Dogwood. Frequent as far north as Grand-Saginaw Valley, usually as a low tree in oak woods. C. & S.

*1589. **C. polygama** Raf. *C. purpuri* Koehne. Bay Co., G. M. Bradford; Detroit, O. A. Farwell.

1590. **C. stolonifera** Michx. Red-osier. Dogwood. Marshes, borders of streams. Very common. Th.

NYSSA L.

*1591. **N. sylvatica** Marsh. Pepperidge. Tupelo. Sour-Gum Tree. Frequent. Alma, Ann Arbor, C. A. Davis; Manistee, F. P. Daniels. L. P. Edges of swamps. Frequent.

PYROLACEÆ Agardh. Wintergreen Family.

PYROLA L.

1592. **P. asarifolia** Michx. Liver-leaf Wintergreen. *P. rotundifolia asarifolia* Hook. The Cove, L. Huron, Winch. Cat.

1593. **P. chlorantha** Swartz. Greenish-flowered Wintergreen. Pine woods. Alma. Rare south of lat. 43. Frequent northward. C. N. & N. R. R.

*1594. **P. elliptica** Nutt. Shin-leaf. Rich woods. Ann Arbor; Ionia Co.; northward to L. Superior. Common. Th.

1595. **P. minor** L. Lesser Wintergreen. Cold woods, L. Superior, A. Gray. L. P.

*1596. **P. rotundifolia** L. Round-leaved Wintergreen. Dry oak woods. Common. Th.

*1597. **P. secunda** L. One-sided Wintergreen. Rich woods. Common. Th. *P. secunda pumila* A. Gray. Grayling, G. H. Hicks; Oscoda; Cheboygan Co., B. & K.; Escanaba, E. J. Hill; Keweenaw Co., O. A. Farwell.

1598. **P. uliginosa** Torr. Bog Wintergreen. *P. rotundifolia uliginosa* A. Gray. L. Superior, Prof. J. Macoun; Quinnesec, E. J. Hill; Livingston Co., Alma, C. A. Davis; Keweenaw Co., O. A. Farwell; Cheboygan Co., Beardslee.

MONESES Salisb.

1599. **M. uniflora** (L.) A. Gray. One-flowered Wintergreen. *M. grahdiflora.* S. F. Gray. Pine forests. Montcalm Co.; Flint.; Indian River, Wheeler; Clarkston, G. H. Hicks; Fort Gratiot, Winch. Cat.; cedar swamps, Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell; Escanaba, E. J. Hill. Rare. Th.

CHIMAPHILA Pursh.

1600. **C. maculata** (L.) Pursh. Spotted Wintergreen. Oak woods. Ionia; Bangor and South Haven, Van Buren Co.; Flint. Rare or local.

*1601. **C. umbellata** (L.) Nutt. Pipsissewa. Prince's Pine. Pine woods. Common. Th.

MONOTROPACEÆ Lindl. Indian-pipe Family.

PTEROSPORA Nutt.

1602. **P. Andromedea** Nutt. Giant Bird's-nest. Pine Drops. Sitting Rabbit, Winch. Cat.; Carp River, U. P., Whitney Cat.; Keweenaw Co., O. A. Farwell; East Tawas; Elk Rapids.

MONOTROPA L.

*1603. **M. uniflora** L. Indian Pipe. Corpse Plant. Damp woods. Frequent. Th.

1637. **L. quadrifolia** L. Crosswort. Sandy soil. Ann Arbor, Winch. Cat.; Ionia Co.; Montcalm Co.; Flint; Macomb Co., Alma; northward to Oscoda Co. L. P.

*1638. **L. terrestris** (L.) B. S. P. Bulb-bearing Loosestrife. *L. stricta* Ait. Borders of marshes. Alma, Ann Arbor, C. A. Davis; Hubbardston, C. F. Wheeler.

STEIRONEMA Raf.

*1639. **S. ciliatum** (L.) Raf. Fringed Loosestrife. Swamps. Common. Th.

1640. **S. lanceolatum** (Walt.) A. Gray. Lance-leaved Loosestrife. Howard City, W. J. B,; Detroit, O. A. Farwell; low grounds, Cheboygan Co., B. & K.

*1642. **S. quadriflorum** (Sims) Hitchc. Prairie Moneywort. *S. longifolium* A. Gray. Banks of streams. Ann Arbor and Ft. Gratiot, Winch. Cat.; Ionia Co.; Flint. Frequent. L. P.

NAUMBERGIA Moench.

*1643. **N. thyrsiflora** (L.) Duby. Tufted Loosestrife. *Lysimachia thyrsiflora* L. Tufted Loosestrife. Swampy soil. Alma, Ann Arbor, C. A. Davis. Common.

TRIENTALIS L.

*1644. **T. Americana** (Pers.) Pursh. Star-flower. Damp woods. Common. Th.

ANAGALLIS L.

1645. **A. arvensis** L. Common Pimpernel. Ann Arbor, Mary H. Clark; St. Clair, C. K. Dodge; Detroit, O. A. Farwell.

DODECATHEON L.

1646. **D. Meadia** L. Shooting Star. Moist, shaded grounds, A. Gray, Dr. A. B. Lyons; Belle Isle, O. A. Farwell.

OLEACEÆ Lindl. Olive Family.

SYRINGA L.

1647. **S Persica** L. Persian Lilac. Keweenaw Co., Detroit, O. A. Farwell.

1648. **S. vulgaris** L. Lilac. Keweenaw Co., Ypsilanti, O. A. Farwell. Escaped from cultivation.

FRAXINUS L.

*1649. **F. Americana** L. White Ash. Common. Th.

*1650. **F. lanceolata** Borck. Green Ash. *F. viridis* Michx. f. Alma, C. A. Davis; Ann Arbor, Winch. Cat.; and northward. Rare.

*1651. **F. nigra** Marsh. Black Ash. *F. sambucifolia* Lam. Common. Th.

*1652. **F. Pennsylvanica** Marsh. Red Ash. *F. pubescens* Lam. Low grounds. Ionia Co.; S. Mich., Wright. Cat.; Drummond's I., Winch. Cat.; along Black River, Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell Scarce.

*1653. **F. quadrangulata** Michx. Blue Ash. Rich woods. Infrequent. Ann Arbor, C. A. Davis; Lenawee Co., W. J. B.

GENTIANACEÆ Dumort. Gentian Family.

ERYTHRAEA Neck.

*1654. **E. Centaurium** (L.) Pers. Centaury. Agricultural College grounds, L. H. Bailey. The only locality known in the State.

SABBATIA Adans.

1655. **S angularis** (L.) Pursh. Marshes, Cass Co., Univ. Herb., 1838; Goguac Lake, near Battle Creek, V. M. Spalding; S. Mich., Wright's Cat.; Berrien Co., E. F. Smith. Rare.

GENTIANA L.

*1656. **G. Andrewsii** Griesb. Closed Gentian. River banks. Frequent.

*1657. **G. crinata** Froel. Fringed Gentian. Low grounds, Lenawee Co., G. F. Comstock; Ann Arbor, Alma, C. A. Davis, Winch. Cat.; South Haven, L. H. Bailey; Mackinac, Whitney; Ionia Co. Frequent. Th.

*1658. **G. detonsa** Röttb. Samller Fringed Gentian. *G. serrata* Gunner. Tamarack swamps. Frequent. Th.

1659. **G. flavida** A. Gray. Yellowish Gentian. *G. alba* A. Gray. Low meadows and borders of woods. Ionia Co.; Clinton Co.; Ann Arbor and S. W., Winch. Cat.; to L. Superior. Rare or local.

1660. **G. linearis** Froel. Narrow-leaved Gentian. *A. latifolia* A. Gray. Keweenaw Co., O. A. Farwell; Escanaba, C. F. Wheeler. Creek bottoms.

*1661. **G. puberula** Michx. Downy Gentian. Barrens. Ann Arbor, Miss Clark, Miss Allmendinger. Rare. S.

1662. **G. quinquefolia** L. Stiff Gentian. Moist hillsides. Ann Arbor, Winch. Cat.; Ionia Co. Occasional. C. & S.

1663. **G. quinquefolia occidentalis** (A. Gray) A. S. Hitchcock. Lenawee Co., G. F. Comstock; Ann Arbor, Winch. Cat.; Hubbardston; Macomb Co. Infrequent.

1664. **G. rubricaulis** Schwein. Red-stemmed Gentian. *G. linearis lanceolata* A. Gray. Pt. au Chene, L. Mich., Winch. Cat.; low ground near Black Lake, Cheboygan Co., B. & K.; Minnesota and along L. Superior, A. Gray. N. & U. P.

1665. **G. Saponaria** L. Soapwort Gentian. Moist woods. Macomb Co.; S. Mich., Wright Cat.; St. Clair, C. K. Dodge. Infrequent. S.

FRASERA Walt.

*1666. **F. Carolinensis** Walt. American Columbo. Jackson Co., and westward, Winch. Cat.; Ann Arbor, Miss Clark; Constantine, Three Rivers, C. F. Wheeler; Ingham Co., W. J. B.; Sturgis, F. P. Daniels; Addison, O. C. McLouth; Lenawee Co., G. F. Comstock; Grand Rapids; Macomb Co.; S. W., H. S. Pepoon. Scarce. C. & S.

TETRAGONANTHUS G. Gmel. HALENIA Borck.

1667. **T. deflexus** (J. E. Smith) Kuntze. Spurred Gentian. *Halenia deflexa* Griseb. Shore of Little Traverse Bay, near Harbor Point; Drummond's Is. and St. Helen's Is., Winch. Cat.; northward to Lake Superior; shore Black Lake, Cheboygan Co.; Keweenaw Co., O. A. Farwell. Frequent.

1668. **T. deflexus Brentonianus** (Griseb.) Britton. Keweenaw Co., O. A. Farwell.

BARTONIA Muhl.

*1669. **B. Virginica** (L.) B. S. P. *B. tenella* Willd. Open woods. S. Mich., Wright Cat.; Algonac; Hubbardston; Flint; Grayling; St. Clair Co., C. K. Dodge; Grass Lake, C. A. Davis. Rare.

MENYANTHACEÆ G. Don. Buckbean Family.

MENYANTHES L.

*1670. **M. trifoliata** L. Buckbeam. Bogs. Th.

LIMNANTHEMUM Gmelin.

1671. **L. lacunosum** (Vent.) Griseb. Floating Hart. Keweenaw Pt., Dr. Robbins.

APOCYNACEÆ Lindl. Dogbane Family.

VINCA L.

1672. **V. minor** L. Periwinkle. Occasionally escaped from cultivation.

APOCYNUM L.

*1673. **A. androsaemifolium** L. Spreading Dogbane. Borders of woods. Common. Th.

*1674. **A. cannabinum** L. Indian Hemp. Low grounds. Exceedingly variable in size, habit, shape of leaves, pubescence. Common. Th.

*1675. **A. cannabinum glaberrimum** DC. Belle Isle, O. A. Farwell.

1676. **A. hypericifolium** Ait. Clasping-leaved Dogbane. Washington, Dr. D. Cooley; Flint, Dr. D. Clark; St. Clair Co., C. K. Dodge.

1677. **A. Milleri** Britton. Miller's Dogbane. Detroit, O. A. Farwell.

1678. **A. pubescens** R. Br. Velvet Dogbane. Detroit, O. A. Farwell.

ASCLEPIADACEÆ Lindl. Milkweed Family.

ASCLEPIAS L.

1679. **A. amplexicaulis** J. E. Smith. *A. obtusifolia* Michx. Baldwin, Lake Co., W. J. B.; Sturgis; Barron Lake. C. F. Wheeler; Niles, O. J. Stilwell. Rare.

*1680. **A. exaltata** (L.) Muhl. Poke Milkweed. *A. phytolaccoides* Pursh. Moist grounds. Ann Arbor; Ft. Gratiot, Winch. Cat.; Alma; Ionia Co.; Flint; Macomb Co. Infrequent. C. & S.

*1681. **A incarnata** L. Swamp Milkweed. Banks of streams. Common. Th.

1682. **A. pulchra** Ehrh. Hairy Milkweed. Hansen's Island. Algonac Co., J. W. Stacy.

1683. **A. purpurascens** L. Purple Milkweed. Woods. Ann Arbor, Winch. Cat.; Ionia Co.; Clinton Co.; Flint; Macomb Co.; Huron Co.; Tuscola Co. S. & S.

1684. **A. quadrifolia** Jacq. Four-leaved Milkweed. Ann Arbor, Winchell's Catalogue.

*1685. **A. Syriaca** L. Common Milkweed. *A. Cornuti* Dec. Fields. Common. Th.

1686. **A. Sullivantii** Engelm. Sullivan's Milkweed. Walpole Island, C. K. Dodge; Tuscola Co., C. A. Davis; Esserville, C. F. Wheeler.

*1687. **A. tuberosa** L. Butterfly-weed. Pleurisy-root. Sandy soil. Ranges from Canada to Florida and from the Saskatchewan Valley to Texas. Common.

1688. **A. verticillata** L. Whorled Milkweed. Barrens, S. Mich., Wright Cat.; Monroe Co., Grand Rapids, C. F. Wheeler; Washington, Macomb Co., Dr. D. Cooley.

ACERATES Ell.

1689. **A. Floridana** (Lam.) Hitchc. Florida Milkweed. *A. longifolia* Ell. South Haven, L. H. Bailey; Bay Co., G. M. Bradford; Detroit, O. A. Farwell; Tuscola Co., C. A. Davis.

1690. **A. viridiflora** (Raf.) Eaton. Green Milkweed. C. & S. Grand Rapids, Geo. D. Sones; Ft. Gratiot, Winch. Cat.; S. W., Wright Cat.; Ann Arbor, C. A. Davis; S. W., H. S. Pepoon; Detroit, O. A. Farwell. Rare.

CYNANCHUM L. VINCETOXICUM Walt., in part.

*1691. **C. nigrum** (L.) Pers. Black Swallow-wort. *Vincetoxicum nigrum* Moench. Escaped from cultivation on College grounds.

CONVOLVULACEÆ Vent. Morning-Glory Family.

CONVOLVULUS L.

1692. **C. Americanus** (Sims.) Greene. S. W., H. S. Pepoon.

*1693. **C. arvensis** L. Bindweed. Alma, Ann Arbor, C. A. Davis; Hubbardston; College grounds, along C. & G. T. R. R. Spreading.

1695. **C. hederacea** Wall. Chinese Morning glory. Bay City, G. M. Bradford.

*1696. **C. repens** L. Trailing Bindweed. *C. sepium repens* A. Gray. Frequent.

*1697. **C. sepium** L. Hedge Bindweed. Low grounds. Common. Th.

1698. **C. spithamaeus** L. Upright Bindweed. Sandy fields. Local. Th.

1698a. **C. stans** Michx. Keweenaw Co., O. A. Farwell.

IPOMOEA L.

1699. **I Pandurata** (L.) Meyer. Wild Potato-vine. Man-of-the-Earth. Win. Cat.; Niles, I. N. Mitchell; Coleman Cat.; Lenawee Co., W. J. B. Infrequent. S.
1700. **I. purpurea** (L.) Roth. Morning Glory Bay Co., G. M. Bradford; Ypsilanti, Detroit, O. A. Farwell; Alma, C. A. Davis; Ann Arbor. Escaped from cultivation.

CUSCUTACEÆ Dumort. Dodder Family.

CUSCUTA L.

*1701. **C. arvensis** Beyrich. Field Dodder. Lansing, C. F. Wheeler; Rochester, W. S. Cooper.
*1702. **C. Cephalanthi** Engelm. Button-bush Dodder. *C. tenuiflora* Elgelm. Maple River Valley; Coleman Cat.; on willows along Cedar River east of the College.
1703. **C. Coryli** Engelm. Hazel Dodder. *C. inflexa* Engelm. Port Huron, C. K. Dodge; Rochester and Algonac, W. S. Cooper.
1704. **C. Epithymum** Murr. Clover Dodder. Introduced at Salem, with alfalfa seed.
*1705. **C. Gronovii** Willd. Gronovius' Dodder. Low grounds. Frequent. C. & S.
1706. **C Polygonorum** Engelm. Smart-weed Dodder. *C. chlorocarpa* Engel. S. Haven, Mrs. Millington. S. & W.

POLEMONIACEÆ DC. Phlox Family.

PHLOX L.

1707. **P. bifida** Beck. Cleft Phlox. Rare. Shores of Barron Lake; Klinger Lake, the farthest station east known for this interesting plant, C. F. Wheeler. S. W.
*1708. **P. divaricata** L. Wild Blue Phlox. S. W., H. S. Pepoon; woods and fields, northward to Kingsley and Frankfort. Common. L. P.
1709. **P. paniculata** L. Garden Phlox. Keweenaw Co., O. A. Farwell. Introduced.
1710. **P. pilosa** L. Downy Phlox. Dry, sandy fields, Ann Arbor, Winch. Cat.; Ionia Co.; Flint; Macomb Co. Infrequent.
1711. **P. subulata** L. Ground or Moss Pink. S. Mich., Winch. Cat.; shore of Barron Lake, C. F. Wheeler; Belle Isle, O. A. Farwell. Common in cultivation.

HYDROPHYLLACEÆ Lindl. Water-leaf Family.

HYDROPHYLLUM L.

*1712. **H. appendiculatum** Michx. Appendaged Water-leaf. Moist hillsides and rich woods, northward to Frankfort. Frequent. L. P.
*1713. **H. Canadense** L. Broad-leaved Water-leaf. Damp, rich woods. Frequent. C. & S.
*1714. **H. Virginicum** L. Virginia Water-leaf. Rich woods. Frequent. C. & S.

PHACELIA Juss.

1715. **P. Franklinii** (R. Br.) A. Gray. Franklin's Phacelia. "Shores of L. Superior, especially on Isle Royale," A. Gray and Dr. A. B. Lyons. U. P.

BORAGINACEÆ Lindl. Borage Family.

CYNOGLOSSUM L.

*1716. **C. officinale** L. Common Hound's-Tongue. Roadsides. Frequent. Th.
*1717. **C. Virginicum** L. Wild Comfrey. Open woods, Comins, Oscoda Co.; Escanaba, where it replaces C. officinale as a weed, E. J. Hill. Frequent. Th.

LAPPULA Moench. ECHINOSPERMUM Sw.

1718. **L. Americana** (A. Gray) Rydberg. Nodding Stickseed. *Echinospermum deflexum Americanum* A. Gray. Cheboygan Co., B. & K.

*1720. **L. Lappula** (L.) Karst. Stickseed. *Echinospermum Lappula* Lehm. A worthless weed along roadsides and in waste places. Th.

*1721. **L. Virginiana** (L.) Greene. Virginia Stickseed. *Echinospermum Virginicum* Lehm. Woods. Frequent. Th.

MERTENSIA Roth.

1722. **M. paniculata** (Ait.) G. Don. Tall Lungwort. Whitney Cat.; Gray; Keweenaw Co., O. A. Farwell. U. P.

1723. **M. Virginica** (L.) DC. Virginia Cowslip. Lungwort. Blue Bells. Near Adrian, Mrs. I. H. Wheeler; S. E. of Grand Rapids, G. D. Sones. Rare.

MYOSOTIS L.

1724. **M. arvensis** (L.) Lehm. Field Scorpion Grass. Lapeer, Mrs. M. Owen.

1725. **M. laxa** Lehm. Smaller Forget-me-not. Lapeer, Mrs. M. Owen.

1726. **M. palustris** (L.) Lam. Escaped from gardens. Port Huron, M. Allenbruch.

*1727. **M. Virginica** (L.) B. S. P. Spring Scorpion Grass. *M. verna* Nutt. Winch. Cat.; Coleman Cat.; Flint, Dr. Clark; Ann Arbor, I. N. Mitchell; shore of Park Lake, Clinton Co., C. F. Wheeler. Rare. S.

LITHOSPERMUM L.

*1729. **L. arvense** L. Wheat-Thief. Red-root. Corn Gromwell. In Wheat fields. Introduced from the old world. Common. C. & S.

*1730. **L. canescens** (Michx.) Lehm. Puccoon of the Indians. Sandy fields, Ann Arbor, Winch. Cat.; Ionia Co.; Macomb Co., and northward. Th.

*1731. **L. Gmelini** (Michx.) Hitch. Hairy Puccoon. *L. hirtum* Lehm. Light sand. Tuscola Co., C. A. Davis; Bay Co., G. M. Bradford; Berrien Co., H. S. Pepoon.

*1732. **L. latifolium** Michx. American Gromwell. Borders of woods S. Mich., Winch. Cat.; Ionia Co.; Flint; Bois Blanc and other islands in Detroit River; Maclagan, Canadian Catalogue. Frequent. C. & S.

1732a. **L. linearifolium** Goldie. Narrow-leaved Puccoon. *L. augustifolium* Michx., not Forsk. Dr. A. B. Lyons; "Mich.," Gray's Manual; Port Huron, C. K. Dodge.

*1733. **L. officinale** L. Common Gromwell. Roadsides. Tuscola Co., C. A. Davis. Infrequent. Th.

ONOSMODIUM Michx.

1734. **O. molle** Michx. Soft-haired False Gromwell. *O. Carolinianum molle* A. Gray. Only noticed by Dr. D. Cooley, Macomb Co.

SYMPHYTUM L.

*1735. **S. officinale** L. Common Comfrey. Escaped from gardens. Infrequent.

BORAGO L.

1736. **B. officinalis** L. A weed in parts of St. Clair Co., C. K. Dodge.

ECHIUM L.

1737. **E. vulgare** L. Blue-weed. Vulcan, E. J. Hill; Bay Co., G. M. Bradford.

VERBENACEÆ J. St. Hil. Vervain Family.

VERBENA L.

1738. **V. angustifolia** Michx. Narrow-leaved Vervain. Dry grounds. Winch. Cat. S.

1739. **V. bracteosa** Michx. Large-bracted Vervain. Waste places. Coleman Cat.; Kalamazoo, L. H. Bailey; Port Huron, C. K. Dodge; Manistee, F. P. Daniels; Roscommon Co., C. A. Davis.

1740. **V. bracteosa x stricta.** Near Port Huron, C. K. Dodge.
*1741. **V. hastata** L. Blue Vervian. Roadsides. Common. Th.
1742. **V. hastata oblongifolia** Nutt. *V. urticifolia riparia* (Raf.) Britton. A probable hybrid between V. hastata and V. urticifolia. South of Marshall, where no other species were found excepting the two last named, W. J. B.; Detroit, O. A. Farwell.
1744. **V. officinalis** L. Detroit, O. A. Farwell.
1745. **V. stricta** Vent. Mullein-leaved Vervain. Grand Rapids, H. C. Skeels; Port Huron, C. K. Dodge; Detroit, O. A. Farwell.
*1746. **V. urticifolia** L. White Vervain. Waste places. Common. Th.
1747. **V. urticifolia riparia** (Raf.) Britton. Detroit, O. A. Farwell.

LIPPIA L.

1748. **L. lanceolata** Michx. Fog-fruit. Coleman Cat.; Niles, I. N. Mitchell; Algonac, W. S. Cooper. S. W.

LABIATÆ B. Juss. Mint Family.

AJUGA L.

*1750. **A. reptans** L. Bugle. Bay City, G. M. Bradford.

TEUCRIUM L.

*1751. **T. Canadense** L. American Germander. Wood Sage. Low grounds. Infrequent at Black Lake, Cheboygan Co., B. & K. Alma, Ann Arbor, C. A. Davis. L. P.
1752. **T. menthifolium** Bickwell, Algonac, W S. Cooper; Alma, where the type was collected.
1753. **T. occidentalis** A. Gray. Hairy Germander. Gaylord, G. L. Stewart.

SCUTELLARIA L.

1754. **S. cordifolia** Muhl. Heart-leaved Skullcap. *S. versicolor* Nutt. Banks of streams. S. Mich., Wright Cat.
*1755. **S. galericulata** L. Marsh Skullcap. Alma, C. A. Davis. Low grounds. Common. Th.
*1756. **S. lateriflora** L. Mad-dog Skullcap. Roscommon, Alma, Ann Arbor, C. A. Davis. Low grounds. Common. Th.
1757. **S. parvula** Michx. Small Skullcap. Ionia, E. F. Smith; islands east of Georgian Bay, Bell, Can. Cat. Infrequent.
1758. **S. pilosa** Michx. Hairy Skullcap. S. W., Winch. Cat.; Gray's Manual, 6th edition.

MARRUBIUM L.

*1759. **M. vulgare** L. Common Horehound. Roadsides, escaped from cultivation. Ionia Co.; Flint; S. Mich., Winch. Cat.; Port Austin; Grand Traverse and Benzie Counties. Frequent.

AGASTACHE Clayt. LOPHANTHUS Benth.

*1760. **A. nepetoides** (L.) Kuntze. Catnep Giant-Hyssop. *Lophanthus nepetoides* Benth. Low grounds. Ionia Co.; Flint; S. W., Wright Cat.; Grosse Isle, Miss Clark. C. & S. Infrequent.
1761. **A. scrophulariaefolia** (Willd.) Kuntze. Figwort. Giant-Hyssop. *Lophanthus scrophulariaefolius* Benth. Low grounds. S. Mich., Wright Cat.; Lenawee Co., G. F. Comstock; Macomb Co.; Hubbardston. Infrequent. C. & S.

NEPETA L.

*1762. **N. Cataria** L. Catnep. Near dwellings. Common. Th.

GLECOMA L. NEPETA L., in part.

*1763. **G. hederacea** L. Gill-over-the-Ground. *Nepeta Glechoma* Benth. Sparingly escaped from culture. Flint; Grand Rapids, Coleman Cat.; Cassopolis; Alma; Ann Arbor. C. & S.

DRACOCEPHALUM L.

1764. **D. parviflorum** Nutt. American Dragon-head. This interesting plant was first detected in the Lower Peninsula in June, 1878, when it was found in Roscommon Co., south of Houghton Lake. Alcona Co.; Hubbardston, Ionia Co.; S. W., H. S. Pepoon. Rare southward.

PRUNELLA L. BRUNELLA.

1765. **P. vulgaris** L. Common Self-heal or Heal-all. *Brunella vulgaris* L. "Brunella" is not the correct name. Fields. Occasionally the flowers are white. Common. Th.

PHYSOSTEGIA Benth.

1766. **P. Virginiana** (L.) Benth. Wet grounds; varies greatly. Ann Arbor, Winch. Cat.; S. W., Wright Cat.; Alma; Muir; Kalamazoo, L. H. Bailey; Flint, Dr. Clark, northward to Keweenaw Co., O. A. Farwell.

GALEOPSIS L.

1767. **G. Ladanum** L. Red Hemp-Nettle. Ft. Gratiot and Sault de Ste. Marie, Winch. Cat.

1768. **G. Tetrahit** L. Common Hemp-Nettle. Rare in C. of the State. Abundant at Mackinac, Winch. Cat.; Cheboygan Co., B. & K.; Alma; Harbor Springs, C. F. Wheeler. Th.

LEONURUS L.

*1769. **L. Cardiaca** L. Common Motherwort. Waste grounds. Common.

LAMIUM L.

*1770. **L. amplexicaule** L. Dead Nettle. Naturalized in gardens. In fields west of Ann Arbor, C. A. Davis.

1771. **L. maculatum** L. Escaped from cultivation. St. Clair Co., C. K. Dodge.

STACHYS L.

1772. **S. aspera** Michx. Rough Hedge Nettle. Alma, Ann Arbor, C. A. Davis. Wet grounds. Infrequent. L. P.

*1773. **S. aspera glabra** Gray. Rare.

1774. **S. hyssopifolia** Michx. Hyssop Hedge Nettle. Wet grounds. S. Mich., Wright Cat.; Stanton, E. J. Quackenbush. Rare. Th.

1775. **S. palustris** L. Hedge Nettle. Keweenaw Co., O. A. Farwell.

MONARDA L.

1776. **M. didyma** L. Bee-Balm. Oswego Tea. Rare in Michigan. Barron Lake, C. F. Wheeler; Flint, Dr. Clark; St. Clair Co., C. K. Dodge. S. W.

*1777. **M. fistulosa** L. Wild Bergamot. Sandy soil. Common. Th.

1778. **M. media** Willd. Purple Bergamot. Near Detroit, O. A. Farwell.

1779. **M. mollis** L. Pale Wild Bergamot. *M. scabra* Beck. Port Huron, C. K. Dodge; Ypsilanti, Belle Isle, O. A. Farwell.

1780. **M. punctata** L. Horse-Mint. Sandy soil. S. Mich., Wright Cat.; S. Haven, L. H. Bailey; Grand Haven, G. D. Sones. Infrequent. S.

BLEPHILIA Raf.

*1781. **B. ciliata** (L.) Raf. Dry ground. Ionia Co.; Ann Arbor, and Sault de Ste. Marie. Winch. Cat. Scarce. Th.

*1782. **B. hirsuta** (Pursh.) Torr. Hairy Blephilia. Low woods. S. Mich., Wright Cat.; Alma; Ionia Co. C. & S.

HEDEOMA Pers.

1783. **H. hispida** Pursh. Rochester, O. A. Farwell.

*1784. **H. pulegioides** (L.) Pers. American Pennyroyal. Fields. Hubbardston; Rochester; S. Mich., Wright Cat.; Macomb Co.; Lenawee Co., W. J. B. C. & S.

SATUREIA L.

1785. **S. hortensis** L. Summer Savory. St. Clair Co. near Capac, C. K. Dodge.

CLINOPODIUM L. CALAMINTHA Moench.

1786. **C. glabrum** (Nutt.) Kuntze. Low Calamint. *Calamintha Nuttallii* Benth. Drummond's Island, and northward, common. Not seen in C. of the State; S. E., Wright Cat.

*1787. **C. vulgare** L. White Basil. *Calamintha Clinopodium* Benth. Fields. Ionia Co.; Fort Gratiot, Winch. Cat., and northward to Lake Superior. Th.

HYSSOPUS L.

1788. **H. officinalis** L. Escaped from gardens. Flint; S. W., Winch. Cat.

KOELLIA Moench. PYCNANTHEMUM Michx.

*1789. **K. flexuosa** (Walt.) MacM. Narrow-leaved Mountain-Mint. *Pycnanthemum linifolium* Pursh. Detroit, O. A. Farwell; Manistee, F. P. Daniels.

1790. **K. Virginiana** (L.) MacM. Virginia Mountain-Mint. *Pycnanthemum lanceolantum* Pursh. Low grounds. Ionia Co.; Flint; Macomb Co.; S. W., H. S. Pepoon, and southward. Frequent. C. & S.

THYMUS L.

1791. **T. Serpyllum** L. Creeping Thyme. Flint, Coleman Cat. Rarely escaped from gardens.

*1792. **T. vulgaris** L. Common Thyme. College grounds.

LYCOPUS L.

*1793. **L. Americanus** Muhl. Cut-leaved Water Hoarhound. *L. sinuatus* Ell. Frequent.

1793a. **L. communis** Bicknell. Bugle-weed. Alma, Ann Arbor, C. A. Davis. Low grounds. Common L. P.

1794. **L. lucidus** Turcz. Western Water Hoarhound. Port Huron, C. K. Dodge.

1794a. **L. Macrophyllus** Benth. Thin-leaved Bugle-weed. *L. membranacea* Bicknell. Keweenaw Co., O. Farwell.

1794b. **L. membranaceus** Bicknell. Thin-leaved Bugle-weed. "Mich." Britton's Manual.

1795. **L. rubellus** Moench. Water Hoarhound. Clinton Co., E. F. Smith; St. Clair Co., Brotherton; Detroit, O. A. Farwell.

MENTHA L.

1797. **M. arvensis** L. Corn Mint. Algonac, C. K. Dodge.

*1798. **M. Canadensis** L. Wild Mint. Low grounds. Common. Th.

1799. **M. Canadensis glabrata** Benth. Keweenaw Co., O. A. Farwell.

1800. **M. citrata** Ehrh. Bergamot Mint. Algonac, Walpole Island, C. K. Dodge.

*1801. **M. piperita** L. Peppermint. Along streams. Extensively cultivated in St. Joseph and Wayne Counties for the oil. Common. Th.

1802. **M. sativa** L. Marsh Whorled Mint. Sturgis, F. P. Daniels.

*1803. **M. spicata** L. Spearmint. *M. viridis* L. Roadsides. Escaped from cultivation. Frequent.

COLLINSONIA L.

*1804. **C. Canadensis** L. Stone-root. Rich-weed. Rich woods. Ionia Co.; Flint; Detroit; Ann Arbor, and S. W., Winch. Cat. Frequent. C. & S.

SOLANACEÆ Pers. Potato Family.

PHYSALODES Boehm. NICANDRA Adans.

*1805. **P. physalodes** (L.) Britton. Apple-of-Peru. *Nicandra physalodes* Gaertn. Gardens. Ann Arbor, Winch. Cat.; Flint; Ionia Co.; Detroit, O. A. Farwell. Scarce.

PHYSALIS L.

1806. **P. heterophylla** Nees. Ground-Cherry. Detroit and Orion, O. A. Farwell; S. W., H. S. Pepoon.

1807. **P. heterophylla ambigua** (A. Gray) Rydberg. *P. Virginiana ambigua* A. Gray. Belle Isle and Keweenaw Co., O. A. Farwell.

1808. **P. heterophylla nyctaginea** (Dunal.) Rydberg. Howard City, W. J. B.; Barron Lake, Ronald, C. F. Wheeler.

1809. **P. ixocarpa** Brot. Tomatillo. Port Huron, C. K. Dodge.

*1810. **P. lanceolata** Michx. Prairie Ground-Cherry. Sandy soil. S. W., H. S. Pepoon; Ionia Co.; Alma. Frequent. Th.

1811. **P. Philadelphica** Lam. Philadelphia Ground-Cherry. Muskegon, C. D. McLouth; Belle Isle, O. A. Farwell.

1812. **P. pubescens** L. Low Hairy Ground-Cherry. Naturalized in West Bay City, G. M. Bradford.

*1813. **P Virginiana** Mill. Virginia Ground-Cherry. Light, sandy soil. Common. Th.

1814. **P. Virginiana vulgaris** Rydb. Grand Rapids, H. C. Skeels; Muskegon, W. J. B.; White Cloud, C. F. Wheeler.

LEUCOPHYSALIS Rydberg.

1815. **L. grandiflora** (Hook.) Rydberg. Large White-flowered Ground-Cherry. *Physalis grandiflora* Hook. "Clearings." First collected in L. P. near Farwell, in June, 1876, where it seemed to be at home. Near the mouth of the Au Sable River, Iosco Co., June, 1878, C. B. Cochran; "S. shore of L. Superior," A. Gray; Keweenaw Co., O. A. Farwell; Escanaba, E. J. Hill; Indian River, C. F Wheeler. N. & U. P.

SOLANUM L.

*1816. **S. Carolinense** L. Horse-Nettle. Introduced from the southwest; Port Huron, C. K. Dodge.

*1817. **S. Dulcamara** L. Bittersweet. Nightshade. Becoming common, especially in low land and swamps. C. & S.

*1818. **S. nigrum** L. Black Nightshade. Moist, cultivated fields. Common. C. & S.

*1819. **S. rostratum** Dunal. Beaked Nightshade. Introduced from the southwest.

LYCIUM L.

*1820. **L. vulgare** (Ait. f.) Dunal. Matrimony Vine. Escaped from gardens in places.

HYOSCYAMUS L.

1821. **H. niger** L. Black Henbane. Macomb Co.; Ft. Gratiot; Mackinac, abundant, Winch. Cat.

DATURA L.

1822. **D. Metel** L. Ypsilanti and Detroit, O. A. Farwell. Waste grounds. Rare.

*1823. **D. Stramonium** L. Common Stramonium, or Thorn-Apple. Alma, Ann Arbor, C. A. Davis; Port Huron, C. K. Dodge. Roadsides. Frequent. C. & S.

*1824. **D. Tatula** L. Purple Thorn-Apple. Flint; Macomb Co.; Berrien Co., H. S. Pepoon; Alma, Ann Arbor, C. A. Davis. C. & S.

NICOTIANA L.

1825. **N. rustica** L. Wild Tobacco. Emmet Co., cultivated by the Indians, Winch. Cat.

PETUNIA Juss.

1826. **P. axillaris** (Lam.) B. S. P. White Petunia. Waste places. Ypsilanti and Detroit, O. A. Farwell.

SCROPHULARIACEÆ Lindl. Figwort Family.

VERBASCUM L.

*1827. **V. Blattaria** L. Moth Mullein. Roadsides. S. Mich.; Ionia Co.; Detroit; Sturgis, F. P. Daniels; Alma, Ann Arbor, C. A. Davis.

*1828. **V. Thapsus** L. Common Mullein. Fields and roadsides everywhere, a common weed. Cultivated in England under the name of Aaron's Rod. Th.

CYMBALARIA Medic.

1829. **C. Cymbalaria** (L.) Wettst. Kenilworth Ivy. Lake shore at Elk Rapids, W. S. Cooper.

LINARIA Hill.

*1830. **L. Canadensis** (L.) Dumont. Wild Toad-Flax. S. shore of Saginaw Bay, Winch. Cat.; barrens in Clare Co. Infrequent or wanting in C. and S. Mich. Grand Rapids, G. D. Sones; Ann Arbor, Grass Lake, C. A. Davis.

*1831. **L. Linaria** (L.) Karst. Butter-and-Eggs. *L. vulgaris* Mill. Alma, Ann Arbor, C. A. Davis. Roadsides. Spreading from cultivation.

CHAENORRHINUM Reichb. LINARIA Hill., in part.

1832. **C. minus** (L.) Lange. *Linaria minor* L. St. Clair Co., C. K. Dodge; Detroit, O. A. Farwell.

SCROPHULARIA L.

1833. **S. leporella** Bicknell. Hare Figwort. Port Huron, C. K. Dodge; Island Lake, Vestaburg, C. F. Wheeler.

*1834. **S. Marilandica** L. Maryland Figwort. *S. nodosa Marylandica* A. Gray. Rich soil. Keweenaw Co., Belle Isle, O. A. Farwell; Alma, Ann Arbor, C. A. Davis, and southward. Frequent. Th.

CHELONE L.

*1835. **C. glabra** L. Snake-head. Borders of streams. Alma, Ann Arbor, C. A. Davis. Frequent. Th.

PENTSTEMON Soland.

1836. **P. canescens** Britton. Beard-tongue. *P. laevigatus canescens* Britton. Detroit, O. A. Farwell.

1837. **P. Digitalis** (Sweet) Nutt. *P. laevigatus Digitalis* A. Gray. Near Adrian, Mrs. I. H. Wheeler.

*1838. **P. hirsutus** (L.) Willd. Hairy Beard-tongue. *P. pubescens* Solander. Dry soil. Ann Arbor, Winch. Cat.; to Mackinac, G. H. Hicks; Alma. Common. Th.

1839. **P. Pentstemon** (L.) Britton. Smooth Beard-tongue. *P. laevigatus* Soland. Grand Rapids, Mrs. A. J. Peters.

COLLINSIA Nutt.

1840. **C. parviflora** Dougl. Small-flowered Collinsia. "Shady, moist ground, Upper Michigan," Gray, Fl. N. A.; Keweenaw Co., O. A. Farwell.

*1841. **C. verna** Nutt. Blue-eyed Mary. A beautiful little fall annual with blue and white flowers. Moist woods. Ann Arbor, Winch. Cat.; S. W., Wright's Cat.; Ionia Co., and probably reaches its N. limits in Gratiot Co. C. & S.

MIMULUS L.

1842. **M. alatus** Soland. Sharp-winged Monkey-flower. Sturgis, F. P. Daniels, S. W.

1843. **M. Jamesii** T. & G. James' Mimulus. Abundant at Mackinac, Winch. Cat.; "Upper Michigan," A. Gray; Petoskey; Hubbardston, Wheeler; Grayling, G. H. Hicks; Comstock, Kalamazoo Co., Tuthill; rare southward; near Millers, Ind., Flora of Cook Co., Illinois. Th.

1844. **M. moschatus** Dougl. Musk-flower. Keweenaw Co., O. A. Farwell.

*1845. **M. ringens** L. Monkey-flower. Wet places. Common. Th.

GRATIOLA L.

1846. **G. Virginiana** L. Clammy Hedge-Hyssop. S. Mich., Winch. Cat.; Keweenaw Pt., Dr. Robbins. Rare.

ILYSANTHES Raf.

1847. **I. attenuata** (Muhl.) Small. Detroit, O. A. Farwell; Bay Co., G. M. Bradford.
*1848. **I. dubia** (L.) Barnhart. *I. gratioloides* (L.) Benth. Long-stalked False Pimpernel. Near Algonac, J. W. Stacey. Low grounds. Frequent. C. & S.

LIMOSELLA L.

1849. **L. tenuifolia** Hoffm. Narrow-leaved Mudwort. *L. aquatica tenuifolia* Hoffmann. U. P., Dr. A. B. Lyons.

SYNTHYRIS Benth. WULFENIA Greene not Jacq.

1850. **S. Bullii** (Eaton) Barnhart. *Wulfenia Houghtoniana* (Benth). Greene. Oak barrens. S. Mich., Wright's Cat.; Ionia Co., three miles S. of Saranac, probably reaches here its N. limits; Barry Co., L. H. Bailey; Sturgis, F. P. Daniels. Rare.

VERONICA L.

*1851. **V. agrestis** L. Field Speedwell. Sandy fields. Coleman Cat.; and Flint. Rare.
*1852. **V. Americana** Schweinitz. American Brooklime. Brooks. Common. Th.
*1853. **V. Anagallis-aquatica** L. Water Speedwell. In springs and brooks. Frequent. Th.
*1854. **V. arvensis** L. Corn Speedwell. Cultivated and waste grounds. Common. Th.
1855. **V. Byzantina** (Sibth. & Smith) B. S. P. Buxbaum's Speedwell. *V. Buxbaumii* Tenore. Waste grounds, Coleman Cat. Corunna, G. H. Hicks; Port Huron, C. K. Dodge.
*1856. **V. Chamaedrys** L. Germander Speedwell. Escaped from cultivation. Port Huron, C. K. Dodge.
1857. **V. officinalis** L. Common Speedwell. Dry hills. Ann Arbor, Winch. Cat.; Alma, Ann Arbor, C. A. Davis. Scarce. S.
*1858. **V. peregrina** L. Neckweed. Purslane Speedwell. Moist grounds everywhere. Common. Th.
*1859. **V. scutellata** L. Marsh Speedwell. Swamps. Frequent. Th.
*1860. **V. serpyllifolia** L. Thyme-leaved Speedwell. Open grounds. Frequent. Th.

LEPTANDRA Nutt.

*1861. **L. Virginica** (L.) Nutt. Culver's-root. *Veronica Virginica* L. Open woods. Common. Th.

BUCHNERA L.

1862. **B. Americana** L. Blue-Hearts. Moist sandy ground. S. W. Wright Cat.; Flint; Macomb Co., Coleman Cat. Infrequent. C. & S.

DASYSTOMA Raf. GERARDIA L., in part.

*1863. **D. flava** (L.) Wood. Downy False Fox glove. *Gerardia flava* L. Open woods. Frequent. C. & S.
1864. **D. laevigata** Raf. Entire-leaved False Fox glove. *Gerardia laevigata* Raf. Coleman's Cat.; Flint; Dr. Lyons. S.
*1865. **D. Pedicularia** (L.) Benth. Fern-leaved Fox glove. *Gerardia pedicularia* L. Woods. Frequent. L. P.
*1866. **D. Virginica** (L.) Britton. Smooth False Fox glove. *Gerardia quercifolia* Pursh. Oak woods. Frequent. C. & S.

GERARDIA L.

1867. **G. aspera** Dougl. Rough Purple Gerardia. "Plains and prairies, Mich.," A. Gray's Manual.
1868. **G. auriculata** Michx. Auricled Gerardia. Wright Cat. Rare. S.
*1869. **G. paupercula** (A. Gray) Britton. Small-flowered Gerardia. *G. purpurea*

paupercula A. Gray. Oakland Co.; shore of Park Lake, Clinton Co., C. F. Wheeler; St. Clair Co., W. S. Cooper.

1870. **G. purpurea** L. Purple Gerardia. Moist, sandy ground, near the Great Lakes; Port Huron, C. K. Dodge; Orion, O. A. Farwell. L. P.

1871. **G. tenuifolia** Vahl. Slender Gerardia. Ann Arbor, Winch. Cat., Macomb Co. Rare.

*1872. **G. tenuifolia asperula** A. Gray. Lenawee Co., G. F. Comstock; Flint; Dr. Clark.

CASTILLEJA Mutis.

*1873. **C. acuminata** (Pursh.) Spreng. Lance-leaved Painted Cup. *C. pallida septentrionalis* A. Gray. Lake Superior, A. Gray; Keweenaw Co., O. A. Farwell. U. P.

*1874. **C. coccinea** (L.) Spreng. Scarlet Painted Cup. Wet and dry grounds. Varies in color of bracts from scarlet to yellow. Common. Th.

EUPHRASIA L.

1875. **E. Canadensis** Townsend. Reported along Lake Superior as *E. officinalis Tartarica.*

PEDICULARIS L.

*1876. **P. Canadensis** L. Common Lousewort. Wood Betony. Moist banks and woodlands. Common. Th.

*1877. **P. lanceolata** Michx. Swamp Lousewort. Swamps. Frequent. Th.

RHINANTHUS L.

1878. **R. Crista-Galli** L. Yellow-Rattle. "Lake Superior," A. Gray. U. P.

MELAMPYRUM L.

1879. **M. latifolium** Muhl. Broad-leaved Cow-Wheat. Keweenaw Co., O. A. Farwell; Bay Co., G. M. Bradford.

*1880. **M. lineare** Lam. Narrow-leaved Cow-Wheat. *M. Americanum* Michx. Sandy woods. A form with broad, spear-shaped leaves is frequent. Common. Th.

LENTIBULARIACEÆ Lindl. Bladderwort Family.

UTRICULARIA L.

*1881. **U. cornuta** Michx. Horned Bladderwort. Shore of Pine Lake, Ingham Co.; Point Sable, Mason Co., C. E. St. John; Petoskey; Cheboygan Co., B. & K.; Keweenaw Co., O. A. Farwell; Alma, Ann Arbor, C. A. Davis. Th.

*1882. **U. gibba** L. Humped Bladderwort. Old Mission, Grand Traverse Co., E. J. Hill; shore of Park Lake, Clinton Co., C. F. Wheeler; Lenawee Co., G. F. Comstock; Sturgis, F. P. Daniels. Rare. L. P.

1883. **U. intermedia** Hayne. Flat-leaved Bladderwort. Shallow water. Ann Arbor, Winch. Cat.; Macomb Co.; Grand Rapids, Sones; northward to Black Lake, Cheboygan Co., B. & K., and Keweenaw Co., O. A. Farwell; Alma. Infrequent. Th.

1884. **U. minor** L. Smaller Bladderwort. Shallow water. Detroit River, D. H. Campbell; Ann Arbor, Winch. Cat.; Hillsdale, University Herb; Hubbardston; Montcalm Co.; northward. Rare.

*1885. **U. purpurea** Walt. Purple Bladderwort. Hamlin Lake, H. T. Blodgett; Marshall, W. J. B.; Pine Lake, Ingham Co., Grass Lake, C. F. Wheeler; Ann Arbor, C. A. Davis.

*1886. **U. resupinata** B. D. Greene. Reversed Bladderwort. Very abundant on the east shore of Woodward Lake, in Ionia Co., E. F. Smith; shore of Pine Lake, Ingham Co., also reported from Whitings, Ind.; Alma, C. A. Davis.

*1887. **U. vulgaris** L. Greater Bladderwort. Slow streams. Throat of corolla-orange, veined with brown-purple. Frequent. Th.

PINGUICULA L.

1888. **P. vulgaris** L. Butterwort. Wet rocks. Whitney's Catalogue; Isle Royale, Dr. A. B. Lyons; Pictured Rocks, abundant, G. H. Hicks. U. P.

OROBANCHACEÆ Lindl. Broom-rape Family.

THALESIA Raf. APHYLLON A. Gray.

1889. **T. fasciculata** (Nutt.) Britton. Yellow Cancer-root. *Aphyllon fasciculatum* A. Gray. "Sandy ground, Lake Michigan," A. Gray; sand dunes, Frankfort.

1890. **T. uniflora** (L.) Britton. Pale Broom-rape. Cancer-root. *Aphyllon uniflorum* Torr. & Gray. Damp woodlands, S. Mich., Wright Cat.; Ann Arbor, G. D. Sones; Flint; Grayling, G. H. Hicks; Lake Superior, Winch. Cat. Infrequent. Th.

CONOPHOLIS Wallr.

1891. **C. Americana** (L. f.) Wallr. Squaw-root. Moist woods, Ann Arbor; S. W. H. S. Pepoon; S. Mich. Wright Cat.; Ionia Co.; Flint; Alma; Macomb Co., northward to Keweenaw Co., F. Infrequent.

LEPTAMNIUM Raf. EPIFAGUS Nutt.

1892. **L. Virginianum** (L.) Raf. Beech-drops. *Epifagus Virginiana* Nutt. Beech woods, Ionia Co.; Flint; Macomb Co. Very common in Grand-Saginaw Valley.

BIGNONIACEÆ Pers. Trumpet-creeper Family.

CATALPA Scop.

1893. **C. speciosa** Warder. Sturgis, F. P. Daniels. Probably introduced from the south.

ACANTHACEÆ J. St. Hil. Acanthus Family.

RUELLIA L.

1894. **R. cilosa** Pursh. Hairy Ruellia. Dry grounds. Wright Cat. Detroit, O. A. Farwell. S.

1895. **R. strepens** L. Smooth Ruellia. Dry soil. Wright Cat. S.

DIANTHERA L.

1896. **D. Americana** L. Dense-flowered Water-willow. In shallow water. Ann Arbor, Winch. Cat.; Put-in-Bay; S. Haven, L. H. Bailey. S.

PHRYMACEÆ Schauer. Lopseed Family.

PHRYMA L.

*1897. **P. Leptostachya** L. Lopseed. Moist woods. Pt. au Chene, Winch. Cat.; Flint; Macomb Co.; Wright Cat.; Ionia Co.; Turin, Marquette Co., B. Barlow; Manistee, F. P. Daniels; Alma, Ann Arbor, C. A. Davis. C. & S.

PLANTAGINACEÆ Lindl. Plantain Family.

PLANTAGO L.

1898. **P. aristata** Michx. Large-bracted Plantain. Introduced with clover seed. Huron Co., C. A. Davis.

*1899. **P. cordata** Lam. Heart-leaved Plantain. Borders of streams. S. Mich., Wright Cat.; Tuscola Co., Winch. Cat.; Ionia Co.; Clinton Co.; Flint; Macomb Co. Frequent. C. & S.

*1900. **P. lanceolata** L. Ripplegrass. Ribgrass. English Plantain. Buck-horn. Meadows and fields. Too common. The seeds are often mixed with clover seed. Th.

*1901. **P. major** L. Common Plantain. Waysides and about dwellings everywhere. Common. Th.

1902. **P. media** L. Hoary Plantain. Bay City, G. M. Bradford.
1903. **P. Purshii** R. & S. S. W., H. S. Pepoon.
*1904. **P. Rugelii** Dec. Rugel's Plantain. Waste places. Often confounded with *P. major*. Frequent.
1905. **P. Virginica** L. White Dwarf Plantain. Algonac, C. K. Dodge.

RUBIACEÆ B. Juss. Madder Family.

HOUSTONIA L.

1906. **H. coerulea** L. Bluets. Innocence. Keweenaw Point, Dr. Robbins.
1907. **H. ciliolata** Torr. Fringed Houstonia. *H. purpurea ciliolata* A. Gray. Dr. Wright. Three Rivers, C. F. Wheeler; Ypsilanti, O. A. Farwell.
1908. **H. longifolia** Gaertn. Long-leaved Houstonia. *H. purpurea longifolia* A. Gray. Hastings, L. H. Bailey; Ionia, common; Clare Co., abundant to L. Sup., Can. Cat.; Vestaburg, Grass Lake, Ann Arbor, C. A. Davis. Th.

CEPHALANTHUS L.

*1909. **C. occidentalis** L. Button-bush. Swamps and flooded river bottoms. Common, Cheboygan Co., B. & K. Very common southward. L. P.

MITCHELLA L.

*1910. **M. repens** L. Partridge-berry. Prefers beech and maple, hemlock, or pine woods, and is seldom found under oaks. Common. Th.

GALIUM L.

*1911. **G. Aparine** L. Cleavers. Goose grass. Common. Th.
*1912. **G. asprellum** Michx. Rough Bedstraw. Frequent. Th.
*1913. **G. boreale** L. Northern Bedstraw. Very common. Th.
*1914. **G. circaezans** Michx. Wild Liquorice. Dry woods, Cheboygan Co., B. & K. Frequent southward. L. P.
1915. **G. Claytoni** Michx. Clayton's Bedstraw. Muskegon, C. D. McLouth; St. Clair Co., C. K. Dodge; Keweenaw Co., O. A. Farwell.
*1916. **G. concinnum** Torr. & Gray. Shining Bedstraw. Flint; Ann Arbor, All. Cat.; Grand Ledge. Rare. C. &. S.
*1917. **G. lanceolatum** Torr. Wild Liquorice. Riverdale, Gratiot Co.; Hubbardston; Flint; Macomb Co.; and S. Mich. Wr. Cat.; also L. Superior, Whitney; Alma, Ann Arbor, C. A. Davis. Th.
*1918. **G. latifolium** Michx. Purple Bedstraw. Infrequent. C. & S.
1919. **G. Mollugo** L. Wild Madder. Bay City, G. M. Bradford.
*1920. **G. pilosum** Ait. Hairy Bedstraw. Frequent as far north as Grand-Saginaw Valley. C. & S.
1921. **G. tinctorium** L. Marshes, Bay Co., G. M. Bradford; Keweenaw Co., O. A. Farwell.
*1922. **G. trifidum** L. Small Bedstraw. Bogs. Common. Th.
*1923. **G. triflorum** Michx. Sweet-scented Bedstraw. Frequent in C. & S., and very abundant northward. Th.

CAPRIFOLIACEÆ Vent. Honeysuckle Family.

SAMBUCUS L.

*1924. **S. Canadensis** L. Common Elder. Follows settlements. Common.
*1925. **S. pubens** Michx. Red-berried Elder. *S. racemosa* HK. More northern in its range than the last. Variety with dissected leaves seen in Clare Co. Th.

VIBURNUM L.

*1926. **V. acerifolium** L. Arrow-wood. Dockmackie. Frequent in Cheboygan Co., B. & K.; Alma, Ann Arbor, C. A. Davis.
1927. **V. alnifolium** Marsh. Hobble-bush. American Wayfaring-tree. *V. lantanoides* Michx. Whitney found none of the sp. abundant in the U. P.

*1928. **V. cassinoides** L. Withe-rod. Macomb Co.; Hubbardston; Stanton; Riverdale; Gratiot Co.; Houghton Lake; to L. Superior. Commonest sp. about Black Lake, B. & K. Frequent. Th.

1929. **V. dentatum** L. Arrow-wood. Sarnia, Ont., C. K. Dodge.

*1930. **V. Lentago** L. Sweet Viburnum. Sheep-berry. Frequent. C. & S., and northward. Cheboygan Co.; Alma. Th.

*1931. **V. Opulus** L. Cranberry-tree. Swamps and borders of streams. Common. Th.

1932. **V. pauciflorum** Pylaie. Few-flowered Cranberry-tree. Dr. Lyons. Gray's Manual. Rare. U. P.

1933. **V. prunifolium** L. Black Haw. Dr. Lyons; Kalamazoo, Tuthill; Barron Lake, Cass Co., C. F. Wheeler; Detroit, O. A. Farwell. Rare. S.

*1934. **V. pubescens** (Ait.) Pursh. Downey Arrow-root. Common. Th.

TRIOSTEUM L.

1934a. **T. aurantiacum** Bicknell. Birmingham, O. A. Farwell.

*1935. **T. perfoliatum** L. Horse-Gentian. Clinton Co.; Ionia Co.; Flint; Macomb Co.; Black River, Cheboygan Co., one plant seen by B. & K. Frequent southward. L. P.

LINNAEA L.

*1936. **L. Americana** Forbes. Twin-flower. *L. borealis Michx.*, not L. Very abundant in N. & U. P.; south to Grand River Valley, and S. E. to Macomb Co.; Port Huron, C. K. Dodge. C. N. & U. P.

SYMPHORICARPOS Juss.

1937. **S. occidentalis** Hook. Wolfberry. Ft. Gratiot, Austin; N. Mich., A. Gray; Port Huron, C. K. Dodge. Rare. C. N. & U. P.

1938. **S. pauciflorus** (Robbins) Britton. Low Snowberry. *S. racemosus pauciflorus* Robbins. Harbor Springs; Keweenaw Co., Dr. Robbins; Hubbardston, C. F. Wheeler.

1939. **S. racemosus** Michx. Snowberry. Along the Great Lakes, Saginaw Bay and Alpena Co., Winchell; L. Sup., Can. Cat. C. N. & U. P.

1940. **S. Symphoricarpos** (L.) MacM. Coral-berry. Indian Currant. *S. vulgaris* Michx. Keweenaw Co., O. A. Farwell.

LONICERA L.

1941. **L. Caprifolium** L. Perfoliate Honeysuckle. *L. grata* Aiton. Dr. Lyons. Gray's Manual.

*1942. **L. ciliata** Muhl. Fly-Honeysuckle. Frequent. Th.

1943. **L. coerulea** L. Mountain Fly-Honeysuckle. Dr. Lyons. Keweenaw Co., O. A. Farwell; Clifton, F. E. Wood, in University Herb.

*1944. **L. dioica** L. Glaucous Honeysuckle. *L. glauca* Hill. Ionia; Montcalm Co.; Ann Arbor, All. Cat. Swamps or dry soil. Common. Th.

*1945. **L. glaucescens** Ryd. Douglas' Honeysuckle. Keweenaw Co., O. A. Farwell; Bay Co., G. M. Bradford.

1946. **L. hirsuta** Eaton. Hairy Honeysuckle. Alma, C. A. Davis. Frequent northward. C. N. & U. P.

1947. **L. involucrata** (Richards.) Banks. Mainland and Isle Royale; Keweenaw Co., O. A. Farwell; Ann Arbor, C. A. Davis.

*1948. **L. Japonica** Thunb. Japanese Honeysuckle. Lansing; Ypsilanti, O. A. Farwell.

*1949. **L. oblongifolia** (Goldie) Hook. Swamp F. Howell Junction, C. F. Wheeler; Macomb Co.; Stanton; Edmore; Isabella Co., Roscommon; Alma. More frequent in N. and U. P. Rare in S. Th.

1950. **L. Sullivantii** A. Gray. Keweenaw Co., O. A. Farwell.

*1951. **L. Tatarica** L. Tartarian Honeysuckle. Escaped from cultivation, Lansing; Detroit; Port Huron; Ypsilanti, O. A. Farwell.

1952. **L. Xylosteum** L. Escaped from cultivation. Detroit. O. A. Farwell.

DIERVILLA Moench.

*1953. **D. Diervilla** (L.) MacM. Bush Honeysuckle. *D. trifida* Moench. Rocky woods and bluffs. Common. Th.

VALERIANACEÆ Batsch. Valerian Family.

VALERIANA L.

1954. **V. edulis** Nutt. Tobacco-root. Ann Arbor and Macomb Co. Rare. S. E.
*1955. **V. uliginosa** (T. & G.) Rydb. Swamp Valerian. *V. sylvatica* Beck, not Banks. Sphagnous swamps, local. Th.

VALERIANELLA Poll.

1956. **V. chenopodifolia** (Pursh) DC. Goose-foot Corn Salad. Flint; Ionia; Lyons. Not common. C. & S.
1957. **V. radiata** (L.) Dufr. Beaked Corn Salad. Mich. A. Gray; Macomb Co., Cooley.

DIPSACACEÆ Lindl. Teasel Family.

DIPSACUS L.

*1958. **D. sylvestris** Huds. Wild Teasel. Fields and roadsides. Gratiot Co.; Ann Arbor; Detroit; Flint; Alma; Addison. Infrequent. C. & S.

CUCURBITACEÆ B. Juss. Gourd Family.

MICRAMPELIS Raf. ECHINOCYSTIS T. & G.

1959. **M. lobata** (Michx.) Greene. Wild Balsam Apple. *Echinocystis lobata* T. & G. Common in low woods along streams; Alma; Ionia; South Haven; Macomb Co.; Ann Arbor; Manistee, F. P. Daniels; Bay Co., G. M. Bradford.

SICYOS L.

1960. **S. angulatus** L. One-seeded Star Cucumber. Ypsilanti, O. A. Farwell; Port Huron, C. K. Dodge; Alma, Ann Arbor, C. A. Davis; Algonac, W. S. Cooper.

CAMPANULACEÆ Juss. Bellflower Family.

CAMPANULA L.

*1961. **C. Americana** L. Tall Bellflower. Moist woodlands. Ionia Co.; Flint; Macomb Co.; Ann Arbor, Winch. Cat.; Alma, C. A. Davis. Infrequent. C. & S.
*1962. **C. aparinoides** Pursh. Marsh Bellflower. Wet grassy grounds. Common. Th.
1963. **C. rapunculoides** L. Creeping European Bellflower. Ann Arbor, A. J. Pieters; Ypsilanti, O. A. Farwell; Bay Co., G. M. Bradford.
*1964. **C. rotundifolia** L. Harebell. Sandy banks and lake shores. Common. Th.
1965. **C. rotundifolia Langsdorfiana** (A. DC.) Britton. *C. rotundifolia alpina* Tuckerman. Keweenaw Co., O. A. Farwell; shore of Black Lake, Cheboygan Co., B. & K.; Petoskey, C. F. Wheeler.
1966. **C. rotundifolia velutina** DC. Sand hills of Burt Lake, E. J. Hill.

SPECULARIA Heist.

*1967. **S. perfoliata** (L.) A. DC. Venus' Looking-glass. Gravelly fields. Clinton Co.; S. E., University Herb.; Niles, I. N. Mitchell; Detroit, A. B. Lyons; Ann Arbor. Scarce. C. & S.

LOBELIA L.

*1968. **L. cardinalis** L. Cardinal-flower. River banks. Flowers rarely rose-color or even white. Common. Th.
1969. **L. Dortmanna** L. Water Lobelia. Isle Royale, Dr. A. B. Lyons. U. P.
*1970. **L. Kalmii** L. Kalm's Lobelia. Wet banks and rocks along shores. Ionia Co.; Ann Arbor; Oakland Co.; Petoskey; to L. Superior. Frequent. Th.

*1971. **L. inflata** L. Indian Tobacco. Pastures. Montcalm Co.; Flint; Macomb Co.; Oakland Co.; Put-in-Bay. Scarce. Th.
*1972. **L. spicata** Lam. Pale Spiked Lobelia. Oak openings. Frequent. Th.
1973. **L. spicata hirtella** A. Gray. Keweenaw Co., Detroit, O. A. Farwell.
*1974. **L. syphilitica** L. Great Lobelia. Low grounds. Flowers vary to white. Common. L. P.

CICHORIACEÆ Reichenb. Chicory Family.

CICHORIUM L.

*1975. **C. Intybus** L. Chicory. Ionia Co.; Flint; Detroit; Keweenaw Co., O. A. Farwell; Alma; Ann Arbor. Frequent. Th.

LAPSANA L.

*1976. **L. communis** L. Nipplewort. Campus of the Agricultural College.

ADOPOGON Neck. KRIGIA Schreb.

1977. **A. Carolinianum** (Walt.) Britton. Carolina Dwarf Dandelion. *Krigia Virginica* Willd. Barrens N. part of Clare Co.; Walton; Elk Rapids; Bay City. Rare.
*1978. **A. Virginicum** (L.) Kuntze. *Krigia amplexicaulis* Nutt. Moist hillsides. Frequent. L. P.

LEONTODON L.

1979. **L. autumnalis** L. Fall Dandelion. Well established at Hubbardston, C. F. Wheeler; Alma, C. A. Davis.

PICRIS L.

1980. **P. hieracioides** L. Hawkweed Picris. Detroit, O. A. Farwell.

TRAGOPOGON L.

1981. **T. porrifolius** L. Oyster Plant. Escaped from cultivation. St. Clair Co.; Alma; Keweenaw Co., O. A. Farwell.
*1982. **T. pratensis** L. Goat's Beard. Spreading. Hubbardston, C. F. Wheeler; Kalamazoo, Tuthill; Keweenaw Co., Ypsilanti, O. A. Farwell.

TARAXACUM Haller.

1983. **T. erythrospermum** Andrz. Red-seeded Dandelion. Detroit, O. A. Farwell.
*1984. **T. Taraxicum** (L.) Karst. Dandelion. *T. officinale* Weber. Fields everywhere.

SONCHUS L.

*1985. **S. arvensis** L. Field Sow-Thistle. Grand Rapids, Coleman's Cat.; Detroit, O. A. Farwell; Tuscola Co., C. A. Davis; Bay Co., G. M. Bradford.
*1986. **S. asper** (L.) Hill. Spiney-leaved Sow-Thistle. Alma, C. A. Davis. Waste places. Frequent.
*1987. **S. oleraceus** L. Common Sow-Thistle. Waste places. Macomb Co.; Montcalm Co.; Alma, C. A. Davis; northward to L. Superior. Frequent.

LACTUCA L.

*1988. **L. Canadensis** L. Wild Lettuce. Rich soil. Frequent. Th.
*1989. **L. Floridana** (L.) Gaertn. Florida Lettuce. Wayne Co., O. A. Farwell; Agricultural College, W. J. B.
*1990. **L. hirsuta** Muhl. Hairy Wood-lettuce. Infrequent. Hubbardston; Macomb Co., Cooley; Alma, Chelsea, C. A. Davis.
1991. **L. pulchella** (Purhs.) DC. Large-flowered Blue lettuce. "Upper Michigan," Prof. T. C. Porter; Caribou I.; Lake Huron, Dr. Todd; Detroit, O. A. Farwell; Keweenaw Co., O. A. Farwell. N. & U. P.
1992. **L. sagittifolia** Ell. Arrow-leaved Lettuce. *L. integrifolia* Bigel. Petoskey, C. F. Wheeler; Belle Isle, O. A. Farwell.

*1994. **L. spicata** (Lam.) Hitchc. Tall Blue Lettuce. *L. leucophaea* A. Gray. Frequent. Th.

1995. **L. spicata integrifolia** (T. &. G.) Britton. *L. leucophaea integrifolia* T. & G. Belle Isle, Detroit, C. F. Wheeler.

*1995a. **L. virosa** L. Prickly Lettuce. Usually given the name *L. Scariola* L., which has pinnatifid leaves. Common.

CREPIS L.

*1996. **C. tectorum** L. Narrow-leaved Hawksbeard. Introduced.

HIERACIUM L.

*1997. **H. Canadense** Michx. Canada Hawkweed. Woods. Frequent. Th.

*1998. **H. Gronovii** L. Hairy Hawkweed. Dry soil. Ionia Co.; Flint; Macomb Co.; S. W. Mich., Wright Cat.; Alma. Infrequent. C. & S.

1999. **H. longipilum** Torr. Long-bearded Hawkweed. Fields. Macomb Co.; Traverse City, S. W., Wright Cat.; Grand Rapids, Coleman; Algonac, W. S. Cooper; Tuscola Co., C. A. Davis. Scarce. L. P.

2000. **H. Marianum** Willd. Maryland Hawkweed. Shore of Grand Traverse Bay, C. F. Wheeler.

*2001. **H. paniculatum** L. Panicled Hawkweed. Macomb Co.; S. Mich., Wright Cat. Scarce. C. & S.

2002. **H. Pilosella** L. Mouse-ear Hawkweed. Introduced at Benzonia, G. A. Clark.

*2003. **H. scabrum** Michx. Rough Hawkweed. Woods; Alma. Common. Th.

*2004. **H. umbellatum** L. Narrow-leaved Hawkweed. Marquette, C. F. Wheeler.

*2005. **H. venosum** L. Rattlesnake-weed. Dry soil in pine woods, or on oak land. Abundant at Point aux Pins, at the entrance to Lake Superior, Macoun, Can. Cat.; Port Austin, Ann Arbor, C. A. Davis. Frequent.

NABALUS Cass. PRENANTHES Vaill.

*2006. **N. albus** (L.) Hook. Rattlesnake-root. *Prenanthes alba* L. Woods. Common. Th.

*2007. **N. altissimus** (L.) Hook. Tall White Lettuce. *Prenanthes altissima* L. Rich woods. Frequent. Th.

*2008. **N. racemosus** (Michx.) DC. Glaucous White-Lettuce. *Prenanthes racemosa* Michx. Lenawee Co., G. F. Comstock; Flint; Macomb Co.; shore of L. Mich., near Sitting Rabbit, Winch. Cat.; S. W., Wright Cat.; Cheboygan Co., Beardslee, Keweenaw Co., Farwell; Kawkawlin, Ann Arbor, Grass Lake, Bay Port, C. A. Davis. Rare. Th.

2009. **N. trifoliatus** Cass. Tall Rattlesnake-root. Washington, Dr. D. Cooley.

AMBROSIACEÆ Reichenb. Ragweed Family.

IVA L.

2010. **I. xanthiifolia** (Fresen.) Nutt. Marsh Elder. Highwater-shrub. Keweenaw Co., O. A. Farwell. The farthest station east known.

AMBROSIA L.

*2011. **A. artemisiaefolia** L. Roman Wormwood. Ragweed. Hog-weed. Bitterweed. Roadsides. Introduced from the west. Abundant. Th.

2012. **A. psilostachya** DC. Manistee, F. P. Daniels; shore of Lake Huron, C. K. Dodge; Keweenaw Co., O. A. Farwell.

*2013. **A. trifida** L. Great Ragweed. Common. Low grounds along Grand and Maple Rivers; northward to Keweenaw Co., O. A. Farwell.

2014. **A. trifida integrifolia** T. & G. Low land. Keweenaw Co., O. A. Farwell. Th.

XANTHIUM L.

*2015. **X. Canadense** Mill. Common in waste places and along river banks.

2015a. **X. commune** Britton. Britton and Brown's Flora.

2016. **X. echinatum** Murr. *X. Canadense echinatum* A. Gray. Shores of Great Lakes.

2017. **X. glabratum** (DC.) Britton. Broad Cocklebur. *X. strumarium* of authors, not of Linnaeus. Detroit, O. A. Farwell.

X. Pennsylvanicum Wallr. Britton and Brown's Flora.

2018. **X. spinosum** L. Spiny Clotbur. Detroit, O. A. Farwell.

COMPOSITÆ Adans. Thistle Family.

VERNONIA Schreb.

2019. **V. Drummondii** Shuttlw. Drummond's Iron-weed. *V. altissima grandiflora* A. Gray. Frequent in the Grand River Valley; Bay Co., G. M. Bradford.

*2020. **V. fasciculata** Michx. River banks; Ionia Co.; Macomb Co.; Detroit, O. A. Farwell; Monroe Co.; Ann Arbor; South Haven, L. H. Bailey. Frequent. C. & S.

2021. **V. glauca** (L.) Britton. Broad-leaved Iron-weed. Three Rivers, C. F. Wheeler; Detroit, O. A. Farwell.

2022. **V. interior** Small. Detroit, O. A. Farwell.

2023. **V. maxima** Small. Tall Iron-weed. *V. gigantea* (Walt.) Britton. Detroit, O. A. Farwell; S. W., H. S. Pepoon. Frequent eastward.

2024. **V. Noveboracensis** (L.) Willd. Manistee, F. P. Daniels.

EUPATORIUM L.

*2025. **E. ageratoides** L. f. White Snake-root. Woods. Frequent. Th.

2026. **E. coelestinum** L. Mist-flower. "Rich soil, Mich." A. Gray's Manual.

2027. **E. maculatum** L. Spotted Joe-Pye Weed. Keweenaw Co., Rochester, O. A. Farwell; near Algonac, C. K. Dodge.

*2028. **E. perfoliatum** L. Thoroughwort. Boneset. Low grounds. Common. Th.

*2029. **E. purpureum** L. Joe-Pye Weed. Keweenaw Co., Rochester, O. A. Farwell. Th.

*2030. **E. purpureum falcatum** (Michx.) Britton. Keweenaw Co., O. A. Farwell.

2031. **E. sessilifolium** L. Upland Boneset. Copses. Macomb Co.; S. Mich., Winch. Cat. S.

2032. **E. trifoliatum** L. Rochester, O. A. Farwell.

WILLUGBAEYA Neck. MIKANIA Willd.

2033. **W. scandens** (L.) Kuntze. Climbing Hemp-weed. *Mikania scandens* Willd. Banks of Muskegon River for a distance of eleven miles above Black Lake, C. D. McLouth.

KUHNIA L.

2034. **K. eupatorioides** L. Ionia Co.; S. Mich., Winch. Cat. Infrequent. C. & S.

LACINARIA Hill. LIATRIS Schreb.

2035. **L. cylindracea** (Michx.) Kuntze. Cylindric Blazing Star. *Liatris cylindracea* Michx. Sterile open places; Ionia Co.; Macomb Co.; Lenawee Co., G. F. Comstock; Tuscola Co.; Grand Rapids to Keweenaw Co., O. A. Farwell. Not common. Th.

*2036. **L. scariosa** (L.) Hill. Large Button Snake-root. *Liatris scariosa* Willd. Rarely the flowers vary to white. Dry soil. Ann Arbor; northward to Keweenaw Co., O. A. Farwell. Frequent.

2036a. **L. scariosa corymbulosa** Sheldon. Orion, O. A. Farwell.

2037. **L. scariosa squarrulosa** (Michx.) Small. Orion, O. A. Farwell.

*2038. **L. spicata** (L.) Kuntze. Dense Button-Snakeroot. *Liatris spicata* Willd. Low grounds. Ionia Co.; Macomb Co.; near Port Huron, C. K. Dodge; Bay Co., G. M. Bradford; Kalamazoo, Tuthill. Infrequent.

GRINDELIA Willd.

2039. **G. squarrosa** (Pursh.) Dunal. Broad-leaved Gum-plant. Escanaba, R. E. Merrell; near Lapeer, Mrs. M. Owen; Marquette, O. A. Farwell. Occasionally introduced in grass seeds from the west.

SOLIDAGO L.

2040. **S. alpestris** Wald. & Kit. Alpine Golden-rod. *S. Virgaurca alpina* Bigel. Gray's Manual, 6th edition. U. P.

*2041. **S. bicolor** L. White Golden-rod. Rochester, W. A. Brotherton; Lansing, W. J. B.; Detroit, O. A. Farwell.

*2042. **S. cæsia** L. Blue-stemmed Golden-rod. Rich woods. Frequent. L. P.

*2043. **S. caesia axillaris** (Pursh.) A. Gray. Frequent near Lansing. L. P.

*2044. **S. Canadensis** L. Canada Golden-rod. Fields. Our most variable and common golden-rod. Th.

*2045. **S. Canadensis procera** (Ait.) T. & G. Common.

2046. **S. Canadensis scabriuscula** T. C. Porter. Keweenaw Co., Detroit, O. A. Farwell.

2047. **S. erecta** Pursh. Slender Golden-rod. *S. speciosa angustata* T. & G. Indian River, Cheboygan Co., C. F. Wheeler.

*2048. **S. flexicaulis** L. Zig-Zag Golden-rod. *S. latifolia* L. Moist woods. Frequent. Th.

*2049. **S. hispida** Muhl. Hairy Golden-rod. *S. bicolor concolor* Torr & Gray. Dry places. Ionia Co.; Flint; Macomb Co.; Alma, Ann Arbor, C. A. Davis; and northward.

2050. **S. Houghtonii** Torr. & Gray. Houghton's Golden-rod. "N. shore of Lake Michigan," Gray's Man.; Drummond's Is., Winch. Cat.

*2051. **S. juncea** Ait. Early Golden-rod. Meadows and fields, variable. Frequent. Th.

2052. **S. juncea scabrella** (T. & G.) A. Gray. Orion, Detroit, O. A. Farwell.

2053. **S. macrophylla** Pursh. Large-leaved Golden-rod. "Shore of Lake Superior and northward," A. Gray's Manual, 6th edition. U. P.

2054. **S. neglecta** Torr. & Gray. Swamp Golden-rod. Swamps. Ionia Co., Orion, O. A. Farwell, and northward. Infrequent.

*2055. **S. nemoralis** Ait. Gray Golden-rod. Dry sandy soil. Common. Th.

2056. **S. Ohioensis** Riddell. Ohio Golden-rod. Moist meadows. Ionia Co.; Flint; Bay Co., G. M. Bradford; Macomb Co. to Grand Detour below Sugar Island, Prof. T. C. Porter. Infrequent.

*2057. **S. patula** Muhl. Rough-leaved Golden-rod. Borders of swamps. Common. C. & S.

2058. **S. Purshii** T. C. Porter. River bank Golden-rod. *S. humilis* Pursh. Frequent at Petoskey; Cheboygan Co., Beardslee; Keweenaw Co., O. A. Farwell.

*2059. **S. rigida** L. Stiff Golden-rod. Dry, sandy ground. Ann Anbor; Ionia Co.; Flint; Manistee, F. P. Daniels, and northward. Infrequent.

*2060. **S. Riddellii** Frank. Riddell's Golden-rod. Swamps, Ionia Co.; Macomb Co.; Detroit, O. A. Farwell; Ann Arbor, and southward. Rare.

*2061. **S. rugosa** Mill. Wrinkle-leaved Golden-rod. Borders of fields, northward to Thunder Bay, Macoun. Common. Th.

*2062. **S. serotina** Ait. Late Golden-rod. Shady places. Frequent. Th.

2063. **S. serotina gigantea** (Ait.) A. Gray. Borders of woods. Frequent. Th.

*2064. **S. speciosa** Nutt. Showy Golden-rod. Margin of woods on light soil. Ionia Co.; Flint; Macomb Co.; Ann Arbor; and northward to Keweenaw Co., O. A. Farwell. Infrequent.

*2065. **S. uliginosa** Nutt. Bog Golden-rod. Swamps. Ann Arbor, Tuscola Co., Alpena Co., C. A. Davis. Infrequent. Th.

*2066. **S. ulmifolia** Muhl. Elm-leaved Golden-rod. Borders of woods. Ionia Co.; Flint; Alma; S. W., Winch. Cat. Infrequent. L. P.

2067. **S. Virgaurea Gillmani** (A. Gray) T. C. Porter. *S. humilis Gillmani* A. Gray. Shores of Lake Mich., from New Buffalo to Mackinaw City, C. F. Wheeler.

EUTHAMIA Nutt.

2068. **E. Caroliniana** (L.) Greene. Slender Fragrant Golden-rod. *Solidago tenuifolia* Pursh. Grand Rapids, Miss E. J. Cole.

*2069. **E. graminifolia** (L.) Nutt. Bushy Golden-rod. *Solidago lanceolata* L. Moist soil. Alma, Ann Arbor, C. A. Davis. Common. Th.

ASTER L.

2069a. **A. amethystinus** Nutt. Detroit, O. A. Farwell.

*2070. **A. azureus** Lindl. Sky-blue Aster. Sterile soil. Flint; Ann Arbor; Ionia Co.; Bay Co., G. M. Bradford; Macomb Co. Infrequent. C. & S.

*2071. **A. cordifolius** L. Blue Wood Aster. *A. cordifolius polycephalus* T. C. Porter. Lansing, C. F. Wheeler; Bay Co., G. M. Bradford. Woods. Common. Th.

2072. **A. divaricatus** L. White Wood Aster. *A. corymbosus* Ait. Woods. Bay Co., G. M. Bradford. Infrequent. Th.

2073. **A. dumosus** L. Bushy Aster. Thickets. Macomb Co.; Petoskey, Dr. D. Clark; Hubbardston, not common, C. F. Wheeler; Cheboygan Co., Beardslee; Orion, O. A. Farwell; Alma, C. A. Davis. L. P.

2074. **A. ericoides** L. White Heath Aster. Petoskey, C. F. Wheeler; Bay Co., G. M. Bradford; shores of the Great Lakes.

2075. **A. ericoides pilosus** (Willd.) T. C. Porter. *A. ericoides villosus* T. & G. Marine City, C. K. Dodge.

2075a. **A. ericoides platyphyllus** T. & G. Detroit, O. A. Farwell.

2076. **A. exiguus** (Fernald) Rydb. Detroit, O. A. Farwell.

2078. **A. Faxoni** Porter. Faxon's Aster. *A. polyphyllus* Willd. Shores of the Great Lakes.

2079. **A. hirsuticaulis** Lindl. Hairy-stemmed Aster. *A. lateriflorus hirsutiusculis* T. C. Porter. Belle Isle, O. A. Farwell.

*2080. **A. junceus** Ait. Rush Aster. Tamarack swamps. Ionia Co.; Flint; Macomb Co.; Alma, Ann Arbor; northward. Frequent. Th.

*2081. **A. laevis** L. Smooth Aster. Border of oak woods. Cheboygan Co., B. & K.; Alma; Ann Arbor. Common southward. L. P.

2082. **A. laevis amplifolius** Porter. Detroit, Birmingham, O. A. Farwell.

2083. **A. lateriflorus** (L.) Britton. Starved Aster. *A. diffusus* Ait. Fields. Exceedingly variable. Common. Th.

2084. **A. lateriflorus glomerellus** (T. & G.) Burgess. Detroit, O. A. Farwell.

2085. **A. lateriflorus horizontalis** (Desf.) Burgess. Belle Isle, O. A. Farwell.

2085a. **A. lateriflorus pendulus** (Ait.) Burgess. Belle Isle, O. A. Farwell.

2086. **A. lateriflorus thyrsoideus** (A. Gray) Sheldon. *A. diffusus thyrsoides* A. Gray. Washington, Macomb Co., Dr. D. Cooley; Belle Isle. O. A. Farwell; and northward.

2087. **A. Lindleyanus** Torr. & Gray. Lindley's Aster. "Labrador to Lake Superior." A. Gray's Manual; Mackinaw City, C. F. Wheeler; Keweenaw Co., O. A. Farwell. Not common. N. & U. P.

2088. **A. longifolius** Lam. Long-leaved Aster. Escanaba, C. F. Wheeler.

2088a. **A. Lowrieanus** Porter. Lowrie's Aster. Lakeville, Brotherton and Farwell.

*2089. **A. macrophyllus** L. Large-leaved Aster. Woods. Common. Th.

*2090. **A. macrophyllus excelsior** Burgess. Keweenaw Co., O. A. Farwell.

2091. **A. multiflorus** Ait. Dense-flowered Aster. Sandy soil. Frequent. C. & S.

2092. **A. nemoralis** Ait. Common. Th.

*2093. **A. Novae-Angliae** L. New England Aster. Moist grounds. Frequent. Th.

2094. **A. Novi-Belgii** L. New York Aster. Detroit, O. A. Farwell.

2095. **A. oblongifolius** Nutt. Detroit. O. A. Farwell.

*2096. **A. paniculatus** Lam. Tall White Aster. Shady banks. Frequent. Th.

2098. **A. paniculatus bellidiflorus** (Willd.) Burgess. Belle Isle, C. F. Wheeler; Lapeer, Mrs. M. Owen.

2099. **A. paniculatus simplex** (Willd.) Burgess. Detroit, O. A. Farwell.

2100. **A. patens** Ait. Late Purple Aster. Dry grounds. Ann Arbor; Macomb Co. S. E.

2101. **A. prenanthoides** Muhl. Keweenaw Co., O. A. Farwell.

*2102. **A. ptarmicoides** (Nees.) Torr. and Gray. Macomb Co.; Clarkston, G. H. Hicks; Keweenaw Co., O. A. Farwell. Infrequent. Th.

*2103. **A. puniceus** L. Purple-stem Aster. Low grounds. Alma, Ann Arbor, C. A. Davis. Common. Th.

2103a. **A. puniceus firmus** (Nees) T. & G. Detroit, O. A. Farwell.

*2104. **A. puniceus lucidulus** Gray. Macomb Co. westward. Abundant.

*2105. **A. sagittifolius** Willd. Arrow-leaved Aster. Dry grounds. Frequent. Th.

2106. **A. sagittifolius urophyllus** (Lindl.) Burgess; Detroit, O. A. Farwell.

2107. **A. salicifolius** Lam. Willow Aster. Moist soil. Flint; Macomb Co., northward. Frequent.

2108. **A. sericeus** Vent. Silky Aster. S. Mich., Winch. Cat., Dr. D. Houghton; north to Keweenaw Co., O. A. Farwell. Rare.

2109. **A. Shortii** Hook. Short's Aster. Pewamo, C. F. Wheeler; Rochester, W. A. Brotherton. Rare.

*2110. **A. Tradescanti** L. Tradescant's Aster. Low grounds. Flint; Macomb Co. Frequent. L. P.

*2111. **A. undulatus** L. Wavy-leaf Aster. Dry copses. Flint; S. Mich.; Cheboygan Co., B. & K.; Ionia Co. Common. L. P.

*2112. **A. vimineus** Lam. Small White Aster. Low grounds. Cheboygan Co., Beardslee; Alma. Frequent southward. L. P.

2112a. **A. vimineus foliolosus** Ait. A. Gray. Detroit, O. A. Farwell.

BRACHYACTIS Ledeb.

2113. **B. angustus** (Lindl.) Britton. Rayless Aster. Port Huron, C. K. Dodge.

ERIGERON L.

2114. **E. acris** L. Blue Fleabane. Keweenaw Co., O. A. Farwell.

2115. **E. acris Droebachianus** (Retz) Willd. "Shores of Lake Superior," A. Gray; Dr. A. B. Lyons; Keweenaw Co., O. A. Farwell. U. P.

*2116. **E. annuus** (L.) Pers. Daisy Fleabane. Sweet Scabious. Fields. Common. Th.

2117. **E. asper** Nutt. Rough Erigeron. *E. glabellus* Nutt. Plains. Dr. A. B. Lyons. U. P.

2118. **E. hyssopifolius** Michx. Hyssop-leaved Erigeron. "Lake Superior, and northward," A. Gray; Dr. Lyons; Keweenaw Co., O. A. Farwell.

*2119. **E. Philadelphicus** L. Common Philadelphia Fleabane. Common. Th.

*2120. **E. pulchellus** Michx. Robin's Plantain. *E. bellidifolius* Muhl. Moist banks. Frequent. C. & S.

*2121. **E. ramosus** (Walt.) B. S. P. Daisy Fleabane. *E. strigosus* Muhl. Fields. Common. Th.

LEPTILON Raf. ERIGERON L., in part.

2122. **L. Canadense** (L.) Britton. Horse-weed. *Erigeron Canadensis* L. Waste grounds. Common. Th.

2122a. **L. divaricatum** (Michx.) Raf. Near Detroit, O. A. Farwell.

DOELLINGERIA Nees.

*2123. **D. umbellata** (Mill.) Nees. Tall Flat-top White Aster. *Aster umbellatus* Mill. Moist places. Th.

2124. **D. umbellata pubens** (A. Gray) Britton. *Aster umbellatus pubens* A. Gray. Upper Mich., A. Gray's Manual; Cheboygan Co., Beardslee; Keweenaw Co., O. A. Farwell.

ANTENNARIA Gaertn.

2125. **A. ambigens** (Greene) Fernald. Bay Co., G. M. Bradford, Detroit, O. A. Farwell.

*2127. **A. campestris** Rydberg. Prairie Cat's-foot. Grayling, C. F. Wheeler; Manistee, F. P. Daniels.

2128. **A. Canadensis** Greene. Shores of Grand Traverse Bay under pines, C. F. Wheeler; Mackinac Island, O. A. Farwell.

2129. **A. fallax** Greene. Detroit, O. A. Farwell.

*2130. **A. Farwellii** Greene. Lansing, C. F. Wheeler; Keweenaw Co., O. A. Farwell; Manistee, F. P. Daniels.

*2131. **A. neglecta** Greene. Field Cat's-foot. Lansing, Brighton, Stockbridge, Wheeler & Longyear; Detroit, Ypsilanti, O. A. Farwell; Manistee, F. P. Daniels.

2131a. **A. neodioica** Greene. Keweenaw Co., O. A. Farwell.

*2132. **A. neodioica attenuata** Fernald. Agricultural College, Brighton, Vestaburg, C. F. Wheeler; Detroit, Mackinac Island, O. A. Farwell.

2133. **A. occidentalis** Greene. Shores of Grand Traverse Bay, C. F. Wheeler.

*2134. **A. Parlinii ambigens** Fernald. Banks of Cedar River at Agricultural College, C. F. Wheeler; Manistee, F. P. Daniels; Port Huron, C. K. Dodge.

*2135. **A. plantaginifolia** (L.) Richards. Plantain-leaved Everlasting. Dry fields. Common. Th.

ANAPHALIS DC.

2136. **A. margaritacea** (L.) Benth. & Hook. Pearly Everlasting. Montcalm Co.; Grayling, G. H. Hicks; Petoskey, Mackinac, C. F. Wheeler; Bay Co., G. M. Bradford.

GNAPHALIUM L.

*2137. **G. decurrens** Ives. Everlasting. Winged Cudweed. Fields. Alma, Ann Arbor, C. A. Davis. Abundant in the pine region, and northward.
*2138. **G. obtusifolium** L. Sweet Balsam. *G. polycephalum* Michx. Common Everlasting. Fields. Common. Th.
2139. **G. purpureum** L. Purplish Cudweed. Macomb and St. Clair Counties, A. F. Foerste; Detroit, O. A. Farwell.
*2140. **G. uliginosum** L. Low Cudweed. Roadsides in clay soil, abundant. Th.

ADENOCAULON Hook.

2141. **A. bicolor** Hook. Moist ground. Ontonagon River. U. P.

INULA L.

*2142. **I. Helenium** L. Elecampane. Roadsides, escaped from gardens. Infrequent.

POLYMNIA L.

*2143. **P. Canadensis** L. Small-flowered Leaf-cup. Shaded river banks. Ionia Co.; Macomb Co.; Flint; S. Michigan. Infrequent. C. & S.
2144. **P. Canadensis radiata** A. Gray. Rochester, O. A. Farwell.
2145. **P. Uvedalia** L. Large-flowered Leaf-cup. S. Michigan, Wright's Cat. S.

SILPHIUM L.

2146. **S. integrifolium** Michx. Entire-leaved Rosin-weed. S. W., University Herb.; Kalamazoo, Tuthill.
2147. **S. laciniatum** L. Rosin-weed. Compass-plant. S. Mich., Wright's Cat.; near Capac, J. W. Stacey. S.
2148. **S. perfoliatum** L. Cup-Plant. S. Mich., Wright's Cat.
2149. **S. terebinthinaceum** Jacq. Prairie Dock. Oak openings. Ionia; Macomb Co.; Ann Arbor; Sturgis, F. P. Daniels; S. W., H. S. Pepoon; Bay Co., G. M. Bradford. Infrequent. C. & S.
2150. **S. trifoliatum** L. Avon, W. A. Brotherton. Rare.
2150a. **H. helianthoides** (L.) B. S. P. Rochester, O. A. Farwell.

HELIOPSIS Pers.

2151. **H. scabra** Dunl. Rough Ox-Eye. Detroit, O. A. Farwell.

RUDBECKIA L.

*2152. **R. hirta** L. Black-eyed Susan. Meadows and low places, apparently introduced. Common. Th.
*2153. **R. laciniata** L. Tall Cone-flower. Low grounds. Frequent. Th.
*2154. **R. speciosa** Wenderoth. Flat-headed Cone-flower. Wet soil. Ionia Co.; Flint; Macomb Co.; Alma; Ann Arbor. Infrequent. C. & S.
2155. **R. triloba** L. Thin-leaved Cone-flower. Washington, Macomb Co., Cooley. Rare.

RATIBIDA Raf. LEPACHYS Raf.

*2156. **R. pinnata** (Vent.) Barnhart. Gray-headed Cone-flower. *Lepachis pinnata* Torr. & Gray. Dry ground. Ionia Co.; Grand Rapids; S. W., H. S. Pepoon. S. Mich. Infrequent. C. & S.

BRAUNERIA Neck. ECHINACEA Moench.

2157. **B. pallida** (Nutt.) Britton. Pale Purple Cone-flower. *Echinacea angustifolia* DC. Keweenaw Co., O. A. Farwell.
2158. **B. purpurea** (L.) Britton. Purple Cone-flower. *Echinacea purpurea* Moench. Univ. Herb. Harrington; Grand Rapids, Delia Bailey; St. Joseph, Dr. Houghton, 1838. Rare. S. W.

HELIANTHUS L.

*2159. **H. decapetalus** L. Thin-leaved Wild Sunflower. Low grounds. Frequent. L. P.

*2160. **H. divaricatus** L. Rough Sunflower. Dry woods. Common. Th.

*2161. **H. giganteus** L. Tall Sunflower. Low grounds; variable. Ionia Co.; Flint; Ann Arbor; Alma; Macomb Co. and northward. Common. Th.

2161a. **H. giganteus subtuberosus** (Bourg.) Britton. Rochester, Birmingham, O. A. Farwell.

2162. **H. grosseserratus** Martens. Saw-toothed Sunflower. Port Huron, C. K. Dodge; S. W., H. S. Pepoon.

2163. **H. hirsutus** Raf. Stiff-haired Sunflower. Dry soil. Ann Arbor, and S. W.

2164. **H. Maximiliani** Schrad. Adventive from the west. Bay Co., G. M. Bradford; Houghton, Detroit, O. A. Farwell.

2165. **H. microcephalus** T. & G. S. W., H. S. Pepoon.

2166. **H. mollis** Lam. Hairy Sunflower. Manistee, F. P. Daniels.

2167. **H. occidentalis** Riddell. Few-leaved Sunflower. Sterile soil. Ionia Co.; Flint; Grand Rapids; Macomb Co. Infrequent. C. & S.

2168. **H. petiolaris** Nutt. Prairie Sunflower. Introduced from the west to Hubbardston, C. F. Wheeler; Port Huron, C. K. Dodge.

2169. **H. scaberrimus** Ell. Stiff Sunflower. *H. rigidus* Desf. Dry soil. Ann Arbor, Winch. Cat. S.

*2170. **H. strumosus** L. Pale-leaved Wood Sunflower. Dry soil. Frequent. C. & S.

2171. **H. strumosus macrophyllus** (Willd.) Britton. *H. strumosus* var. *mollis* T. & G. Sturgis, F. P. Daniels.

2172. **H. tracheliifolius** Mill. Throatwort Sunflower. Copses. S. Mich., Wright's Cat.; Macomb Co., Dr. D. Cooley. C. & S.

*2173. **H. tuberosus** L. Jerusalem Artichoke. Sparingly escaped from gardens.

VERBESINA L. ACTINOMERIS Nutt.

2174. **V. alternifolia** (L.) Britton. *Actnomeris squarrosa* Nutt. Rich soil. S. Mich., Wright's Cat.; Dundee, C. F. Wheeler; Hudson, W. J. B.; islands in the Detroit River, Maclagan. Can Cat. S.

COREOPSIS L.

2175. **C. lanceolata** L. Lance-leaved Tickseed. *C. lanceolata angustifolia* T. & G. Sandy shores of Lake Huron; sand hills, Cheboygan Co., B. & K.; sandy plains near Baldwin, W. J. B.; south to Macomb Co., Cooley; Elk Rapids, W. S. Cooper. Th.

2176. **C. palmata** Nutt. Stiff Tickseed. S. Mich., Wright's Cat.; shore of Barron Lake, C. F. Wheeler, 1890; Gull Prairie, Dr. Houghton, 1838.

2177. **C. tinctoria** Nutt. Garden Tickseed. Bay City, G. M. Bradford.

2178. **C. tripteris** L. Tall Coreopsis. Tall Tickseed. Detroit; Macomb Co.; Oakland Co.; S. Mich., Wright's Cat.; Ionia; Grand Rapids; Ann Arbor. S.

2179. **C. verticillata** L. Whorled Tickseed. Samaria, Macomb Co., Grand Rapids, C. F. Wheeler.

BIDENS L.

2180. **B. aristosa** (Michx.) Britton. Belle Isle, O. A. Farwell.

*2180. **B. Beckii** Torr. Water Marigold. S. Mich., Wright's Cat.; Ann Arbor; Wexford Co., Cooley; Manistee, E. J. Hill; Indian River, C. F. Wheeler; Orion, O. A. Farwell; St. Clair Lake and River, C. K. Dodge. Rare.

*2182. **B. cernua** L. Smaller Bur-Marigold. Wet places. Cheboygan Co., Kofoid; Orion, Detroit, O. A. Farwell; Algonac, W. S. Cooper. Southward. Frequent. Th.

2182a. **B. cernua elliptica** Weigand, in part. Belle Isle, O. A. Farwell.

*2183. **B. laevis** (L.) B. S. P. Large Bur-Marigold. *B. chrysanthemoides* Michx. Swamps. Common. Th.

*2184. **B. comosa** (A. Gray) Wiegand. Leafy-bracted Tickseed. *B. connata comosa* A. Gray. Detroit, O. A. Farwell; Alma; Ann Arbor. Infrequent.

*2185. **B. connata** Muhl. Swamp Beggar-ticks. Low grounds. Common. Th.

2185a. **B. connata anomala** O. A. F. Awns upwardly barbed, O. A. F. Detroit, O. A. Farwell.

2185b. **B. connata petiolata** (Nutt.) O. A. F. Detroit, O. A. Farwell.

2186. **B. discoidea** (T. & G.) Britton. Small Beggar-ticks. *Coreopsis discoidea* Torr. & Gray. Wet ground. Ionia Co.; Manistee, F. P. Daniels. Scarce.

*2187. **B. frondosa** L. Common Beggar-ticks. Stick-tight. Low grounds. A troublesome weed. Common. Th.

2187a. **B. trichosperma** (Michx.) Britton. Orion, O. A. Farwell.

2188. **B. trichosperma tenuiloba** (A. Gray) Britton. *Coreopsis trichosperma tenuiloba* A. Gray. Swamps. Flint; Montcalm Co.; Ionia Co.; S. Mich., Wright's Cat.; Orion, O. A. Farwell. Common in pine country. Infrequent elsewhere.

2188a. **B. vulgaris** Greene. Detroit, O. A. Farwell.

2188b. **B. vulgaris puberula** (Wieg.) Greene. Detroit, O. A. Farwell.

GALINSOGA R. & P.

2189. **G. parviflora** Cav. Detroit, W. S. Cooper. Introduced.

HELENIUM L.

*2190. **H. autumnale** L. Sneeze-weed. River banks. Common. C. & S.

2191. **H. tenuifolium** Nutt. Fine-leaved Sneeze-weed. Port Huron, C. K. Dodge.

ACHILLEA L.

2192 **A. lanulosa** Nutt. Western Yarrow. Rochester, W. S. Cooper; Frequent in U. P., O. A. Farwell.

*2193. **A. Millefolium** L. Common Yarrow or Milfoil. Fields. Common.

2193a. **A. Ptarmica** L. Sneezewort. Mich., Gray's Manual. Rare. Th.

ANTHEMIS L.

2194. **A. arvensis** L. Corn Chamomile. Introduced. Three Rivers, C. F. Wheeler; Keweenaw Co., near Detroit, O. A. Farwell; St. Clair Co., C. K. Dodge.

*2195. **A. Cotula** DC. May-weed. Dog-Fennel. Roadsides. Common. Th.

CHRYSANTHEMUM L.

2196. **C. Balsamita tanacetoides** Boiss. Mint-Geranium. Escaped from gardens.

*2197. **C. Leucanthemum** L. Ox-eye or White Daisy. Meadows and pastures. Th.

2197a. **C. Parthenium** (L.) Pers. Keweenaw Co., O. A. Farwell.

MATRICARIA L.

2198. **M. inodora** L. Scentless Camomile. Flint, Dr. D. Clark.

TANACETUM L.

2199. **T. Huronense** Nutt. Lake Huron Tansy. Sand dunes at the head of Little Traverse Bay, and northward. Frequent; Elk Rapids, W. S. Cooper.

*2200. **T. vulgare** L. Common Tansy. Escaped from gardens. Frequent.

*2201. **T. vulgare crispum** DC. More common than the species.

ARTEMISIA L.

2202. **A. Abrotanum** L. Southernwood. Escaped in Keweenaw Co.. O. A. Farwell.

2203. **A. Absinthium** L. Common Wormwood. Ionia Co.; Flint; Gratiot Co.; Keweenaw Co., O. A. Farwell. Sparingly escaped from gardens.

2104. **A. annua** L. Annual Wormwood. Detroit, O. A. Farwell; New Baltimore, J. W. Stacey.

*2205. **A. biennis** Willd. Biennial Wormwood. A roadside weed, lately introduced from the west and extending throughout the State.

2206. **A. Canadensis** Michx. Canada Wormwood. Sand dunes. Barron Lake; New Buffalo, C. F. Wheeler; Ottawa Co.; Emmet Co.; to Lake Superior. Th.

*2207. **A. caudata** Michx. Tall Wormwood. Bay Co., G. M. Bradford. Sandy fields. Ionia Co.; Montcalm Co.; S. Haven, L. H. Bailey. Infrequent. C. & S.

2208. **A. gnaphaloides** Nutt. Keweenaw Co., O. A. Farwell.

2209. **A. Ludoviciana** Nutt. Western Mugwort. Dry banks. Niles, J. T. Scoville. Univ. Herb.; Keweenaw Co., O. A. Farwell. Rare.

2210. **A. Stellariana** Bess. Port Austin. C. A. Davis; Keweenaw Co., O. A. Farwell.

2211. **A. vulgaris** L. Common Mugwort. Waste places. Infrequent.

TUSSILAGO L.

2212. **T. Farfara** L. Sault de Ste. Marie, Whitney's Cat.; Washington, Macomb Co., W. A. Brotherton.

PETASITES Gaertn.

2213. **P. palmata** (Ait.) A. Gray. Harrisville, Oscoda, Alger, and westward; Keweenaw Co., O. A. Farwell; St. Clair Co., C. K. Dodge.

ARNICA L.

2214. **A. lanceolata** Nutt. Northern Arnica. Shores of Lake Superior, Gray's Manual; Copper Harbor, Whitney's Catalogue. "It has been confused with *A. Chamissonis* of the west." Britton's Manual.

ERECHTITES Raf.

*2215. **E. hieracifolia** (L.) Raf. Fireweed. New clearings. Common northward. Th.

MESADENIA Raf. CACALIA L., in part.

2216. **M. atriplicifolia** (L.) Raf. Pale Indian Plantain. *Cacalia atriplicifolia* L. Woods. Ionia Co.; Alma; Ann Arbor; Manistee and Sturgis, F. P. Daniels; Battle Creek, W. S. Cooper; Berrien Co., H. S. Pepoon. Frequent.

2217. **M. tuberosa** (Nutt.) Britton. Tuberous Indian Plantain. *Cacalia tuberosa* Nutt. S. Mich., Wright's Cat.; three miles east of Kalamazoo, 1838, Dr. D. Houghton; Huron and Tuscola counties; Mottville, I. N. Mitchell; Bay Co., G. M. Bradford.

SYNOSMA Raf. CACALIA L., in part.

2218. **S. suaveolens** (L.) Raf. Sweet-scented Indian Plantain. *Cacalia suaveolens* L. Lodi, Miss Clark, in Winch. Cat.; Gray's Manual. Rare.

SENECIO L.

*2219. **S. aureus** L. Golden Ragwort. Squaw-weed. Very variable. Common. Th.

2220. **S. aureus gracilis** (Pursh.) Britt. A slender form of the Life-root, is frequent at Linden Park. Detroit, O. A. Farwell.

2221. **S. aureus lancelatus** Oakes. Muskegon Co., C. D. McLouth.

*2222. **S. Balsamitae** Muhl. Balsam Groundsel. *S. aureus Balsamitae* T. & G. Berrien Co., H. S. Pepoon. Northward to Keweenaw Co.

*2223. **S. discoideus** (Hook.) Britton. North of Chandler's marsh near Lansing, W. J. B.; Keweenaw Co., O. A. Farwell.

*2224. **S. obovatus** Muhl. Round-leaf Squaw-weed. *S. aureus obovatus* T. & G. More common southward.

*2225. **S. vulgaris** L. Common Groundsel., Flint; Macomb Co.; S. W. Mich. Wright's Cat.; Keweenaw Co., O. A. Farwell.

ARCTIUM L.

*2226. **A. Lappa** L. Great Burdock. Waste places. Common. Th.

2227. **A. minus** Schk. Common Burdock. *A. Lappa minus* A. Gray. Keweenaw Co., O. A. Farwell; Bay Co., G. M. Bradford.

CARDUUS L. CNICUS L. in part.

*2228. **C. altissimus** L. Tall Thistle. *Cnicus altissimus* Willd. Ionia Co.; Ann Arbor; Flint; Macomb Co.; Bay Co., G. M. Bradford. Infrequent. C. & S.

*2229. **C. arvensis** (L.) Robs. Canada Thistle. *Cnicus arvensis* Hoffm. Fields. A vile pest, introduced from Europe. Th.

2230. **C. crispus** L. Port Huron, C. K. Dodge. Introduced from Europe.

*2231. **C. discolor** (Muhl.) Nutt. Field Thistle. *Cnicus altissimus discolor* A. Gray. Meadows. Montcalm Co.; Tuscola Co.; Ann Arbor; Ionia Co.; Flint; and southward. Flowers sometimes white. Infrequent. C. & S.

2232. **C. Hillii** (Canby.) Porter. Hill's Thistle. *Cnicus Hillii* Canby. Muskegon, C. F. Wheeler; Rochester, O. A. Farwell.

*2233. **C. lanceolatus** L. Common or Bull Thistle. Roadsides and fields. Common. Th.

*2234. **C. muticus** (Mich.) Pers. Swamp Thistle. *Cnicus muticus* Pursh. Swamps, low land. Frequent. Th.

2235. **C. odoratus** (Muhl.) Porter. *C. pumilus* Nutt. S. W. Mich., Wright's Cat.; Macomb Co., Dr. Cooley; Baldwin, W. J. B.; Cheboygan Co., B. & K.

2236. **C. Pitcheri** (Torr.) Porter. Pitcher's Thistle. *Cnicus Pitcheri* Torr. Shores of the Great Lakes; sand dunes.

2237. **C. spinosissimus** Walt. Yellow Thistle. *Cnicus horridus* Pursh. N. shore Lake Superior, Agassiz; Grand Detour, T. C. Porter.

2238. **C. undulatus** Nult. Wavy-leaved Thistle. *Cnicus undulatus* A. Gray. Islands of Lakes Huron and Michigan, A. Gray; Drummond's I., Winch. Cat. N. & U. P.

MARIANA Hill. SILYBUM Gaertn.

*2239. **M. Mariana** (L.) Hill. Milk Thistle. *Silybum Marianum* Gaertn. Occasional.

ONOPORDON L.

2240. **O. Acanthium** L. Cotton or Scotch Thistle. Grand Rapids; Detroit, O. A. Farwell. Rare.

CENTAUREA L.

2241. **C. Cyanus** L. Blue-bottle. Escaped from gardens.

2242. **C. nigra** L. Black Knapweed. Centaury. Bay City, G. M. Bradford.

ECHINOPSIS.

2243. **E. sphaerocephalus** L. Well established at Three Rivers, C. F. Wheeler. Grand Rapids, H. C. Skeels. Introduced from Europe as a plant for bees.

INDEX.

Page.

Abele 67
Abies 39
Abutilon 100
Acacia, False 92
  Three-thorned 91
Acalypha 96
Acanthaceæ 122
Acanthus Family 122
Aceraceæ 98
Acer 98
Acerates 112
Achillea 134
Achroanthes 66
Acnida 74
Acorus 58
Actæa 78
Actinomeris 133
Adam-and-Eve 66
Adder's-Mouth 66
  Tongue 34, 61
Adenocaulon 132
Adder's-Tongue Family 61
Adder's-Tongue, Yellow ... 61
Adiantum 37
Adicea 71
Adlumia 81
Adopogon 126
Æsculus 98
Agassiz, Louis, Catalogue of 9
Agastache 115
Agrimonia 88
Agrimony 88
Agropyron 49
Agrostemma 75
Agrostis 46
Ailanthus 95
Aizoaceæ 74
Ajuga 115
Alder 69
  Black 97
  Green or Mountain 69
  Smooth 69
  Speckled or Hoary 69
Aletris 61
Alexanders 106
Alfalfa 92
Alfilarilla 94
Alisma 42
Alismaceæ 42
Allionia 74
Allium 61
Almendinger, Miss E. C., regarding flora 9
Alnus 69
Alopecurus 45
Alsine 75, 76
Althaea 99
Alum-root 85
Alyssum 84
Amaranthaceæ 74
Amaranth 74
Amaranth Family 74
Amaranthus 74

Page.

Amaryllidaceæ 63
Amaryllis Family 63
Ambrosia 127
Ambrosiaceæ 127
Amelanchier 89
Ammophila 46
Amorpha 92
Ampelopsis 99
Amphicarpæa 94
Anacardiaceæ 97
Anagallis 110
Anaphalis 131
Andromeda 108
Andropogon 42
Androsace 108
Anemone 78
  False Rue 78
  Long-fruited 78
  Rue 78
  Wind 78
  Wood 78
Anemonella 78
Angelica 104
Anonaceæ 77
Antennaria 131
Anthemis 134
Anthoxanthum 44
Anychia 76
Apera 46
Aphyllon 122
Apios 94
Aplectrum 66
Apocynaceæ 111
Apocynum 112
Apple Family 89
Apple 89
  American Crab 89
  Cultivated 89
  May 80
  of Peru 117
  Thorn 118
Aquilegia 78
Arabis 83
Araceæ 58
Aralia 104
Araliaceæ 104
Arbor Vitæ 39
Arbutus, Trailing 108
Arceuthobium 71
Archangelica 104
Arctium 135
Arctostaphylos 108
Arenaria 76
Arethusa 65
Argentina 87
Arisæma 58
Aristolochia 71
Aristolochiaceæ 71
Aristida 44
Arnica 135
Aronia 89
Arrhenatherum 46
Arrow-head 42
  wood 123, 124

Page.

Artemisia 134
Artichoke, Jerusalem 133
Arum, Arrow 58
  Dragon 58
Arum Family 58
Arum, Water 58
Asarum 71
Asclepiadaceæ 112
Asclepias 112
Ash, American Mountain.. 89
  Black 110
  Blue 110
  Green 110
  Northern Prickly 95
  Prickly 95
  Red 110
  White 110
Asimina 77
Asparagus 62
Aspen 67
  American 68
  Large-tooth 67
Asphodel, False 60
Aspidium 36
Asplenium 37
Asprella 50
Aster 129, 130, 131
Atheropogon 47
Astragalus 92
Atragene 79
Atriplex 73
Avena 46
Avens, Purple 88
  Water 88
  White 88

Bailey, L. H., as collector. 4
Bailey, L. H., Carices Keweenaw Penin. 10
Bailey, L. H., plants at South Haven 9
Balsaminaceæ 98
Balsam 99
  Wild Apple 125
Baneberry 78
  Red 78
  White 78
Baptisia 91
Barbarea 82
Barberry 80
Barberry Family 80
Barley 50
Barren Soil Indicated by Plants 16
Bartonia 111
Basil 117
Basswood 99
Batrachium 79
Bayberry 67
Beal, W. J., as author of this work 3
Beal, W. J., Flora of Sandy Pine Plains 16

Page.

Bean, Sacred 93
Wild 93
Bearberry 108
Beardslee, H. C., List of Cheboygan Plants 11
Beard-tongue 119
Beaver-Poison 106
Bedstraw, Northern 123
Rough 123
Small 123
Sweet-scented 123
Bee-Balm 116
Beech 69
American 69
Blue or Water 69
Beech-drops 122
Bee Plants 24, 84
Beggar-tricks, Common. 133, 134
Swamp 133
Bellflower 125
Marsh 125
Tall 125
Bellwort 61
Benjamin-bush 80
Benzoin 80
Berberidaceæ 80
Berberis 80
Bergamot 116
Berry, Pigeon 74
Berula 106
Betula 69
Betulaceæ 69
Bibliography 9
Bicuculla 81
Bidens 133
Bignoniaceæ 122
Bilberry 109
Bindweed 112
Black 72
Hedge 72
Birch 69
Black 69
Cherry 69
Dwarf 69
Low 69
Paper or Canoe 69
White 69
Yellow or Gray 69
Birthroot 62
Birthwort Family 71
Bishop's Cap 85
Bitter-nut 67
Bittersweet 118
Climbing or Shrubby... 97
Bitted-weed 127
Birthwort 71
Birthwort Family 71
Blackberry, Common or High 87
Low 87
Running Swamp 86, 87
Bladder Ketmia 100
Bladder-nut 98
Bladderwort Family 121
Bladderwort 121
Blazing-Star 60, 128
Blephariglottis 65
Blephilia 116
Blite, Strawberry 73
Blodgett, H. T., Plants of Mason County 10
Blitum 73
Blood-root 80
Bloodwort Family 80
Blueberry, Canada 109
Dwarf 109
Low 109
Swamp 109
Blue Bells 125
Blue-bottle 136
Blue-eyed Mary 119
Blue Grass, Kentucky 48
Blue-hearts 120
Bluets 123
Blue-weed 114

Page.

Bœhmeria 71
Boneset 128
Borage Family 113
Borraginaceæ 113
Borago 114
Botrychium 35
Bouncing Bet 75
Bouteloua 47
Bowman's Root 86
Box-Elder 98
Brachyactis 131
Brachyelytrum 45
Bradford, G. M., as Collector 5
Brake 35, 36, 37
Cliff 37
Rock 37
Bramble 37, 86, 87
Brasenia 76
Brassica 81, 82
Brauneria 132
Britton, N. L., Manual followed 5, 9
Briza 48
Bromus 49
Brooklime, American 120
Broom-rape Family 122
Broom-rape 122
Brook-weed 109
Brotherton, W. K., as Collector 5
Brunella (see Prunella) ... 116
Buchnera 120
Buckbean 111
Buckeye, Ohio 98
Buckthorn 98
Buckthorn Family 98
Buckwheat 72
Buckwheat Family 72
Buckwheat, Clambing, False 72, 73
Buda 76
Buffalo-berry 102
Bugbane 78
Bugle-weed 115, 117
Bugloss, Viper's 114
Bug-seed 74
Bulrush 59, 60
Bunch-berry 106
Bunch-flower Family 60
Burdock 135
Burnet, Canadian, Great.. 88
Burning-Bush 97
Bursa 83
Burt, W. A., Catalogue of. 9
Bush-Clover 93
Strawberry 97
Butter and Eggs 119
Buttercup 79
Butterfly-weed 112
Butternut 67
Butter-weed 135, 136
Butterwort 121
Button-bush 123
Buttonwood 86

Cabbage, Skunk 124
Cacalia 135
Cactaceæ 102
Cactus Family 102
Caesalphinaceæ 91
Cakile 81
Calamagrostis 46
Calamintha 117
Calamint 117
Calamovilfa 46
Calamus 58
Calceolaria 102
Calla 58
Callitrichaceæ 96
Callitriche 96
Calopogon 65
Caltha 77
Calypso 66

Page.

Camassia 61
Camelina 59, 83
Cammelinaceæ 59
Camomile 134
Campanula 125
Campanula Family 125
Campanulaceæ 125
Campbell, D. H., Plants of Detroit river 9
Campion, Bladder 75
Campion, Starry 75
White 75
Camptosorus 37
Cancer-root 122
Cannabis 71
Caper Family 84
Capnoides 81
Capparidaceæ 84
Caprifoliaceæ 123
Capriola 83
Capsella 83
Caraway 106
Cardamine 82
Cardinal-flower 125
Carduus 135, 136
Carex 53
Carnation 75
Carpet-weed 74
Carpinus 69
Carrion-Flower 63
Carrot 104
Carrot Family 104
Carum 106
Carya 67
Caryophyllaceæ 75
Cassandra 108
Cassia 91
Castalia 77
Catalpa 122
Castanea 69
Castilleia 121
Cat-brier 63
Catchfly 75
Night-flowering 75
Sweet-William 75
Sleepy 75
Catgut 92
Cat-Mint 115
Catnep 115
Cat-tail 40
Cat-tail Family 40
Caulophyllum 80
Ceanothus 99
Cedar, Red 40
White 39
Celandine 80
Celastraceæ 97
Celastrus 97
Celtis 70
Cenchrus 44
Centaurea 136
Centaury 110, 136
Cephalanthus 123
Cerastium 76
Ceratophyllaceæ 77
Ceratophyllum 77
Cercis 91
Chænorrhinum 119
Chærophyllum 105
Chætochloa 44
Chamædaphne 108
Chætochloa 44
Chamælirium 60
Chamænerion 103
Chamomile, Wild 134
Charlock, English 82
Cheat 49
Chelone 119
Chelidonium 80
Chenopodiaceæ 73
Chenopodium 73
Cherry, Black 91
Choke 91
Dwarf 91
Ground 118

Page.

Cherry, Red 91
Sand 91
Chervil 105
Chess, Wild 49
Chestnut 69
Chickweed 76
Field 76
Forked 76
Larger Mouse-ear 76
Mouse-ear 76
Chicory 126
Chicory Family 126
Chimaphila 107
Chinquapin, Water 77
Chiogenes 109
Choke-berry 89
Chrysanthemum 134
Chrysopogon 42
Chrysosplenium 85
Cichoriaceæ 126
Cichorium 126
Cicuta 106
Cimicifuga 78
Cinque-foil, Common 88
Marsh 87
Shrubby 87
Silvery 88
Tall 87
Three-toothed 87
Cinna 45
Circæa 103
Cistaceæ 100
Cladium 53
Clammy-weed 84
Clark, H. L., Notes on the Flora of Eaton Co. 10
Claytonia 74
Clearweed 71
Cleavers 123
Climate 12
Clematis 78, 79
Cleome 84
Clinopodium 117
Clintonia 62
Clotbur 128
Clover 92
Alsike 92
Bush 93
Low Hop 92
Mammoth 92
Prairie 92
Rabbit-foot or Stone... 92
Red 92
Running Buffalo 92
Sweet 92
White 92
Yellow or Hop 92
Club-Moss 38
Club-Moss Family 38
Cnicus 135, 136
Cockle 75
Cocklebur 128
Coeloglossum 64
Coffee-tree, Kentucky 91
Cohosh 78
Black 78
Blue 80
Cole, Miss Emma J., as Collector 4
Cole, Miss Emma J., List of Plants Collected in and about Grand Rapids. 10
Coleman, N., Catalogue of. 9
Colic-root 61
Collinsia 119
Collinsonia 117
Coltsfoot 135
Sweet 135
Columbine 78
Columbo, American 111
Comandra 71
Comarum 87
Comfrey 114
Wild 113

Page.

Commelina 83
Comparison of trees of Michigan with those of other countries 132
Compass-Plant 128
Compositæ found on Pine Plains 17
Composite Family 128
Comptonia 67
Cone-flower, Purple 132
Coniferæ 39
Conioselinum 105
Conium 106
Conopholis 122
Conringia 81
Contents 7
Convallariaceæ 62
Convolvulaceæ 112
Convolvulus 112
Convolvulus Family 112
Cooley, D., Catalogue of.... 9
Cooper, W. S., as Collector 5
Coptis 78
Coral-berry 124
Corallorhiza 66
Coral-root 66
Cockle, Corn 75
Cord-Grass, Fresh-water.. 47
Coreopsis 133, 134
Corispermum 74
Cornaceæ 106
Cornel 106
Dwarf 107
Panicled 107
Silky 106
Corn Salad 125
Cornus 106
Corpse-plant 107
Corydalis 81
Golden 81
Pale 81
Corylus 69
Cotton-wood 67
Cowbane, Spotted 105
Cowberry 87
Cow-herb 75
Cowslip, American 110
Virginian 114
Cow-Wheat 121
Crab-grass 43
Cracca 92
Cranberry 43
Large or American 109
Small 109
Cranberry-tree 124
Cranesbill 94
Crassulaceæ 85
Cratægus 90
Creeper, Virginia 99
Crepis 127
Cress, Bitter 79, 82
Balbous 82
Cow 81
Lake 82
Marsh 79
Purple 82
Rock 79, 81, 83
Small Bitter 79
Spring 79
True Water 79, 82
Water 79
Winter 82
Yellow 82
Crosswart 110
Crotalaria 91
Crowberry Family 97
Crowberry, Black 97
Crowfoot 79
Crowfoot Family 77
Crowfoot, Bristly 79
Bulbous 79
Common White Water. 79
Cursed 79
Early 79

Page.

Crowfoot, Hooked 79
Rock 79
Sea-side 79
Small-flowered 79
Stiff Water 79
Tall 79
Yellow Water 79
Cruciferæ 81
Cryptogramma 37
Cryptotænia 106
Cuckoo Flower 82
Cucumber 125
Cucumber-root, Indian 62
Cucurbitaceæ 125
Cudweed 132
Low 132
Purplish 132
Culver's-root 120
Cup, Plant 132
Cupuliferæ 69
Currant, Fetid 86
Indian 124
Red 86
Wild Black 86
Cuscutaceæ 113
Cuscuta 113
Custard-apple Family 77
Cut-grass, Rice 44
Cycloloma 73
Cymbalaria 119
Cynanchum 112
Cynodon 47
Cynoglossum 113
Cynosurus 48
Cyperaceæ 50
Cyperus 50
Cypripedium 64
Cystopteris 36

Dactylis 48
Daisy, Ox-eye 134
White 134
Dalibarda 87
Dandelion 126
Dwarf 126
Fall 126
Daniels, Francis P., as Collector 5
Daniels, Francis P., Flora of the Vicinity of Manistee 10
Daniels, Francis P., Ecology of the Flora of Sturgis and vicinity 10
Danthonia 46
Darnel 49
Dasiphora 87
Dasystoma 120
Datura 118
Daucus 104
Davis, C. A., The Flora of Michigan Lakes 10
Davis, C. A., Flora of Tuscola County 10
Davis, C. A., Botanical Notes on Huron Co. 10
Davis, C. A., Herbarium of 5
Davis, Geo. W., as Collector 5
Day-flower 83
Decodon 102
Deerberry 108
Deer-grass 102
Dennstædtia 35
Dentaria 83
Deringa 106
Deschampsia 46
Desmodium 93
Dewberry 87
Dewey, L. H., List of Tecumseh Plants 11
Dianthera 122
Dianthus 75

Page.

Diarrhena 48
Dicentra 81
Dicksonia 35
Diervilla 124
Dioscorea 63
Dioscoreaceæ 63
Diplotaxis 82
Dipsaceæ 125
Dipsacus 125
Dirca 102
Distribution of Plants 12
Dock 72
  Bitter 72
  Broad-leaved 72
  Curled 72
  Narrow 72
  Pale 72
  Patience 72
  Prairie 132
  Swamp 72
  White 72
Dockmackie 123
Dodder 113
Dodecatheon 110
Dodge, C. K., as Collector. 5
Dodge, C. K., List of Port Huron Plants 10
  Flora of St. Clair Co. . 10
Doellingeria 131
Dogbane 112
Dogbane Family 111
Dogwood Family 106
Dogwood, Flowering 107
  Poison 97
  Round-leaved 107
Draba 83
Dracocephalum 116
Dragon-head 116
  False 116
Dragon-root 58
Drosera 84
Droseraceæ 84
Drupaceæ 90
Drymocallus 87
Duck's-meat 58
Duckweed 58
Duckweed Family 58
Dulichium 51
Dutchman's Breeches 81

Eagle Fern 37
Eatonia 47
Echinacea 132
Echinochloa 43
Echinocystis 125
Echinodorus 42
Echinops 136
Echinospermum 114
Echium 114
Ecology 3
Elæagnaceæ 102
Elder 123
  Marsh 127
  Red-berried 123
  Wild 104
Elecampane 132
Eleocharis 51, 52
Eleusine 47
Elm 70
  American 70
  Cork 70
  Red 70
  Rock 70
  Slippery 70
Elodea 42, 100
Elymus 50
Empetraceæ 97
Empetrum 97
Epigæa 108
Epilobium 103
Epiphegus 122
Equisetaceæ 38
Equisetum 38

Page.

Eragrostis 47
Erechtites 135
Ericaceæ 108
Ericaceæ, Found on Pine Plains 17
Erigenia 106
Erigeron 131
Eriocaulaceæ 59
Eriocaulon 59
Eriophorum 52
Erodium 94
Eryngium 105
Erysimum 83
Erythræa 110
Erythronium 61
Euonymus 97
Eupatorium 128
Euphorbia 96
Euphorbiaceæ 96
Euphrasia 121
Euthamia 129
Evening Primrose 103
Evening Primrose Family. 102
Everlasting, Pearly 132
Eyebright 121

Fagopyrum 72
Fagaceæ 69
Fagus 69
Falcata 94
False Heather 100
False Mermaid Family 97
Families of Plants not represented on the Pine Plains 17
Families, Sequence of 5
Farwell, O. A., List of Plants of Detroit 10
Farwell, O. A., as Collector 5
Fennel 105
Fennel-flower 80
Fennel, Dog 134
Fern, Beech 36
  Chain 37
  Christmas 36
  Cinnamon 35
  Flowering 35
  Male 36
  Sensitive 35
  Shield 36
Fertile Soil, Plants Indicating 18
Fertile Soil Indicated by Plants 18
Fescue, Meadow 49
  Taller 49
  Sheep's 49
Festuca 49
Fever-bush 80
Fever-wort 124
Fig, Indian 102
Figwort Family 119
Figwort 115, 119
Filbert 69
Filices 34, 35, 36
Filix 36
Fimbristylis 51
Finger-Grass 42, 43
Fir, Balm of Gilead 39
  Balsam 39
Fire Pink 75
Fireweed 135
Five-Finger, Marsh 87, 88
Flag, Blue 63
  Cat-tail 40
  Sweet 58
Flax 95
  False 83
Flax Family 95
Fleabane, Common 131
  Daisy 131
Flixweed 83
Floating Heart 111

Page.

Flœrkea 97
Flora, Eastern and Western Sides Contrasted ... 20
Flora of Pine Plains 16
Flower-de-Luce 63
Fœniculum 105
Fœrste, A. F., Plants of Belle Isle 9
Fog-fruit 115
Forage Plants 23
Forget-me-not 114
Foster, J. W., Catalogue of 9
Four-o'clock 74
Four-o'clock Family 74
Foxglove, Downy False ... 120
  Smooth False 120
Foxtail, Meadow 45
Fragaria 87
Frasera 111
Fraxinus 110
Fringed-Orchis, Ragged ... 65
Frost-weed 100
Fuirena 52
Fumaria 81
Fumariaceæ 80
Fumitory Family 80
Fumitory, Climbing 81

Galeopsis 116
Galeorchis 64
Gale, Sweet 67
Galinsoga 134
Galium 123
Garlic, Wild 61
Gaultheria 108
Gaura 103
Gaylussacia 108
Gentianaceæ 110
Gentian Family 110
Gentian, Closed 111
  Fringed 111
  Horse 124
  Smaller Fringed 111
  Soapwort 111
  Spurred 111
Gentiana 111
Geraniaceæ 94
Geranium 94
Geranium Family 94
Gerardia 120
  Purple 120
  Slender 121
Germander, American 115
Geum 88
Gillenia 86
Gill-over-the-Ground 115
Ginger, Wild 71
Ginseng Family 104
Ginseng 104
  Dwarf 104
Glecoma 115
Gleditsia 91
Globe Flower 77
Glyceria 48
Gnaphalium 132
Goat's-beard 126
Goat's Rue 92
Golden-rod 129
Golden Seal 77
Goldthread 78
  Three-leaved 78
Good-King-Henry 73
Goodyera 66
Gooseberry Family 86
Gooseberry, Missouri 86
  Swamp 86
Goosefoot Family 73
Goosefoot 73, 125
  Maple-leaved 73
  Oak-leaved 73
Gourd Family 125
Gramineæ 17, 42
Grape 99
  Summer 99

Page.

Graphephorum 48
Grass Family 42
Grass, Arrow 42
Barnyard 43
Beard 42
Bent 46
Bermuda 47
Black 60
Blue 48
Blue-eyed 63
Blue-Joint 46
Bottle 60
Bottle-brush 50
Brome 49
Brown Bent 46
Bur 44
Canary 44
Cockspur 43
Cord 47
Cotton 52
Couch 49
Crab 43
Deer 102
Dog's-tail 47, 48
Drop-seed 45
Eel 42
English Blue 49
Feather 45
Fescue 49
Fiorin 46
Fowl Meadow 48
Foxtail 44, 45
Goose 123
Grama 47
Hair 46
Holy 44
Hungarian 44
Indian 42
June 48
Knot 72
Lyme 50
Manna 48
Meadow 48, 49
Millet 44, 45
Oat 45, 46
Old-witch 43
Orchard 48
Panic 43
of Parnassus 85
Pepper 81
Pigeon 44
Porcupine 45
Quack 49
Quaking 48
Quick 49
Quitch 49
Rattlesnake 48
Ray 49
Red Top 46, 47
Reed 47
Reed Canary 44
Rush 45
Rye 49
Sand 47
Sand-Reed 46
Scorpion 114
Spear 48
Squirrel-tail 50
Star 63
Stink 47
Sweet Vernal 44
Thin 46
Triple-awned 44
White 44
Whitlow 83
Wire 47, 48
Wood Reed 45
Yard 47
Yellow-eyed 63
Gratiola 120
Gray, Dr. Asa, on Causes for Distribution of Trees 22
Manual of Botany Consulted 9

Page.

Greenbrier 63
Common 63
Grindelia 128
Gromwell 114
Corn 114
False 114
Ground-nut 94
Grossulariaceæ 86
Ground Pine 38
Groundsel 135
Gum-plant 128
Gymnandeniopsis 64
Gymnocladus 91
Gypsophila 75
Gyrostachys 65

Habenaria 64, 65
Hackberry 70
Helenia 111
Haloragidaceæ 104
Hamamelidaceæ 86
Hamamelis 86
Harbinger-of-Spring 106
Hardhack 86
Harebell 125
Harrington, M. W., Herbarium of 11
Hartmannia 103
Haw, Black 124
Hawkbit 126
Hawksbeard 127
Hawkweed 126, 127
Long-bearded .....
Hawthorn 90
Hazel-nut 69
Beaked 69
Wild 69
Heal-all 116
Heart's-ease 101
Heath Family 108
Hedeoma 116
Hedgehog-grass 44
Helenium 134
Helianthemum 100
Helianthus 133
Heliopsis 132
Hemerocallis 61
Hemicarpha 53
Hemlock 39
Ground 40
Parsley 105
Poison 106
Water 106
Hemp 71,
Indian 112
Hemp-Nettle 116
Hemp-weed, Climbing 128
Henbane, Black 118
Hepatica 78
Heracleum 105
Herbaria Consulted 10
Herbarium of Alma College, made by O. J. Stillwell and C. A. Davis.... 10
Herbarium of Geo. L. Ames 11
W. J. Beal 11
Dr. D. Clark 10
Miss Mary H. Clark.... 11
G. F. Comstock 11
Dr. D. Cooley 10
C. A. Davis 11
M. W. Harrington..... 11
G. H. Hicks 11
Dr. D. Houghton 11
Geo. D. Sones 11
State University noticed 11
F. E. Wood 11
Herb Robert 94
Herd's-Grass 45
Hesperis 84
Heteranthera 59
Heuchera 85

Page.

Hyssop 117
Hibiscus 100
Hicks, G. H., Herbarium of 11
Hicks, G. H., as collector.. 5
Hicks, G. H., New and Rare Plants of Michigan 10
Hickory, Shag-bark 67
Hicoria 67
Hieracium 127
Hierochloe 44
Hill, E. J., Plants of Menominee Region 9
Hill, E. J., Flora of Lake Superior Region 9
Hippocastanaceæ 98
Hippuris 104
Hobble-bush 123
Hog-weed 127
Holly Family 97
Holly 97
Mountain 97
Homalocenchrus 44
Honeysuckle Family 123
Honeysuckle 124
Fly 124
Honewort 106
Hop 70
Hop-Hornbean 69
American 69
Hop-tree 95
Hordeum 50
Horehound 115
Water 117
Hornbeam 69
American 69
Hornwort Family 77
Hornwort 75, 106
Horse-brier 63
Horse-chestnut 98
Horseradish 82
Horsetail Family 38
Horsetail 38
Horseweed 131
Houghton, Dr. D., Herbarium of 11
Hound's Tongue 113
Houstonia 123
Huckleberry 108
Black 108
Hudsonia 100
Hull, W. C., List of Albion Plants 11
Humulus 70
Huntman's Cup 84
Hyacinth, Grape 61
Wild 61
Hydrastis 77
Hydrocotyle 106
Hydrophyllaceæ 113
Hydrophyllum 113
Hypericaceæ 100
Hypericum 100
Hypopitys 108
Hypoxys 63
Hyssopus 117
Hyssop 117
Giant 115
Hedge 120
Hyoscyamus 118
Hystrix 50

Iberis 81
Ilex 97
Ilicaceæ 97
Iliciodes 97
Ilysanthes 120
Impatiens 98
Imperatoria 105
Indian Physic 86
Indigo, False 91
White 91
Wild 91
Innocence 123

Page.

Ionidium 102
Inula 132
Ipomœa 113
Iridaceæ 63
Iris 63
  Lake Dwarf 63
Iris Family 63
Iron-weed 128
Iron-wood 69
Isnardia 102
Isoetaceæ 39
Isoetes 39
Isopyrum 78
Isotria 65
Iva 127
Ivy, American 99
  Kenilworth 119
  Poison 97

Jamestown-Weed 118
Jeffersonia 80
Jewel-weed 98
Joe-Pye Weed 128
Jointweed 73
Judas-tree 91
Juglandaceæ 67
Juglans 67
Juncaceæ 59
Juncoides 60
Juncus 59
June-berry 89
Juniperus 40
Juniper 40

Kalmia 108
King-nut 67
Kinnikinnik 86
Knawel 76
Kneiffia 103
Knotweed 73
Kœleria 47
Kœllia 117
Kofoid, C. A., List of Cheboygan Plants 11
Korycarpus 60
Krigia 126
Kuhnia 128
Kuhnistera 92
Kyllinga 51

Labiatæ 115
Labrador Tea 108
Lacinaria 128
Lactuca 126
Ladies' Tresses 65
Lady's Slipper 64
  Larger Yellow 64
  Showy 64
  Smaller Yellow 64
  Stemless 64
  Small White 64
Lady's Thumb 72
Lambkill 108
Lamb's-Quarters 73
Lamium 116
Laportea 71
Lappula 114
Lapsana 126
Larch 39
  American or Black 39
Larix 39
Lathyrus 94
Lauraceæ 80
Laurel Family 80
Laurel, American 108
  Ground 108
  Sheep 108
Lead-plant 92
Leaf-cup 132
Leather-leaf 108
Leatherwood 102
Lechea 101

Page.

Ledum 108
Leek, Wild 61
Leersia 44
Leguminosæ 91
Lemnaceæ 58
Lemna 58
Lentibulariaceæ 121
Leontodon 126
Leonurus 116
Lepachys 132
Lepidium 81
Leptamium 122
Leptandra 120
Leptilon 131
Leptorchis 66
Lespedeza 93
Lettuce 126, 127
  Florida 126
  Prickly 127
  Wild 126
Leucophysalis 118
Lever-wood 69
Liatris 128
Lilac 110
Liliaceæ 61
Lilium 61
Lily Family 61
Lily 61
  Day 61
  Turk's Cap 61
  Water Tuber-bearing .. 77
  Wild Yellow 61
  Yellow Pond 77
  Wild Orange-red 61
Lily-of-the-Valley 62
  False 62
Limnanthaceæ 97
Limnanthemum 111
Limnorchis 64
Limodorum 65
Limosella 120
Linaceæ 95
Linaria 119
Linden 99
Linden Family 99
Lindera 80
Linnæa 124
Linum 95
Liparis 66
Lippia 115
Liquorice, Wild 123
Liriodendron 77
Listera 65
Lithospermum 114
Live-for-ever 85
Liver-leaf 78
Livingston, Burton E., The Distribution of the Plant Societies of Kent Co. 10
Livingston, Burton E., The Relation of Soils to Natural Vegetation 10
Lizard's-tail 67
Lobeliaceæ 125
Lobelia Family 125, 126
Lobelia, Great 126
  Water 125
Locust, Clammy 92
  Common 92
  False Acacia 92
  Honey 91
Locust-tree 92
Lolium 49
Longyear, B. O., as Collector 4
Lonicera 124
Looking-glass, Venus's 125
Loosestrife Family 102
Loosestrife 102, 110
  Swamp 102
  Tufted 110
Lophanthus 115
Lopseed 122
Loranthaceæ 71
Lotus 77

Page.

Lousewort 121
Lucerne 92
Ludwigia 102, 103
Lungwort 114
Lupine, Wild 91
Lupinus 91
Luzula 60
Lychnis 75
  Evening 75
Lycium 118
Lycopodiaceæ 38
Lycopus 117
Lycopodium 38
Lyonia 108
Lyons, A. B., Medicinal Plants 9
Lysias 64
Lysiella 65
Lysimachia 109, 110
Lythraceæ 102
Lythrum 102

Macoun, John, Catalogue of Canadian Plants 10
Madder Family 123
Malanthemum 62
Maidenhair 37
Magnoliaceæ 77
Magnolia Family 77
Mallow 97
Mallow Family 99
Mallow, Common 99
  Curled 99
  High 99
  Musk 99
  Rose 100
  Swamp Rose 100
Malus 89
Malvaceæ 99
Malva 99
Mandrake 80
Man-of-the-Earth 113
Maple Family 98
Maple 98
  Ash-leaved 98
  Black Sugar 98
  Mountain 98
  Norway 98
  Red 98
  Rock 98
  Striped 98
  White or Silver 98
Mare's Tail 104
Mariana 136
Marigold, Bur 133
  Larger Bur 133
  Marsh 77
  Smaller Bur 133
Marrubium 115
Masterwort 105
Matricaria 134
Matrimony-Vine 118
Matteuccia 35
Matthiola 82
May-apple 80
May-weed 134
McLouth, C. D., as Collector 5
Meadow-Beauty Family ... 102
Meadow-Rue, Early 79
  Purplish 79
  Tall 79
Meadow-Sweet 62, 86
Medeola 62
Medicago 92
Medick, Black 92
Melbomia 93
Melampyrum 121
Melanthaceæ 60
Melastomaceæ 102
Melilot, Yellow 92
  White 92
Melilotus 92
Menispermaceæ 80

Page.

Menispermum 80
Mentha 117
Menyanthaceæ 111
Menyanthes 111
Mercury, Three-seeded ... 96
Mermaid, False 104
Mermaid-weed 104
Mertensia 114
Mesadenia 135
Mexican Tea 73
Mezereon Family 102
Micrampelis 125
Microstylis 66
Mignonette 84
Mikania 128
Milkweed Family 112
Milkweed 112
  Green 112
  Purple 112
  Swamp 112
Milkwort 95
Milium 45
Millet 45
Mimulus 119
Mint Family 115
Mint, Corn 117
  Horse 116
  Mountain 117
  Wild 117
Mistletoe 71
Mistletoe Family 71
Mist-flower 128
Mitchella 123
Mitchell, I. N., List of Michigan Plants 11
Mitella 85
Mitre-wort 85
  False 85
Moccason-flower 64
Mocker-nut 67
Moehringia 76
Mollugo 74
Monarda 116
Moneses 107
Moneywort 109, 110
Monkey-flower 119
Monotropa 107, 108
Monotropaceæ 107
Moonseed 80
Moonseed Family 80
Moonwort 108
Moosewood 102
Morning Glory 112, 113
Moraceæ 70
Morus 70
Moseley, E. H., Plants of Union City 11
Motherwort 116
Mudwort 120
Mugwort 134
Muhlenbergia 45
Mulberry 70
  Red 70
  White 70
Mullein 119
  Moth 119
  Pink 75
Musk-flower 119
Muscari 61
Musquash Root 106
Mustard Family 81
Mustard, Black 82
  Hedge 81
  Indian 82
  Tansy 83
  Tower 83
  White 81
  Worm-seed 83
Myrica 67
Myricaceæ 67
Myriophyllum 104
Myosotis 114

Nabalus 127

Page.

Naiadaceæ 40
Naias 40
Nasturtium 82
Naumbergia 110
Neckweed 120
Negundo 98
Nelumbo 77
Nemopanthes 97
Nepeta 115
Nettle Family 71
Nettle 71
  Dead 116
  False 71
  Hedge 116
  Horse 118
  Wood 70
Nettle Tree .....
Nicandra 117
Nicotiana 118
Nigella 80
Nightshade 118
  Enchanter's 103
Nightshade Family 118
Nimble Will 45
Nine-bark 86
Nipplewort 126
Nonesuch 92
Northern Plants Found on the East Side of the State 20
Nuphar 77
Nyctaginaceæ 74
Nymphæa 77
Nymphæaceæ 76
Nyssa 107

Oak Family 70
Oak 70
  Alexander's 70
  Black 70
  Bur 70
  Chestnut 70
  Dwarf Chestnut 70
  Gray 70
  Hill's 70
  Jerusalem 73
  Laurel 70
  Over-cup or Mossy-cup. 70
  Pin 70
  Poison 97
  Red 70
  Scarlet 70
  Schneck's 70
  Shingle 70
  Spanish 70
  Swamp White 70
  White 70
  Yellow 70
  Yellow-barked 70
Oat 46
Oats, Water 44, 46
Oat-grass 44
Œnothera 103
Oleaceæ 110
Oleaster Family 102
Olive Family 110
Onagra 103
Onagraceæ 102
Onion, Wild 61
Onoclea 35
Onopordon 136
Onosmodium 114
Ophioglossaceæ 34
Ophioglossum 34
Opulaster 86
Opuntia 102
Orache 73
Orchidaceæ 64
Orchis Family 64
Orchis 64
  Crane-fly 66
  Fen 66
  Showy 64

Page.

Orchis, White, Fringed.... 65
  Yellow, Fringed 65
Orpine Family 85
Orpine 85
  Garden 85
Orobanchaceæ 122
Orth, S. P., List of Plants of Imlay City 11
Oryzopsis 45
Osier, Basket 68
Osmorrhiza 106
Osmundaceæ 35
Osmunda 35
Ostrya 69
Overlapping of Northern and Southern Species 19
Oxalidaceæ 95
Oxalis 95
Ox-eye 132
Oxycoccus 109
Oxygraphis 79
Oxyphabus 74
Oxypolis 105
Oyster Plant 126

Painted-cup 121
  Scarlet 121
Palmer, Elmore, Catalogue of 9
Panicularia 48
Panicum 43
Pansy 101
Papaveraceæ 80
Papaver 80
Papaw 77
Papilionaceæ 91
Pappoose-root 80
Parnassia 85
Parnassiaceæ 85
Parsnip 105, 106
  Water 106
Parsley Family 104
Parthenocissus 99
Pasque Flower 78
Paspalum 43
Partridge-berry 123
Pastinaca 105
Pea Family 91
Pea, Beach 94
  Marsh 94
Pea-nut, Hog 94
Pear 89
Pear, Prickly 102
Pearlwort 76
Pedicularis 121
Pellæa 37
Peltandra 58
Pennycress 81
Pennyroyal, American 116
Penthonaceæ 85
Pennywort, Water 106
Penthorum 85
Pentstemon 119
Pepper Family 67
Peppergrass 81
Pepperidge 107
Peppermint 117
Peramium 66
Periwinkle 111
Persicaria 72
Perularia 64
Petalostemon 92
Petasites 135
Petunia 118
Phaca 92
Phacelia 113
Phalaris 44
Phegopteris 36
Philotria .....
Phleum 45
Phlox 113
Phlox Family 113
Phragmites 47
Phryma 122

Page.

Phrymaceæ 122
Physalis 118
Physalodes 117
Physocarpa 86
Physostegia 116
Phytolaccaceæ 94
Phytolacca 74
Picea 39
Pickerel-weed 59
Pickerel-weed Family 59
Picris 126
Pieters, A. J., Plants of Lake St. Clair 10
Pigeon-berry 74
Pig-nut 67
Pig-weed 73, 74
  Winged 73
Pilea 71
Pimpernel 110
  False 120
  Water 109
Pimpinella 105
Pine Family 39
Pine, Ground 38
  Jack 39
  Norway 39
  Red 39
  Sap 108
  White 39
Pine Drops 107
Pine Plains, Flora of 16
Pine-weed 100
Pine-sap 108
Pinguicula 121
Pink Family 75
Pink 75
  Fire 75
  Ground or Moss 113
  Mullein 75
Pinus 39
Pinweed 101
Pipe, Indian 107
Pipewort Family 59
Pipsissewa 107
Pitcher Plants 84
Plane-tree Family 86
Plant, Lead 92
Plantaginaceæ 122
Plantago 122
Plantain Family 122
Plantain 122, 123
  English 122
  Indian 135
  Mud 58
  Pale Indian 135
  Rattlesnake 66
  Tuberous Indian 135
Plant Lists Consulted, Furnished by:
  Beardslee, H. C. 11
  Bradford, G. M. 11
  Cooper, W. S. 11
  Daniels, F. P. 11
  Davis, C. A. 11
  Dewey, L. H. 11
  Dodge, C. K. 11
  Farwell, O. A. 11
  Foerste, A. F. 11
  Hull, W. 11
  Kofoid, C. A. 11
  Mitchell, I. N. 11
  Moseley, E. H. 11
  Orth, S. P. 11
  Pepoon, H. S. 11
  Stacey, I. W. 11
  St. Johns 11
Plants for bees .....
  Fast disappearing and how 28
  Immigrated from the Northeast 20
  Immigrated from the North and West 21
  Indicating fertile soil .. 18

Page.

Plants Overlapping in the G River Valley 19
Platanaceæ 86
Platanus 86
Pleurisy-root 112
Plum Family 90
Plum, Cherry 90
  Wild 90, 91
  Yellow 90
Poa 48
Podophyllum 80
Podostemaceæ 84
Podostemon 84
Pogonia 65
Poisonous Plants 27
Poke 74
Pokeweed 74
Pokeweed Family 74
Polanisia 84
Polemoniaceæ 113
Polycodium 108
Polygala 95
Polygalaceæ 95
Polygonaceæ 72
Polygonatum 62
Polygonella 73
Polygonum 72
Polypodiaceæ 35
Polypodium 37
Polypody 37
Polymnia 132
Polystichum 36
Polytænia 105
Pomaceæ 89
Pondweed 40
Pondweed Family 40
Pontederia 59
Pontederiaceæ 59
Poplar 67
  Balsam 67
  Downy 68
  Necklace 67
  White 67
Poppy Family 80
Poppy 80
  Celandine 80
Populus 67
Porteranthus 86
Portulaca 74
Portulacaceæ 74
Potato Family 117
Potomageton 40
Potato-vine, Wild 113
Potentilla 87, 88
Poterium 88
Prairie Plants 19
Prenanthes 127
Primrose Family 109
Primrose, Bird's-eye 109
  Evening 103
  Showy 103
Primula 109
Primulaceæ 109
Prince's Feather 72
Prince's Pine 107
Proserpinaca 104
Prunella 116
Prunus 90
Psilocarya 51
Ptelea 95
Pteridium 37
Pteris 37
Pterospora 107
Puccinelia 49
Puccoon 77
  of the Indians 114
  Yellow 77
Pulsatilla 78
Pulse Family 91
Purse, Shepherd's 83
Purslane Family 74
Purslane, Marsh 102
Putty-root 66
Pycnanthemum 117

Page.

Pyrola 107
  One-flowered 107
Pyrolaceæ 107
Pyrus 89

Quamasia 61
Queen of the Prairie 88
Quercitron 70
Quercus 70
Quillwort 39

Radish 82
Ragweed 127
Ragwort, Golden 135
Ranunculaceæ 77
Ranunculus 79
Raphanus 82
Raspberry, Black 86
  Dwarf 86
  Purple Flowering 87
  Wild Red 86
Ratibida 132
Rattle-box 91
Rattlesnake-Master 66
  Plantain 66
  Root 127
  Weed 127
Razoumofskya 71
Red-bud 91
Red-osier 107
Red-root 99, 114
Red-top 46
Reed, Bur 40
Relghard, J. E., Biological Examinations of Lake St. Clair 10
Reseda 84
Resedaceæ 84
Rhamnaceæ 98
Rhamnus 98
Rheumatism-root 80
Rhexia 102
Rhinanthus 121
Rhus 97
Rhynchospora 53
Ribes 86
Ribgrass 122
Ribwort 122
Rice, Indian 44
  Mountain 45
  Water 44
Richweed 71, 117
Ripplegrass 122
River-weed 84
River-weed Family 84
Robinia 92
Rocket, Sand 82
  Yellow 82
Rock-rose 100
Rock-rose Family 100
Rocky Mountain Bee-plant 84
Rominger, C., Geology of Lower Peninsula 12
Roripa 82
Rosa 88, 89
Rosaceæ 17, 86
Rosaceæ, Found on Pine Plains 17
Rose Family 86
Rose, Climbing 89
  Prairie 89
Rosemary, Wild 108
Rosin-weed 132
Rubiaceæ 123
Rubus 86, 87
Rudbeckia 132
Rue Family 95
  Goat's 92
Ruellia 122
Rumex 72
Rutaceæ 95
Rush Family 59
Rush 59
  Scouring 38

Page.
Rush, Soft 60
Spike 60
Wood 60
Russian Thistle 74
Rye, Wild 50

Sabbatia 110
Sage, Wood 115
Sagina 76
Sagittaria 42
Salicaceæ 67
Salix 68
Salmon-berry 87
Salomonia 62
Salsola 74
Sambucus 123
Samolus 109
Sandalwood Family 71
Sandwort 76
Thyme-leaved 76
Sanguinaria 80
Sanguisorba 88
Sanicle 105
Sanicula 105
Santalaceæ 71
Saponaria 75
Sarothra 100
Sarracenia 84, 104
Sarraceniaceæ 84
Sarsaparilla, Wild 104
Bristly 104
Sassafras 80
Satureia 117
Saururus 67
Savory, Summer 117
Saxifraga 85
Saxifragaceæ 85
Saxifrage 85
Early 85
Golden 85
Saxifrage Family 85
Scabious, Sweet 131
Scheuchzeriaceæ 42
Scheuchzeria 42
Schollera 109
Scirpus 52
Scleranthus 76
Scleria 53
Scoke 74
Scrub Pine, Gray or Northern 39
Scrophularia 119
Scrophulariaceæ 119
Scutellaria 115
Seal, Golden 77
Sea-Rocket 81
Secale 50
Sedge 53
Sedge Family 53
Sedum 85
Senna Family 91
Seed-box 103
Selaginella 38
Selaginellaceæ 38
Self-heal 116
Senna, Wild 91
Senecio 135
Service-berry 89
Setaria 44
Sevastana 44
Shad-bush 89
Sheep-berry 124
Shell-bark 67
Shepherdia 102
Shepherd's Purse 83
Shin-leaf 107
Shooting-Star 110
Shrubs, List of 30
Sibbaldiopsis 87
Sickle-pod 83
Sicyos 125
Sida 99
Side-saddle Flower 84
Sieversia 88

Page.
Silene 75
Silkweed 112
Silene 75
Silphium 132
Silver-weed 87
Simarubaceæ 95
Sinapis 81
Sisymbrium 81, 83
Sisyrinchium 63
Sium 106
Skeels, H. C., as Collector. 5
Skullcap 115
Mad-dog 115
Smartweed 72
Water 72
Smilacina 62
Smilaceæ 63
Smilax Family 63
Smilax 63
Smith, E. F., Catalogue of 3
Smith, E. F., Notes on Climate 12
Smith, E. F., Notes on Distribution of Plants 12
Snake-mouth 65
Snakeroot, Black 78
Button 105, 128
Seneca 96
Virginia 71
White 128
Snake-head 119
Sneeze-weed 124
Sneezewort 134
Snowberry 109, 124
Soapwort 75
Soil, Quality Indicated by Plants 16, 18
Solanaceæ 117, 118
Solanum 118
Solea 102
Solidago 129
Solomon's Seal 62
False 62
Great 62
Smaller 62
Sonchus 126
Sones, G. D., Herbarium of 11
Sophia 83
Sorbus 89
Sorghastrum 43
Sorrel 72
Dock 72
Sheep 72
Sour-Gum Tree 107
Southernwood 134
Spalding, V. M., Medicinal Plants 9
Sparganiaceæ 40
Sparganium 40
Spartina 47
Spathyema 58
Spearmint 117
Spearwort, Creeping 79
Specularia 125
Speedwell 120
Corn 120
Field 120
Marsh 120
Purslane 120
Thyme-leaved 120
Water 120
Spergula 76
Spergularia 76
Spice-bush 80
Spiderwort 59
Spiderwort Family 59
Spikenard 104
Spindle-tree 97
Spiræa 86
Spiranthes 65
Spirodela 58
Spleenwort 37
Sporobolus 45
Spring-Beauty 74
Spruce 39

Page.
Spruce, Black 39
White 39
Spurge 96
Spurge Family 96
Spurry 76
Corn 76
Squaw-root 122
Squaw-weed 135
Squirrel Corn 81
Stachys 116
Stacey, I. W., as Collector. 5
Staff-tree 97
Staff-tree Family 97
Staphylea 98
Staphyleaceæ 98
Star-flower 110
Star-grass 61, 63
Starwort 75
Steeple-Bush 86
Steironema 110
Stellaria 75
Stenophyllus 51
Stitchwort, Fleshy 75
Long-leaved 75
Long-stalked 75
Northern 75
Stickseed 114
Stick-tight 114
Stipa 45
St. John, C. E., List of Plants in Mason Co. 11
St. John's-wort 100
St. John's-wort Family 100
Canadian 100
Great 100
Kalm's 100
Marsh 100
Shrubby 100
Spotted 100
Stillwell, O. J., Herbarium of 11
Stone-crop 85
Ditch 85
Mossy 85
Stone-root 117
Storksbill 94
Stramonium, Common 118
Strawberry 87
Barren 88
Streptopus 62
Strophostyles 93
Stylophorum 80
Succory 126
Sugarberry 70
Sumach, Dwarf 97
Fragrant 97
Poison 97
Smooth 97
Staghorn 97
Sundew Family 84
Sundew, Oblong-leaved 84
Round-leaved 84
Slender 84
Sundrops 103
Sunflower 133
Swallow-wort, Black 112
Sweet-brier 89
Sweet Cicely 106
Sweet-fern 67
Sweet Flag 58
Sweet-Gale Family 67
Sweetbrier 89
Sweet William 75
Sycamore 86
Sylabum 136
Symphoricarpos 124
Symphytum 114
Symplocarpus 58
Syndesmon 78
Synosma 135
Syntherisma 43
Synthyris 120
Syringa 110

Page.

Tænidia 105
Tamarack 39
Tanacetum 134
Tansy 134
Taraxacum 126
Tare 94
Taxaceæ 40
Taxus 40
Tea, Labrador 108
Mexican 73
New Jersey 99
Oswego 116
Tear-thumb, Arrow-leaved. 72
Halberd-leaved 72
Teasel Family 125
Teasel 125
Tephrosia 92
Tetragonanthus 111
Teucrium 115
Thalesia 122
Thalictrum 79
Thaspium 105
Thimbleberry 87
Thistle, Bull 136
Canada 135
Hill's 135
Milk 136
Pasture 136
Pitcher's 136
Russian 74
Scotch 136
Sow 126
Spiny-leaved Sow 126
Swamp 136
Tall 135
Wavy-leaved 136
Yellow 136
Thlaspi 81
Thorn, Cockspur 90
Thorn-Apple 118
Thoroughwort 128
Thread-foot 84
Thuya 39
Thyme 117
Creeping 117
Thymelæaceæ 102
Thymus 117
Tiarella 85
Tickseed 133
Tiedemannia 105
Tilia 99
Tiliaceæ 99
Timothy 45
Tipularia 66
Tissa 76
Toad-flax 119
Bastard 71
Tobacco, Indian 126
Root 125
Wild 118
Tofieldia 60
Tomatillo 118
Toothache Tree 95
Toothwort 83
Topography 11
Touch-me-not 98
Tradescantia 59
Tragopogon 126
Tree-of-Heaven 95
Trees of Michigan 28
Trees, Distribution of, Explained by Dr. A. Gray.. 22
Trees, List of 28
Trees of Michigan Contrasted with those of Europe 21
Trees, Why has Great Britain so few? 22
Trees, Why has Michigan so many? 22
Trefoil 93, 95
Shrubby 93, 95
Tick 93
Tricuspis 47
Triadenum 100

Page.

Trientalis 110
Trifolium 92
Triglochin 42
Trillium 62
Dwarf White 62
Painted 62
Triodia 47
Triosteum 46, 124
Triphora 65
Triplasis 47
Trisetum 46
Trollius 77
Tsuga 39
Tulip-tree 77
Tumble Weed 74
Tupelo 107
Tussilago 135
Tuthill, F. H., Notes by.... 9
Turnip 82
Indian 58
Turtle-head 119
Twayblade 65, 66
Twin-flower 124
Twin-leaf 80
Twisted-stalk 62
Typha 40
Typhaceæ 40

Ulmaceæ 70
Ulmaria 88
Ulmus 70
Umbelliferæ 104
Umbel-wort 74
Unifolium 62
Urtica 71
Urticaceæ 71
Urticastrum 71
Utricularia 121
Uvularia 61

Vaccaria 75
Vaccinium 108, 109
Vagnera 62
Valerian 125
Valeriana 125
Valerianaceæ 125
Valerianella 125
Valerian Family 125
Valisneriaceæ 125
Vallisneria 42
Velvet-leaf 100
Verbascum 119
Verbena 114, 115
Verbenaceæ 114
Verbesina 133
Vernal-grass, Sweet 44
Vernonia 128
Veronica 120
Vervain 114, 115
Vervain Family 114
Vervain, Blue 114
White 114
Vetch 94
Cooper's 92
Milk 92
Viburnum 123, 124
Vicia 94
Vinca 111
Vincetoxicum 112
Vine Family 99
Viola 101
Violaceæ 101
Violet Family 101
Violet, Arrow-leaved 101
Bird-foot 101
Canada 101
Common Blue 101
Dame's 84
Dog's-tooth 61
Downy Yellow 101
Great-spurred 101
Green 101
Lance-leaved 101

Page.

Violet, Long-spurred 101
Pale 101
Round-leaved 101
Sweet White 101
White Dog's-tooth 61
Virgin's-Bower 78
Purple 79
Vitaceæ 99
Vitis 99

Waahoo 97
Waldsteinia 88
Walking-Leaf 37
Walnut 67
Walnut Family 67
Walnut, Black 67
Wake Robin 62
Washingtonia 106
Water-Dock, Great 72
Water-Plantain 42
Water-Plantain Family ... 42
Waterleaf 113
Waterleaf Family 113
Water-Hemp 74
Water-Lily Family 76
Water-Lily 77
Water-Milfoil 104
Water-Milfoil Family 104
Water-Pepper 72
Water-Shield 76
Water-Starwort 96
Wayfaring-tree, American. 123
Wax-work 97
Wax-Myrtle 67
Weeds from Europe 25
Native 26, 27
and introduced 25
Weymouth Pine 39
Wheat-Grass, Awned 49
Wheat-Thief 114
Wheeler, C. F., Cyperaceæ of Central Michigan 10
Wheeler, C. F., as Collector 4
Wheeler, C. F., Catalogue of 3
Herbarium of 11
White-rod 124
Whitewood 77
Whitney, W. D., Catalogue of 9
Wicky 108
Willow Family 67
Willow 68
Basket 69
Bebb's 68
Black 68
Dwarf Gray 69
Glaucous 68
Golden 68
Heart-leaved 68
Hoary 68
Myrtle 68
Prairie 68
Shining 68
Silky 68
Tomentose 68
Waten 122
Willow-herb 103
Great 103
Willugbaeya 128
Winchell, Alex., Climate, Notes on 13
Winchell, N. H., Catalogue of 9
Wind-flower 78
Wintergreen Family 107
Winterberry 97
Wintergreen, Chickweed .. 110
Creeping 108
Flowering 95
Spotted 107
Witch-Hazel 86
Witch-Hazel Family 86
Wolfberry 124

	Page.
Wolffia	58
Wood, F. E., Herbarium of	11
Woodsia	35
Wood-Sorrel	95
Woodwardia	37
Wormseed	73
Wormwood	127, 134
Wright, John, Catalogue of	9
Wulfenia	120
Xanthium	127
Xanthoxylum	95
Xolisma	108
Xyridaceæ	59
Xyris	59
Yam	63
Yam Family	63
Yam-root, Wild	63
Yarrow	134
Yellow-eyed-grass Family	59
Yellow-Rattle	121
Yew	40
Zannichellia	41
Zizania	44
Zizia	106
Zygadenus	60

ADDITIONS TO THE MICHIGAN FLORA.

W. J. BEAL.

[Reprinted from the Tenth Annual Report of the Michigan Academy of Science.]

ADDITIONS TO THE MICHIGAN FLORA AS PUBLISHED IN THE FIFTH REPORT OF THE MICHIGAN ACADEMY OF SCIENCE, 1904.

W. J. Beal.

The list includes 122 species and varieties of which H. S. Pepoon furnishes 66, collected in Counties Van Buren, Cass and Berrien.

These added to the former list make 2365.

The following seem to be the most interesting.

Andropogon Virginicum L.

Aristida tuberculosa Mutt.

Boltonia asteroides (L) L'Her.

Cassia Chamaecrista L.

Elatine Americana Arn. Growing about an inch high in mud, probably usually overlooked.

Quercus Michauxii Nutt.

Triosteum aurantiacum Bicknell.

Prunus Virginiana L. var. with cream-colored fruit of mild flavor. Collected at Menominee by A. L. Sawyer.

This variety was collected some years ago in Massachusetts by Mrs. H. L. T. Woolcott, Dedham, Mass.

Agrostis intermedia Scribn.

Upland Bent-grass. Cass Co., H. S. Pepoon.

Andropogon Virginicus L.

Virginia Beard-grass. Keeler, Van Buren Co., H. S. Pepoon.

Antennaria ambigens (Greene) Fernald.

Van Buren Co., H. S. Pepoon.

Antirrhinum majus L. Detroit, O. A. Farwell.

Arctium minus Schk.

Common Burdock, Keeler, Van Buren Co., H. S. Pepoon.

Aristida tuberculosa Nutt.

Sea Beach Aristida, Keeler, Van Buren Co., H. S. Pepoon.

Artemisia Pontica L.

Roman Wormwood, Bay City, Geo. M. Bradford.

Ascyrum hyperocoides L.

Near Detroit, O. A. Farwell.

Boltonia asteroides L. L'Her.

Aster-like Boltonia, Van Buren Co., H. S. Pepoon.

Briza media L.

Detroit River Front, O. A. Farwell.

Calamagrostis inexpansa A. Gray.

Bog Reed-grass, Keeler, Van Buren Co., H. S. Pepoon.

Campanula uliginosa Ryd.

Van Buren Co., H. S. Pepoon.

Cardamine Pennsylvanica Muhl.

Pennsylvania Bitter Cress. Keeler, Van Buren Co., H. S. Pepoon.

Carex Baileyi Brit.

Van Buren Co., H. S. Pepoon.

Carex conjuncta Boot.
Keeler, Van Buren Co., H. S. Pepoon.
Carex glaucodea Tuck.
Keeler, Van Buren Co., H. S. Pepoon.
Carex Jamesii Schwein.
Keeler, Van Buren Co., H. S. Pepoon.
Carex paupercula irrigua (Wahl.) Fernald.
Keweenaw Co., O. A. Farwell.
Carex paupercula pallens Fernald.
Keweenaw Co., O. A. Farwell.
Carex marcida Boot.
Thunder Bay Island, Port Huron, C. K. Dodge.
Carex teretiuscula prairea (Dewey). Britton.
Keeler, Van Buren Co., H. S. Pepoon.
Cassia Chamaecristata L.
Partridge Pea. Pipestone, Berrien Co., H. S. Pepoon.
Caucalis nodosa (L) Huds.
In Ballast, Detroit, O. A. Farwell.
Chelone obliqua L. Red Turtle head.
Near Huron River, Ann Arbor, S. Alexander.
Commelina communis L.
Asiatic Day-Flower, Dowagiac, Cass Co., H. S. Pepoon.
Corallorhiza multiflora flavida Peck. Large Coral-root.
Birmingham, S. Alexander.
Crataegus Margaretta Ash.
Dowagiac, H. S. Pepoon.
Cuscuta compacta Juss.
Compact Dodder. Keeler, Van Buren Co., H. S. Pepoon.
Cuscuta paradoxa Raf.
Glomerate Dodder. Berrien Co., H. S. Pepoon.
Dasystoma laevigata Raf.
Birmingham, O. A. Farwell.
Delphinium Ajacis L.
Keeler, Van Buren Co., H. S. Pepoon.
Dipsacus laciniatus L. Lansing, a mile and a half below the College, W. J. B. Cup-leaved Teasle.
Elatine Americana Arnott.
Mud Purslane, Border Round Lake, Van Buren Co., H. S. Pepoon.
Elatine brachisperma A. Gray.
Border of Lake Anne, Houghton Co., C. E. Davis.
Eleocharis melanocarpa Torr.
Priest's Lake, Cass Co., H. S. Pepoon.
Eleocharis palustris calva A. Gray.
Belle Isle, O. A. Farwell.
Equisetum palustre L.
Marsh Horsetail. Silver Creek, Van Buren Co., H. S. Pepoon.
Euphorbia serpyllifolia Pers.
Thyme-leaved Spurge, Agricultural College, escaped from cultivation W. J. B.
Euthamia Caroliniana (L) Greene. *Solidago tenuifolia* Pursh.
Slender Fragrant Golden-rod.
Dewey Lake, Cass Co. H. S. Pepoon.

Eupatorium trifoliatum L.
Crenate-leaved Joe-pye weed. Van Buren Co., H. S. Pepoon.
Galinsoga parviflora hispida D. C.
Detroit, O. A. Farwell.
Galium circaezans glabellum Britton.
Van Buren Co., H. S. Pepoon.
Galium Claytoni Michx.
Clayton's Bedstraw, Dewey Lake, Cass Co., H. S. Pepoon.
Galium tricorne Stokes.
In ballast, Detroit, O. A. Farwell.
Genista tinctoria L.
Detroit, O. A. Farwell.
Geranium Carolinianum L.
Carolina Crane's Bill. Cass Co., and Van Buren Co., H. S. Pepoon.
Helianthus annuus L.
Detroit, O. A. Farwell.
Helianthus atrorubens L.
Hairy-Wood Sunflower, Keeler, Van Buren Co., H. S. Pepoon.
Helianthus laetiflorus Pers.
Showy Sunflower, Keeler, Van Buren Co., H. S. Pepoon.
Helianthus lenticularis Dougl.
Ann Arbor, S. Alexander; Ypsilanti, O. A. Farwell.
Helianthus strumosus mollis T. & G.
Ypsilanti, O. A. Farwell; Ann Arbor, S. Alexander.
Heliopsis helianthoides (L.) B. S. P. H. *laevis* Pers, Addison, W. J. B.; Port Huron, C. K. Dodge.
Hemerocallis flava L.
Yellow Day-Lily. Dowagiac, Cass Co., H. S. Pepoon.
Hemicarpha micrantha (Vahl.) Brit.
Cable Lake, Van Buren Co., H. S. Pepoon.
Hicoria microcarpa (Nutt.) Brit.
Small-fruited Hickory. Keeler, Van Buren Co., H. S. Pepoon.
Hieracium marianum Willd.
Maryland Hawkweed. Keeler, Van Buren Co., H. S. Pepoon.
Hypoxis hirsuta (L.) Coville.
Star grass. Keeler, Van Buren Co., H. S. Pepoon.
Ilex Bronxensis Britton.
Cass Co., H. S. Pepoon.
Juncus marginatus aristatus (Michx.) Coville, Detroit, O. A. Farwell.
Kneiffia linearis (Michx.) Beach.
St. Clair Co., C. K. Dodge.
Lactuca Morssii Robinson. Morss Wild Lettuce.
Turin, U. P., Brunson Barlow.
Lechea racemulosa Michx.
Oblong-fruited Pin-weed, H. S. Pepoon.
Lycopus Virginicus L. Purple Bugle Weed. Near Magician Lake, Van Buren Co., H. S. Pepoon.
Mitchella repens alba.
Partridge-berry. Fruit white, Mrs. Helen W. Paul, Ontonagon.
Naias gracillima (A. Br.) Morong.
Thread-like Naias, H. S. Pepoon.
Onagra biensis cruciata (L.)
Belle Isle, Detroit. O. A. Farwell.

Onagra biennis grandiflora (Ait.) Lind. Detroit. O. A. Farwell.
Onagra biennis muricata L.
Keweenaw Co., O. A. Farwell.
Onagra biennis parviflora.
Detroit, O. A. Farwell.
Panicularia Torreyana.
Van Buren Co., H. S. Pepoon.
Panicum Ashei G. Pearson.
Ashe's Panicum (No specimen), H. S. Pepoon.
Panicum barbulatum Michx.
Keeler, Van Buren Co., H. S. Pepoon.
Panicum lanuginosum Ell. Van Buren Co., H. S. Pepoon.
Panicum laxiflorum Lam.
Lax-flowered Panicum (No specimen), H. S. Pepoon.
Panicum microcarpon Muhl.
Keeler, Van Buren Co., H. S. Pepoon.
Panicum pubescens Lam.
Pine Lake, Van Buren Co., H. S. Pepoon.
Parietaria Pennsylvanica Muhl.
Pennsylvania Pellitory. Keeler, Van Buren Co., H. S. Pepoon.
Polygala cruciata L.
Cross-leaved Milkwort. Sandy Shore, Dewey Lake, Cass Co., H. S. Pepoon.
Polygonum punctatum lepostachyum (Meisn.) Small.
Belle Isle, O. A. Farwell.
Potamogoton lonchitis L.
Detroit, O. A. Farwell.
Potamogeton nitens L.
Detroit, O. A. Farwell.
Prunus avium L.
Mazard Cherry, near Magician Lake, Cass Co., H. S. Pepoon.
Prunus cerasifera Ehrh.
Belle Isle, Detroit, O. A. Farwell.
Prunus Virginiana L.
Choke Cherry, Variety with cream colored fruit of mild flavor. A. L. Sawyer, Attorney, Menominee.
Quercus Michauxii Nutt.
Cow Oak, Basket Oak, Moist wood. N. Keeler, Van Buren Co., very rare, H. S. Pepoon.
Ranunculus Macounii Brit.
Macoun's Buttercup. Hartfort, Van Buren Co., H. S. Pepoon.
Rhynchospora corniculata (Lam.) A. Gray.
Horned Rush. Fox Lake, Van Buren Co., H. S. Pepoon.
Rhexia Virginica L.
Detroit, O. A. Farwell.
Rhus glabra borealis Britt.
Belle Isle, Detroit, O. A. Farwell.
Robinia hispida L.
Rose Acacia, Keeler, Van Buren Co., H. S. Pepoon.
Robinia viscosa Vent.
Clammy Locust. Near Silver Creek, Cass Co., H. S. Pepoon.
Rosa lucida Ehrh.
Glossy Rose. Van Buren Co., H. S. Pepoon.

Rotala ramosior (L) Koehne.
Van Buren Co., H. S. Pepoon.
Rubus Alleghaniensis Porter.
Keweenaw Co., O. A. Farwell.
Rudbeckia Sullivantii Boynton & Beadle.
North Lansing on River bank. Inserted in last catalogue as R. speciosa. W. J. B.
Salix eriocephala Michx.
Pussy Willow. Van Buren Co., H. S. Pepoon.
Salix fluviatilis Nutt.
Sand bar Willow. H. S. Pepoon.
Salix servissima (Bailey) Fernald.
Lakeville. Brotherton & Farwell.
Scandix Pecten-Veneris L.
In Ballast. Detroit, O. A. Farwell.
Scleria pauciflora Muhl.
Nut-rush. Keeler, Van Buren Co., H. S. Pepoon.
Sida spinosa Lin.
Detroit, O. A. Farwell.
Sisymbrium canescens Nutt.
Thunder Bay Island, C. F. Wheeler.
Smilax pulverulenta Michx.
Dowagiac, H. S. Pepoon.
Solidago juncea ramosa Porter.
Detroit, O. A. Farwell.
Solidago rigidiuscula (T. & G.) Porter.
Keeler, Van Buren Co., H. S. Pepoon.
Specularia perfoliata (L.) A. DC.
Venus' Looking Glass. Keeler, Van Buren Co., H. S. Pepoon.
Stenophragma Thaliana (L.) Celak. *Arabis Thaliana L.*
Mouse-ear Cress. H. S. Pepoon.
Symphoricarpos recemosus L.
Snow-berry. Silver Creek, Cass Co., H. S. Pepoon.
Taraxacum erythrospermum Andrez.
Previously only found in Wayne Co., Berrien Co., H. S. Pepoon.
Teucrium menthifolium Bicknell.
Detroit, O. A. Farwell.
Triosteum auriantiacum Bicknell.
Scarlet-fruited Horse-Gentian. Keeler, Van Buren Co., H. S. Pepoon.
Tunica Saxifraga Scop.
Escaped from cultivation, Mrs. Ralph Ballard, Niles.
Utricularia biflora Lam.
Two-flowered Bladder-wort. Silver Creek, Van Buren Co., H. S. Pepoon.
Vaccinium atrococcum (A. Gray) Heller.
Black Blueberry. Keeler, Van Buren Co., H. S. Pepoon.
Verbesina alternifolia (L.) Britt.
Ypsilanti, O. A. Farwell; Ann Arbor, S. Alexander.
Vernonia Illinoensis Gleason, *V. grandiflora*, O. A. Farwell.
Vernonia Noveboracensis (L.) Willd.
Detroit, O. A. Farwell.
Vibernum dentatum L.
Arrow-wood. N. Keeler, Van Buren Co., H. S. Pepoon.
Washingtonia divaricata Britt.
Thunder Bay Island, Alpena Co., C. K. Dodge.

Michigan Fl.a. 1904

BIOLOGY
LIBRARY
G

U.C. BERKELEY L
C026264

293907

Beal

QK167
B3
1904

BIOLOGY
LIBRARY
G

UNIVERSITY OF CALIFORNIA LIBRARY